

- 1 -

## Breast Cancer Progression Signatures

Inventors:

Xiao-Jun Ma

Dennis Sgroi

Mark G. Erlander

### FIELD OF THE INVENTION

The invention relates to the identification and use of gene expression profiles, or patterns, involved in breast cancer progression. The gene expression profiles, whether embodied in nucleic acid expression, protein expression, or other expression formats, are used in the study and/or diagnosis of cells and tissue during breast cancer progression as well as for the study and/or determination of prognosis of a patient. When used for diagnosis or prognosis, the profiles are used to predict the status and/or phenotype of cells and tissues relative to breast cancer and the treatment thereof.

### BACKGROUND OF THE INVENTION

Breast cancer is by far the most common cancer among women. Each year, more than 180,000 and 1 million women in the U.S. and worldwide, respectively, are diagnosed with breast cancer. Breast cancer is the leading cause of death for women between ages 50-55, and is the most common non-preventable malignancy in women in the Western Hemisphere. An estimated 2,167,000 women in the United States are currently living with the disease (National Cancer Institute, Surveillance Epidemiology and End Results (NCI SEER) program, *Cancer Statistics* sd-71385

- 2 -

Review (CSR), [www-seer.ims.nci.nih.gov/Publications/CSR1973](http://www-seer.ims.nci.nih.gov/Publications/CSR1973) (1998)). Based on cancer rates from 1995 through 1997, a report from the National Cancer Institute (NCI) estimates that about 1 in 8 women in the United States (approximately 12.8 percent) will develop breast cancer during her lifetime (NCI's Surveillance, Epidemiology, and End Results Program (SEER) publication *SEER Cancer Statistics Review 1973-1997*). Breast cancer is the second most common form of cancer, after skin cancer, among women in the United States. An estimated 250,100 new cases of breast cancer are expected to be diagnosed in the United States in 2001. Of these, 192,200 new cases of more advanced (invasive) breast cancer are expected to occur among women (an increase of 5% over last year), 46,400 new cases of early stage (*in situ*) breast cancer are expected to occur among women (up 9% from last year), and about 1,500 new cases of breast cancer are expected to be diagnosed in men (Cancer Facts & Figures 2001 American Cancer Society). An estimated 40,600 deaths (40,300 women, 400 men) from breast cancer are expected in 2001. Breast cancer ranks second only to lung cancer among causes of cancer deaths in women. Nearly 86% of women who are diagnosed with breast cancer are likely to still be alive five years later, though 24% of them will die of breast cancer after 10 years, and nearly half (47%) will die of breast cancer after 20 years.

Every woman is at risk for breast cancer. Over 70 percent of breast cancers occur in women who have no identifiable risk factors other than age (U.S. General Accounting Office. Breast Cancer, 1971-1991: Prevention, Treatment and Research. GAO/PEMD-92-12; 1991). Only 5 to 10% of breast cancers are linked to a family history of breast cancer (Henderson IC, Breast Cancer. In: Murphy GP, Lawrence WL, Lenhard RE (eds). *Clinical Oncology*. Atlanta, GA: American Cancer Society; 1995:198-219).

Each breast has 15 to 20 sections called lobes. Within each lobe are many smaller lobules. Lobules end in dozens of tiny bulbs that can produce milk. The lobes, lobules, and bulbs are all linked by thin tubes called ducts. These ducts lead to the nipple in the center of a dark area of skin called the areola. Fat surrounds the lobules and ducts. There are no muscles in the breast, but muscles lie under each breast and cover the ribs. Each breast also contains blood

- 3 -

vessels and lymph vessels. The lymph vessels carry colorless fluid called lymph, and lead to the lymph nodes. Clusters of lymph nodes are found near the breast in the axilla (under the arm), above the collarbone, and in the chest.

Breast tumors can be either benign or malignant. Benign tumors are not cancerous, they do not spread to other parts of the body, and are not a threat to life. They can usually be removed, and in most cases, do not come back. Malignant tumors are cancerous, and can invade and damage nearby tissues and organs. Malignant tumor cells may metastasize, entering the bloodstream or lymphatic system. When breast cancer cells metastasize outside the breast, they are often found in the lymph nodes under the arm (axillary lymph nodes). If the cancer has reached these nodes, it means that cancer cells may have spread to other lymph nodes or other organs, such as bones, liver, or lungs.

Major and intensive research has been focussed on early detection, treatment and prevention. This has included an emphasis on determining the presence of precancerous or cancerous ductal epithelial cells. These cells are analyzed, for example, for cell morphology, for protein markers, for nucleic acid markers, for chromosomal abnormalities, for biochemical markers, and for other characteristic changes that would signal the presence of cancerous or precancerous cells. This has led to various molecular alterations that have been reported in breast cancer, few of which have been well characterized in human clinical breast specimens. Molecular alterations include presence/absence of estrogen and progesterone steroid receptors, HER-2 expression/amplification (Mark HF, et al. HER-2/neu gene amplification in stages I-IV breast cancer detected by fluorescent in situ hybridization. Genet Med; 1(3):98-103 1999), Ki-67 (an antigen that is present in all stages of the cell cycle except G0 and used as a marker for tumor cell proliferation, and prognostic markers (including oncogenes, tumor suppressor genes, and angiogenesis markers) like p53, p27, Cathepsin D, pS2, multi-drug resistance (MDR) gene, and CD31.

Examination of cells by a trained pathologist has also been used to establish whether ductal epithelial cells are normal (i.e. not precancerous or cancerous or having another

- 4 -

noncancerous abnormality), precancerous (i.e. comprising hyperplasia, atypical ductal hyperplasia (ADH)) or cancerous (comprising ductal carcinoma *in situ*, or DCIS, which includes low grade ductal carcinoma *in situ*, or LG-DCIS, and high grade ductal carcinoma *in situ*, or HG-DCIS) or invasive (ductal) carcinoma (IDC). Pathologists may also identify the occurrence of lobular carcinoma *in situ* (LCIS) or invasive lobular carcinoma (ILC). Breast cancer progression may be viewed as the occurrence of abnormal cells, such as those of ADH, DCIS, IDC, LCIS, and/or ILC, among normal cells.

It remains unclear whether normal cells become hyperplastic (such as ADH) and then progressing on to become malignant (DCIS, IDC, LCIS, and/or ILC) or whether normal cells are able to directly become malignant without transitioning through a hyperplastic stage. It has been observed, however, that the presence of ADH indicates a higher likelihood of developing a malignancy. This has resulting in treatment of patients with ADH to begin treatment with an antineoplastic/antitumor agent such as tamoxifen. This is in contrast to the treatment of patients with malignant breast cancer which usually includes surgical removal.

The rational development of preventive, diagnostic and therapeutic strategies for women at risk for breast cancer would be aided by a molecular map of the tumorigenesis process. Relatively little is known of the molecular events that mediate the transition of normal breast cells to the various stages of breast cancer progression. In particular, there is a significant paucity of information regarding the genetic changes that are associated with the earliest stages of human breast cancer, which include the transition of normal breast cells to atypical hyperplastic and/or pre-invasive malignant cells (carcinoma *in situ*).

Molecular means of identifying the differences between normal, non-cancerous cells and cancerous cells (in general) have also been the focus of intense study. The use of cDNA libraries to analyze differences in gene expression patterns in normal versus tumorigenic cells has been described (USP 4,981,783). DeRisi et al. (1996) describe the analysis of gene expression patterns between two cell lines: UACC-903, which is a tumorigenic human melanoma cell line, and UACC-903(+6), which is a chromosome 6 suppressed non-tumorigenic form of UACC-903.

- 5 -

Labeled cDNA probes made from mRNA from these cell lines were applied to DNA microarrays containing 870 different cDNAs and controls. Genes that were preferentially expressed in one of the two cell lines were identified.

Golub et al. (1999) describe the use of gene expression monitoring as means to cancer class discovery and class prediction between acute myeloid leukemia (AML) and acute lymphoblastic leukemia (ALL). Their approach to class predictors used a neighborhood analysis followed by cross-validation of the validity of the predictors by withholding one sample and building a predictor based only on the remaining samples. This predictor is then used to predict the class of the withheld sample. They also used cluster analysis to identify new classes (or subtypes) within the AML and ALL.

Gene expression patterns in human breast cancers have been described by Perou et al. (1999), who studied gene expression between cultured human mammary epithelia cells (HMEC) and breast tissue samples by use of microarrays comprising about 5000 genes. They used a clustering algorithm to identify patterns of expression in HMEC and tissue samples. Perou et al. (2000) describe the use of clustered gene expression profiles to classify subtypes of human breast tumors. Hedenfalk et al. describe gene expression profiles in BRCA1 mutation positive, BRCA2 mutation positive, and sporadic tumors. Sgroi et al. also analyzed gene expression of normal and breast cancer cells from a single patient. Using gene expression patterns to distinguish breast tumor subclasses and predict clinical implications is described by Sorlie et al. and West et al.

All of the above described approaches, however, utilize heterogeneous populations of cells found in culture or in a biopsy to obtain information on gene expression patterns. The use of such populations may result in the inclusion or exclusion of multiple genes from the patterns. For this and the lack of statistical robustness reasons, the gene expression patterns observed by the above described approaches provide little confidence that the differences in gene expression may be meaningfully associated with the stages of breast cancer.

## SUMMARY OF THE INVENTION

The present invention relates to the identification and use of gene expression patterns (or profiles or “signatures”) which are correlated with (and thus able to discriminate between) cells in various stages of breast cancer. Broadly defined, these stages are non-malignant versus malignant, but may also be viewed as normal versus atypical (optionally including reactive and pre-neoplastic) versus cancerous. Another definition of the stages is normal versus precancerous (e.g. atypical ductal hyperplasia (ADH) or atypical lobular hyperplasia (ALH)) versus cancerous (e.g. carcinoma *in situ* such as DCIS and/or LCIS) versus invasive (e.g. carcinomas such as IDC and/or ILC). DCIS may be further viewed as low grade versus high grade or grade I through grade III.

The gene expression patterns comprise one or more than one gene capable of discriminating between various stages of breast cancer with significant accuracy. The gene(s) are identified as correlated with various stages of breast cancer such that the levels of their expression are relevant to a determination of the stage of breast cancer of a cell. Thus in one aspect, the invention provides a method to determine the stage of breast cancer of a subject afflicted with, or suspected of having, breast cancer by assaying a cell containing sample from said subject for expression of one or more than one gene disclosed herein as correlated with one or more stages of breast cancer.

Gene expression patterns of the invention are identified by analysis of gene expression in multiple samples of each stage to be studied. The overall gene expression profile of each sample is obtained by analyzing the expressed or unexpressed state of genes in each stage relative to each other (one gene to another across all genes). This overall profile is then analyzed to identify genes that are positively, or negatively, correlated, with a stage of breast cancer relative to other genes. An expression profile of a subset of human genes may then be identified by the methods of the present invention as correlated with breast cancer. The use of multiple samples increases the confidence which which a gene may be believed to be correlated with a particular stage.

- 7 -

Without sufficient confidence, it remains unpredictable whether a particular gene is actually correlated with a stage of breast cancer and also unpredictable whether a particular gene may be successfully used to identify the stage of an unknown breast cancer cell sample.

A profile of genes that are highly correlated with one stage relative to another may be used to assay a sample from a subject afflicted with, or suspected of having, breast cancer to identify the stage of breast cancer to which the sample belongs. Such an assay may be used as part of a method to determine the therapeutic treatment for said subject based upon the stage(s) of breast cancer identified.

The correlated genes may be used singly with significant accuracy or in combination to increase the ability to accurately discriminate between various stages of breast cancer. The present invention thus provides means for correlating a molecular expression phenotype with a physiological (cellular) stage or state. This correlation provides a way to molecularly diagnose and/or monitor a cell's status in comparison to different cancerous versus non-cancerous phenotypes as disclosed herein. Additional uses of the correlated gene(s) are in the classification of cells and tissues; determination of diagnosis and/or prognosis; and determination and/or alteration of therapy.

The ability to discriminate is conferred by the identification of expression of the individual genes as relevant and not by the form of the assay used to determine the actual level of expression. An assay may utilize any identifying feature of an identified individual gene as disclosed herein as long as the assay reflects, quantitatively or qualitatively, expression of the gene. Identifying features include, but are not limited to, unique nucleic acid sequences used to encode (DNA), or express (RNA), said gene or epitopes specific to, or activities of, a protein encoded by said gene. All that is required is the identity of the gene(s) necessary to discriminate between stages of breast cancer and an appropriate cell containing sample for use in an expression assay.

In one aspect, the invention provides for the identification of the gene expression patterns by analyzing global, or near global, gene expression from single cells or homogenous cell

- 8 -

populations which have been dissected away from, or otherwise isolated or purified from, contaminating cells beyond that possible by a simple biopsy. Because the expression of numerous genes fluctuate between cells from different patients as well as between cells from the same patient sample, multiple individual gene expression patterns are used as reference data to generate models which in turn permit the identification of individual gene(s) that are most highly correlated with particular breast cancer stages and/or have the best the ability to discriminate cells of one stage from another.

In another aspect, the invention provides physical and methodological means for detecting the expression of gene(s) identified by the models generated by individual expression patterns. These means may be directed to assaying one or more aspect of the DNA template(s) underlying the expression of the gene(s), of the RNA used as an intermediate to express the gene(s), or of the proteinaceous product expressed by the gene(s).

In a further aspect, the gene(s) identified by a model as capable of discriminating between breast cancer stages may be used to identify the cellular state of an unknown sample of cell(s) from the breast. Preferably, the sample is isolated via non-invasive means. The expression of said gene(s) in said unknown sample may be determined and compared to the expression of said gene(s) in reference data of gene expression patterns from the various stages of breast cancer. Optionally, the comparison to reference samples may be by comparison to the model(s) constructed based on the reference samples.

One advantage provided by the present invention is that contaminating, non-breast cells (such as infiltrating lymphocytes or other immune system cells) are not present to possibly affect the genes identified or the subsequent analysis of gene expression to identify the status of suspected breast cancer cells. Such contamination is present where a biopsy is used to generate gene expression profiles.

While the present invention has been described mainly in the context of human breast cancer, it may be practiced in the context of breast cancer of any animal known to be potentially afflicted by breast cancer. Preferred animals for the application of the present invention are



- 9 -

mammals, particularly those important to agricultural applications (such as, but not limited to, cattle, sheep, horses, and other “farm animals”) and for human companionship (such as, but not limited to, dogs and cats).

#### BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is a schematic representing a data matrix of a pair-wise comparison between Grade I and Grade III DCIS among 16 samples (across the top) and a large number of genes identified by “CloneID”) along the left hand side.

Figure 2 is a table showing the actual weight data corresponding to Example II, where the data from ten genes (by CloneID number vertically) are compared to DCIS and ADH samples (across the top). Some data in the table has been vertically presented to permit the table to be displayed on a single sheet. The use of “-“ with data in the table reflects genes that are more highly expressed in ADH relative to DCIS. The absence of “-“ reflects genes that are more highly expressed in DCIS relative to ADH.

Figure 3 is a table showing the actual weight data corresponding to Example VII, where the data from over 300 genes (by CloneID number vertically) are compared to DCIS and ADH samples (across the top). Some data in the table has been vertically presented solely for display purposes. The use of “-“ with data in the table reflects genes that are more highly expressed in ADH relative to DCIS. The absence of “-“ reflects genes that are more highly expressed in DCIS relative to ADH.

Figure 4 is a table showing the actual weight data corresponding to Example VIII, where the data from over 300 genes (by CloneID number vertically) are compared to samples (across the top) from two grades of DCIS. The use of “-“ with data in the table reflects genes that are more

- 10 -

highly expressed in grade I relative to grade III. The absence of “-“ reflects genes that are more highly expressed in grade III relative to grade I.

## DETAILED DESCRIPTION OF THE SPECIFIC EMBODIMENTS

Definitions of terms as used herein:

A gene expression “pattern” or “profile” or “signature” refers to the relative expression of a gene between two or more stages of breast cancer which is correlated with being able to distinguish between said stages.

A “gene” is a polynucleotide that encodes a discrete product, whether RNA or proteinaceous in nature. It is appreciated that more than one polynucleotide may be capable of encoding a discrete product. The term includes alleles and polymorphisms of a gene that encodes the same product, or a functionally associated (including gain, loss, or modulation of function) analog thereof, based upon chromosomal location and ability to recombine during normal mitosis.

A “stage” or “stages” (or equivalents thereof) of breast cancer refer to a physiologic state of a breast cell as defined by known cytological or histological (including immunohistology, histochemistry, and immunohistochemistry) procedures and are readily known to skilled in the art. Non-limiting examples include normal versus abnormal, non-cancerous versus cancerous, the different stages described herein (e.g. hyperplastic, carcinoma, and invasive), and grades within different stages (e.g. grades I, II, or III or the equivalents thereof within cancerous stages).

The terms “correlate” or “correlation” or equivalents thereof refer to an association between expression of one or more genes and a physiologic state of a breast cell to the exclusion of one or more other stages and/or identified by use of the methods as described herein. A gene may be expressed at higher or lower levels and still be correlated with one or more breast cancer stages.

- 11 -

A “polynucleotide” is a polymeric form of nucleotides of any length, either ribonucleotides or deoxyribonucleotides. This term refers only to the primary structure of the molecule. Thus, this term includes double- and single-stranded DNA and RNA. It also includes known types of modifications including labels known in the art, methylation, “caps”, substitution of one or more of the naturally occurring nucleotides with an analog, and internucleotide modifications such as uncharged linkages (e.g., phosphorothioates, phosphorodithioates, etc.), as well as unmodified forms of the polynucleotide.

The term “amplify” is used in the broad sense to mean creating an amplification product can be made enzymatically with DNA or RNA polymerases. “Amplification,” as used herein, generally refers to the process of producing multiple copies of a desired sequence, particularly those of a sample. “Multiple copies” mean at least 2 copies. A “copy” does not necessarily mean perfect sequence complementarity or identity to the template sequence.

By corresponding is meant that a nucleic acid molecule shares a substantial amount of sequence identity with another nucleic acid molecule. Substantial amount means at least 95%, usually at least 98% and more usually at least 99%, and sequence identity is determined using the BLAST algorithm, as described in Altschul et al. (1990), J. Mol. Biol. 215:403-410 (using the published default setting, i.e. parameters w=4, t=17). Methods for amplifying mRNA are generally known in the art, and include reverse transcription PCR (RT-PCR) and those described in U.S. Patent Application (number to be assigned) entitled “Nucleic Acid Amplification” filed on October 25, 2001 as attorney docket number 485772002900 as well as U.S. Provisional Patent Applications 60/298,847 (filed June 15, 2001) and 60/257,801 (filed December 22, 2000), all of which are hereby incorporated by reference in their entireties as if fully set forth. Alternatively, RNA may be directly labeled as the corresponding cDNA by methods known in the art.

A “microarray” is a linear or two-dimensional array of preferably discrete regions, each having a defined area, formed on the surface of a solid support such as, but not limited to, glass, plastic, or synthetic membrane. The density of the discrete regions on a microarray is

- 12 -

determined by the total numbers of immobilized polynucleotides to be detected on the surface of a single solid phase support, preferably at least about 50/cm<sup>2</sup>, more preferably at least about 100/cm<sup>2</sup>, even more preferably at least about 500/cm<sup>2</sup>, but preferably below about 1,000/cm<sup>2</sup>. Preferably, the arrays contain less than about 500, about 1000, about 1500, about 2000, about 2500, or about 3000 immobilized polynucleotides in total. As used herein, a DNA microarray is an array of oligonucleotides or polynucleotides placed on a chip or other surfaces used to hybridize to amplified or cloned polynucleotides from a sample. Since the position of each particular group of primers in the array is known, the identities of a sample polynucleotides can be determined based on their binding to a particular position in the microarray.

Because the invention relies upon the identification of genes that are over- or under-expressed, one embodiment of the invention involves determining expression by hybridization of mRNA, or an amplified or cloned version thereof, of a sample cell to a polynucleotide that is unique to a particular gene sequence. Preferred polynucleotides of this type contain at least about 20, at least about 22, at least about 24, at least about 26, at least about 28, at least about 30, or at least about 32 consecutive basepairs of a gene sequence that is not found in other gene sequences. The term "about" as used in the previous sentence refers to an increase or decrease of 1 from the stated numerical value. Even more preferred are polynucleotides of at least about 50, at least about 100, and at least about 150 basepairs of a gene sequence that is not found in other gene sequences. The term "about" as used in the preceding sentence refers to an increase or decrease of 10% from the stated numerical value.

Alternatively, and in another embodiment of the invention, gene expression may be determined by analysis of expressed protein in a cell sample of interest by use of one or more antibodies specific for one or more epitopes of individual gene products (proteins) in said cell sample. Such antibodies are preferably labeled to permit their easy detection after binding to the gene product.

The term "label" refers to a composition capable of producing a detectable signal indicative of the presence of the labeled molecule. Suitable labels include radioisotopes,

- 13 -

nucleotide chromophores, enzymes, substrates, fluorescent molecules, chemiluminescent moieties, magnetic particles, bioluminescent moieties, and the like. As such, a label is any composition detectable by spectroscopic, photochemical, biochemical, immunochemical, electrical, optical or chemical means.

The term “support” refers to conventional supports such as beads, particles, dipsticks, fibers, filters, membranes and silane or silicate supports such as glass slides.

As used herein, a “breast tissue sample” or “breast cell sample” refers to a sample of breast tissue or fluid isolated from an individual suspected of being afflicted with, or at risk of developing, breast cancer. Such samples are primary isolates (in contrast to cultured cells) and may be collected by any non-invasive means, including, but not limited to, ductal lavage, fine needle aspiration, needle biopsy, the devices and methods described in U.S. Patent 6,328,709, or any other suitable means recognized in the art. Alternatively, the “sample” may be collected by an invasive method, including, but not limited to, surgical biopsy.

“Expression” and “gene expression” include transcription and/or translation of nucleic acid material.

As used herein, the term “comprising” and its cognates are used in their inclusive sense; that is, equivalent to the term “including” and its corresponding cognates.

Conditions that “allow” an event to occur or conditions that are “suitable” for an event to occur, such as hybridization, strand extension, and the like, or “suitable” conditions are conditions that do not prevent such events from occurring. Thus, these conditions permit, enhance, facilitate, and/or are conducive to the event. Such conditions, known in the art and described herein, depend upon, for example, the nature of the nucleotide sequence, temperature, and buffer conditions. These conditions also depend on what event is desired, such as hybridization, cleavage, strand extension or transcription.

Sequence “mutation,” as used herein, refers to any sequence alteration in the sequence of a gene disclosed herein interest in comparison to a reference sequence. A sequence mutation includes single nucleotide changes, or alterations of more than one nucleotide in a sequence, due

- 14 -

to mechanisms such as substitution, deletion or insertion. Single nucleotide polymorphism (SNP) is also a sequence mutation as used herein. Because the present invention is based on the relative level of gene expression, mutations in non-coding regions of genes as disclosed herein may also be assayed in the practice of the invention.

“Detection” includes any means of detecting, including direct and indirect detection of gene expression and changes therein. For example, “detectably less” products may be observed directly or indirectly, and the term indicates any reduction (including the absence of detectable signal). Similarly, “detectably more” product means any increase, whether observed directly or indirectly.

Unless defined otherwise all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention belongs.

### Specific Embodiments

The present invention relates to the identification and use of gene expression patterns (or profiles or “signatures”) which discriminate between (or are correlated with) cells in various stages of breast cancer. Such patterns may be determined by the methods of the invention by use of a number of reference cell or tissue samples, such as those reviewed by a pathologist of ordinary skill in the pathology of breast cancer, which reflect various stages of breast cancer. Because the overall gene expression profile differs from person to person, cancer to cancer, and cancer cell to cancer cell, correlations between certain cell states and genes expressed or underexpressed may be made as disclosed herein to identify genes that are capable of discriminating between different breast cancer states.

The present invention may be practiced with any number of genes believed, or likely to be, differentially expressed in breast cancer cells. In Example I below, approximately 12,000 genes were used to identify hundreds of genes capable of discriminating between various stages of breast cancer as shown in Examples 2-9. The identification may be made by using expression

- 15 -

profiles of various homogenous normal and breast cancer cell populations, which were isolated by microdissection, such as, but not limited to, laser capture microdissection (LCM) of 100-1000 cells. Each gene of the expression profile may be assigned weights based on its ability to discriminate between two or more stages of breast cancer (see Example I). The magnitude of each assigned weight indicates the extent of difference in expression between the two groups and is an approximation of the ability of expression of the gene to discriminate between the two groups (and thus stages). The magnitude of each assigned weight also approximates the extent of correlation between expression of individual gene(s) and particular breast cancer stages.

It should be noted that merely high levels of expression in cells from a particular stage or stages does not necessarily mean that a gene will be identified as having a high absolute weight value.

Genes with top ranking weights (in absolute terms) may be used to generate models of gene expressions that would maximally discriminate between the two groups. Alternatively, genes with top ranking weights (in absolute terms) may be used in combination with genes with lower weights without significant loss of ability to discriminate between groups. Such models may be generated by any appropriate means recognized in the art, including, but not limited to, cluster analysis, supported vector machines, neural networks or other algorithm known in the art. The models are capable of predicting the classification of a unknown sample based upon the expression of the genes used for discrimination in the models. "Leave one out" cross-validation may be used to test the performance of various models and to help identify weights (genes) that are uninformative or detrimental to the predictive ability of the models. Cross-validation may also be used to identify genes that enhance the predictive ability of the models.

The gene(s) identified as correlated with particular breast cancer stages by the above models provide the ability to focus gene expression analysis to only those genes that contribute to the ability to identify a cell as being in a particular stage of breast cancer relative to another stage or stages. The expression of other genes in a breast cancer cell would be relatively unable to provide information concerning, and thus assist in the discrimination of, different stages of

- 16 -

breast cancer. For example, the alpha subunit of human topoisomerase II (identified by CloneID 825470) has been found to be useful in discriminations between normal and atypical cells (ADH and DCIS and IDC and LCIS), between normal and ADH cells compared to DCIS and IDC cells, between normal and DCIS cells, between ADH and DCIS cells, between grade I and grade III DCIS cells, and between grade I and grade III IDC cells but not between normal and ADH cells (see Examples II to IX below). Thus expression of this topoisomerase II subunit would be utilized in models to discriminate between the above listed stages but not for discerning normal from ADH cells. This type of analysis is readily incorporated into algorithms used to generate models with reference gene expression data.

As will be appreciated by those skilled in the art, the models are highly useful with even a small set of reference gene expression data and can become increasingly accurate with the inclusion of more reference data although the incremental increase in accuracy will likely diminish with each additional datum. The preparation of additional reference gene expression data using genes identified and disclosed herein for discriminating between different stages of breast cancer is routine and may be readily performed by the skilled artisan to permit the generation of models as described above to predict the status of an unknown sample based upon the expression levels of those genes.

To determine the expression levels of genes in the practice of the present invention, any method known in the art may be utilized. In one preferred embodiment of the invention, expression based on detection of RNA which hybridizes to the genes identified and disclosed herein is used. This is readily performed by any RNA detection or amplification+detection method known or recognized as equivalent in the art such as, but not limited to, reverse transcription-PCR, the methods disclosed in U.S. Patent Application (number to be assigned) entitled "Nucleic Acid Amplification" filed on October 25, 2001 as attorney docket number 485772002900 as well as U.S. Provisional Patent Applications 60/298,847 (filed June 15, 2001) and 60/257,801 (filed December 22, 2000), and methods to detect the presence, or absence, of RNA stabilizing or destabilizing sequences.



- 17 -

Alternatively, expression based on detection of DNA status may be used. Detection of the DNA of an identified gene as methylated or deleted may be used for genes that have decreased expression in correlation with a particular breast cancer stage. This may be readily performed by PCR based methods known in the art. Conversely, detection of the DNA of an identified gene as amplified may be used for genes that have increased expression in correlation with a particular breast cancer stage. This may be readily performed by PCR based, fluorescent in situ hybridization (FISH) and chromosome in situ hybridization (CISH) methods known in the art.

Expression based on detection of a presence, increase, or decrease in protein levels or activity may also be used. Detection may be performed by any immunohistochemistry (IHC) based, blood based (especially for secreted proteins), antibody (including autoantibodies against the protein) based, exfoliate cell (from the cancer) based, mass spectroscopy based, and image (including use of labeled ligand) based method known in the art and recognized as appropriate for the detection of the protein. Antibody and image based methods are additionally useful for the localization of tumors after determination of cancer by use of cells obtained by a non-invasive procedure (such as ductal lavage or fine needle aspiration), where the source of the cancerous cells is not known. A labeled antibody or ligand may be used to localize the carcinoma(s) within a patient.

A preferred embodiment using a nucleic acid based assay to determine expression is by immobilization of one or more of the genes identified herein on a solid support, including, but not limited to, a solid substrate as an array or to beads or bead based technology as known in the art. Alternatively, solution based expression assays known in the art may also be used. The immobilized gene(s) may be in the form of polynucleotides that are unique or otherwise specific to the gene(s) such that the polynucleotide would be capable of hybridizing to a DNA or RNA corresponding to the gene(s). These polynucleotides may be the full length of the gene(s) or be short sequences of the genes that are optionally minimally interrupted (such as by mismatches or

- 18 -

inserted non-complementary basespairs) such that hybridization with a DNA or RNA corresponding to the gene(s) is not affected.

The immobilized gene(s) may be used to determine the state of nucleic acid samples prepared from sample breast cell(s) for which the pre-cancer or cancer status is not known or for confirmation of a status that is already assigned to the sample breast cell(s). Without limiting the invention, such a cell may be from a patient suspected of being afflicted with, or at risk of developing, breast cancer. The immobilized polynucleotide(s) need only be sufficient to specifically hybridize to the corresponding nucleic acid molecules derived from the sample. While even a single correlated gene sequence may be able to provide adequate accuracy in discriminating between two breast cancer cell stages, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, or eleven or more of the genes identified herein may be used as a subset capable of discriminating may be used in combination to increase the accuracy of the method. The invention specifically contemplates the selection of more than one, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, or eleven or more of the genes disclosed in the tables and figures herein for use as a subset in the identification of whether an unknown or suspicious breast cancer sample is normal or is in one or more stages of breast cancer. Optionally, the genes used will not include CloneID 809507, which is also known as GenBank accession number AA454563, described as an EST with high similarity to CD63 but of unknown function.

In embodiments where only one or a few genes are to be analyzed, the nucleic acid derived from the sample breast cancer cell(s) may be preferentially amplified by use of appropriate primers such that only the genes to be analyzed are amplified to reduce contaminating background signals from other genes expressed in the breast cell. Alternatively, and where multiple genes are to be analyzed or where very few cells (or one cell) is used, the nucleic acid from the sample may be globally amplified before hybridization to the immobilized

- 19 -

polynucleotides. Of course RNA, or the cDNA counterpart thereof may be directly labeled and used, without amplification, by methods known in the art.

The above assay embodiments may be used in a number of different ways to identify or detect the breast cancer stage, if any, of a breast cancer cell sample from a patient. In many cases, this would reflect a secondary screen for the patient, who may have already undergone mammography or physical exam as a primary screen. If positive, the subsequent needle biopsy, ductal lavage, fine needle aspiration, or other analogous methods may provide the sample for use in the above assay embodiments. The present invention is particularly useful in combination with non-invasive protocols, such as ductal lavage or fine needle aspiration, to prepare a breast cell sample. The current analysis of ductal lavage samples is by cytological examination by a trained pathologist who classifies the samples in terms that are at least partly subjective: unsatisfactory (too few cells), benign (including fibrocystic change), atypical (or mild atypia), suspicious (or marked atypia), or malignant.

The present invention provides a more objective set of criteria, in the form of gene expression profiles of a discrete set of genes, to discriminate (or delineate) between meaningful stages (or classes) of breast cancer cells. In particularly preferred embodiments of the invention, the assays are used to discriminate between ADH and DCIS or otherwise malignant cells, which is a critical determination for decisions concerning subsequent treatment and therapy for the patient. Another particularly preferred determination is between the three grades (I, II, III) of carcinomas *in situ* as well as the discrimination between grade III carcinomas *in situ* and invasive carcinomas. Other pairwise comparisons that are provided by the invention include, but are not limited to, normal versus ADH, normal versus carcinoma *in situ*, normal versus invasive, normal versus cancerous (i.e. carcinoma present), ADH versus cancerous, and carcinoma *in situ* versus invasive. With the use of alternative algorithms, such as neural networks, comparisons that discriminate between multiple (more than pairwise) classes may also be performed. It is believed by the inventors that the present invention is the first example of objective, molecular criteria for making these discriminations.

- 20 -

In one embodiment of the invention, the isolation and analysis of a breast cancer cell sample may be performed as follows:

- (1) Ductal lavage or other non-invasive procedure is performed on a patient to obtain a sample.
- (2) Sample is prepared and coated onto a microscope slide. Note that ductal lavage results in clusters of cells that are cytologically examined as stated above.
- (3) Pathologist or image analysis software scans the sample for the presence of atypical cells.
- (4) If atypical cells are observed, those cells are harvested (e.g. by microdissection such as LCM).
- (5) RNA is extracted from the harvested cells.
- (6) RNA is purified, amplified, and labeled.
- (7) Labeled nucleic acid is contacted with a microarray containing polynucleotides of the genes identified herein as correlated to discriminations between two or more stages of breast cancer under hybridization conditions, then processed and scanned to obtain a pattern of intensities of each spot (relative to a control for general gene expression in cells) which determine the level of expression of the gene(s) in the cells.
- (8) The pattern of intensities is analyzed by comparison to the expression patterns of the genes in known samples of normal and breast cancer cells (relative to the same control).

A specific example of the above method would be performing ductal lavage following a primary screen, observing and collecting atypical cells for analysis. The comparison to known expression patterns, such as that made possible by a model generated by an algorithm (such as, but not limited to nearest neighbor type analysis, SVM, or neural networks) with reference gene expression data for the different breast cancer stages, identifies the cells as being most likely ADH.

Alternatively, the sample may permit the collection of both normal and atypical cells for analysis. The gene expression patterns for each of these two samples will be compared to each

- 21 -

other as well as the model and the normal versus individual abnormal comparisons therein based upon the reference data set. This approach can be significantly more powerful than the atypical cells only approach because it utilizes significantly more information from the normal cells and the differences between normal and atypical cells (in both the sample and reference data sets) to determine the status of the atypical cells from the sample.

By appropriate selection of the genes used in the analysis, identification of the relative amounts of atypical cells may also be possible, although in most clinical settings, the identification of the highest grade of breast cancer with confidence makes identification of lower grades less important. Stated differently, the identification of invasive cancer determines the clinical situation regardless of the presence of carcinoma *in situ* or hyperplastic cells, or the identification of carcinoma *in situ* makes determines the clinical situation regardless of the presence of hyperplastic cells.

With use of the present invention, skilled physicians may prescribe treatments based on non-invasive samples that they would have prescribed for a patient which had previously received a diagnosis via a solid tissue biopsy.

The above discussion is also applicable where a palpable lesion is detected followed by fine needle aspiration or needle biopsy of cells from the breast. The cells are plated and reviewed by a pathologist or automated imaging system which selects cells for analysis as described above. This again provides a means of linking visual to molecular cytology and provides a less subjective means of identifying the physiological state of breast cancer cells without the need for invasive solid tissue biopsies.

The present invention may also be used, however, with solid tissue biopsies. For example, a solid biopsy may be collected and prepared for visualization followed by determination of expression of one or more genes identified herein to determine the stage of breast cancer, if any. One preferred means is by use of *in situ* hybridization with polynucleotide or protein identifying probe(s) for assaying expression of said gene(s).

- 22 -

In an alternative method, the solid tissue biopsy may be used to extract molecules followed by analysis for expression of one or more gene(s). This provides the possibility of leaving out the need for visualization and collection of only those cells suspected of being atypical. This method may of course be modified such that only cells suspected of being atypical are collected and used to extract molecules for analysis. This would require visualization and selection as an prerequisite to gene expression analysis.

In a further modification of the above, both normal cells and cells suspected of being atypical are collected and used to extract molecules for analysis of gene expression. The approach, benefits and results are as described above using non-invasive sampling.

In a further alternative to all of the above, the gene(s) identified herein may be used as part of a simple PCR or array based assay simply to determine the presence of atypical cells in a sample from a non-invasive sampling procedure. This is simple to perform and utilizes genes identified to be the best discriminators of normal versus abnormal cells without the need for any cytological examination. If no atypical cells are identified, no cytological examination is necessary. If atypical cells are identified, cytological examination follows, and a more comprehensive analysis, as described above, may follow.

The genes identified herein may be used to generate a model capable of predicting the breast cancer stage (if any) of an unknown breast cell sample based on the expression of the identified genes in the sample. Such a model may be generated by any of the algorithms described herein or otherwise known in the art as well as those recognized as equivalent in the art using gene(s) (and subsets thereof) disclosed herein for the identification of whether an unknown or suspicious breast cancer sample is normal or is in one or more stages of breast cancer. The model provides a means for comparing expression profiles of gene(s) of the subset from the sample against the profiles of reference data used to build the model. The model can compare the sample profile against each of the reference profiles or against model defining delineations made based upon the reference profiles. Additionally, relative values from the sample profile may be used in comparison with the model or reference profiles.

- 23 -

In a preferred embodiment of the invention, breast cell samples identified as normal and abnormal (atypical) from the same subject may be analyzed for their expression profiles of the genes used to generate the model. This provides an advantageous means of identifying the stage of the abnormal sample based on relative differences from the expression profile of the normal sample. These differences can then be used in comparison to differences between normal and individual abnormal reference data which was also used to generate the model.

The detection of gene expression from the samples may be by use of a single microarray able to assay gene expression from all pairwise comparisons disclosed herein for convenience and accuracy.

Other uses of the present invention include providing the ability to identify breast cancer cell samples as being those of a particular stage of cancer for further research or study. This provides a particular advantage in many contexts requiring the identification of breast cancer stage based on objective genetic or molecular criteria rather than cytological observation. It is of particular utility to distinguish different grades of a particular breast cancer stage for further study, research or characterization because no objective criteria for such delineation was previously available.

The materials for use in the methods of the present invention are ideally suited for preparation of kits produced in accordance with well known procedures. The invention thus provides kits comprising agents for the detection of expression of the disclosed genes for identifying breast cancer stage. Such kits optionally comprising the agent with an identifying description or label or instructions relating to their use in the methods of the present invention, is provided. Such a kit may comprise containers, each with one or more of the various reagents (typically in concentrated form) utilized in the methods, including, for example, pre-fabricated microarrays, buffers, the appropriate nucleotide triphosphates (e.g., dATP, dCTP, dGTP and dTTP; or rATP, rCTP, rGTP and UTP), reverse transcriptase, DNA polymerase, RNA polymerase, and one or more primer complexes of the present invention (e.g., appropriate length

- 24 -

poly(T) or random primers linked to a promoter reactive with the RNA polymerase). A set of instructions will also typically be included.

The methods provided by the present invention may also be automated in whole or in part. All aspects of the present invention may also be practiced such that they consist essentially of a subset of the disclosed genes to the exclusion of material irrelevant to the identification of breast cancer stages in a cell containing sample.

Having now generally described the invention, the same will be more readily understood through reference to the following examples which are provided by way of illustration, and are not intended to be limiting of the present invention, unless specified.

## EXAMPLES

### Example I: Materials and Methods

#### Clinical specimens

Clinical biopsies from 30 patients were obtained from the Massachusetts General Hospital with Institutional Review Board approval. The tissue from one of the patients was not associated with breast cancer of any kind since it was from a breast reduction procedure. Pathological and histological information for the biopsies were also obtained. Three independent captures of about 1000 breast epithelial cells of one or more of the four different disease stages (normal, N; atypical ductal hyperplasia, ADH; ductal carcinoma *in situ*, DCIS; invasive ductal carcinoma, IDC) were procured from each biopsy using Laser Capture Microdissection (LCM, Arcturus Engineering). Three independent captures of LCIS (lobular carcinoma in situ) in one biopsy were also made. Total RNA was extracted from the captured (procured) cells and amplified with a T7-promoter based RNA amplification protocol. The human universal



- 25 -

reference RNA (Stratagene, La Jolla), was similarly amplified and used as the reference channel in a two-color microarray hybridization.

#### Microarrays

To maximize coverage of breast cancer-related genes on the microarrays used, 11,435 cDNA clones from the IMAGE consortium (Research Genetics) were obtained. These clones were selected based on literature knowledge (such as, but not limited to, preferential expression in cancer versus normal cells) and after mining (such as, but not limited to, preferential expression in breast tissues) gene expression information in the expressed sequence tags (EST) databases and the Serial Analysis of Gene Expression (SAGE) data sets available from the National Center for Biotechnology Information (NCBI, <http://www.ncbi.nlm.nih.gov>).

#### Microarray data processing

Microarray images were analyzed with ImaGene (BioDiscovery) to find and quantitate each spot on the microarray. Spots flagged by ImaGene as poor spots using standard criteria used with the software for the standardization of signals were excluded from further analysis. Raw Cy5 (sample channel) and Cy3 (reference channel) intensities and associated local background estimates for each spot were then examined. The signal/noise ratio, defined as the spot intensity over background intensity, was used as the second criteria for spot exclusion; spots with signal/noise ratio < 3.0 in the reference channel or < 1.5 in the sample channel were excluded from further analysis. Background-subtracted intensities across the chip were normalized to the 75th-percentile of the spot intensity distribution on the entire chip (alternative normalizations to the mean, median or other point may also be used as known in the art). Cy5/Cy3 ratios of each spot for each cellular state were averaged across each of six measurements (3 LCM captures x 2 chips/capture = 6 chips); outliers among the 6 data points were removed before taking the average. The resulting data were formatted as a data matrix

- 26 -

(samples along the top horizontal axis and gene identity along the vertical axis) for data mining (see Figure 1 with data).

#### Microarray data analysis

Before further analysis, each value in a row (gene) of the gene expression matrix was divided by the median value for the row, and the resulting matrix log-transformed. Normalized, median-centered, and log-transformed, gene expression data matrix was loaded into GeneMaths software (Applied-Maths, Belgium). Clustering and discriminant analysis were performed to identify sets of genes associated with different cellular states. For each pair-wise comparison between two breast cancer stages, samples are assigned to either the positive group or negative group, and genes were sorted by their discriminatory weights. The absolute value of the weight of a gene indicates the extent of difference in expression between the two groups; the positively signed genes are expressed higher in one group and the negatively signed genes are expressed higher in the other group.

The utility of the top-ranking genes as a diagnostic test was evaluated using the support vector machines (SVMs) algorithm (see Yeang, C. H., S. Ramaswamy, et al. (2001). "Molecular classification of multiple tumor types." *Bioinformatics* 17 Suppl 1: S316-22; Xiong, M., X. Fang, et al. (2001). "Biomarker identification by feature wrappers." *Genome Res* 11(11): 1878-87 **this one used linear discriminate analysis, logistic regression and svm**; Furey, T. S., N. Cristianini, et al. (2000). "Support vector machine classification and validation of cancer tissue samples using microarray expression data." *Bioinformatics* 16(10): 906-14; and Brown, M. P., W. N. Grundy, et al. (2000). "Knowledge-based analysis of microarray gene expression data by using support vector machines." *Proc Natl Acad Sci U S A* 97(1): 262-7, who state "SVMs are considered a supervised computer learning method because they exploit prior knowledge of gene function to identify unknown genes of similar function from expression data. SVMs avoid several problems associated with unsupervised clustering methods, such as hierarchical

- 27 -

clustering and self-organizing maps.”) Other algorithms, such as, but not limited to, linear discriminate analysis, logistic regression, cluster analysis, K-th nearest neighbor, or neural nets.

The support vector machines algorithm finds the maximal margin hyperplane that separate the two groups under comparison. The method of leave-one-out cross-validation was used to test the performance of a given set of genes; one sample was taken in turn out of the training set and a model is built using the rest of the training set, which is in then applied to classify the left-out sample. The accuracy of the genes in the cross-validation procedure is the percentage of correct classifications over the total number of the training samples.

#### Example II: 10 Genes for discriminating between ADH and DCIS

Based upon the methodology of Example I above, 10 genes identified as capable of discriminating between ADH and DCIS are listed in Table 1 below along with a brief description of the gene. CloneID as used in the context of the present invention refers to the IMAGE Consortium clone ID number of each gene, the sequences of which are hereby incorporated by reference in their entirety as they are available from the Consortium at <http://image.llnl.gov/> as accessed on the filing date of the present application. Weight refers to the absolute value indicating the extent of difference in expression between ADH and DCIS where the positively signed values are expressed higher in ADH and the negatively signed values are expressed higher in DCIS; Chromosome Location refers to the human chromosome to which the gene has been assigned, and Description provides a brief identifier of what the gene encodes. The actual data corresponding to the assigned weights are shown in Figure 2.

**Table 1**

GeneID	Weight	Chromosome Location	Description
825470	0.9946555	17q21-q22	topoisomerase (DNA) II alpha (170kD)
595213	0.9884884	8	hypothetical protein
796694	0.9852686	17q25	baculoviral IAP repeat-containing 5 (survivin)
1404774	-1.5638738	12p12.1-p11.2	parathyroid hormone-like hormone
823871	-1.5260464	7	SPARC-like 1 (mast9, hevin)
1882697	-1.401878	17q22-q23	peanut (Drosophila)-like 2
140071	-1.3881954	2	frizzled-related protein
160192	-1.2551663	5	ESTs, Weakly similar to 2004399A
796542	-1.1401853	3q28	chromosomal protein [H.sapiens]
611532	-1.1227597	11p15.5	ets variant gene 5 (ets-related molecule)
			troponin I, skeletal, fast

Example III: Genes for discriminating between normal and non-normal (a combination of ADH, DCIS, and IDC) cells from breast tissue

As shown in Table 2 below, 850 genes were identified as being able to discriminate between normal and "abnormal" (defined in this instance as any sample that was not normal), which includes ADH, DCIS and IDC.

**Table 2**

CloneID	Weight	Description
1323448	1.5470535	cysteine-rich protein 1 (intestinal)
788654	1.4818381	growth factor receptor-bound protein 2
35147	1.3764654	"ESTs, Weakly similar to unnamed protein product [H.sapiens]"
745606	1.3470375	hypothetical protein PP591
1500000	1.3204029	"H2B histone family, member B"
595037	1.3057353	retinoic acid induced 3

2024

- 30 -

503215	0.9757636	pilin-like transcription factor
1911343	0.967716	"RAB26, member RAS oncogene family"
2029173	0.9672743	"ESTs, Weakly similar to N-WASP [H.sapiens]"
470061	0.9665514	seven in absentia (Drosophila) homolog 2
2016908	0.9615102	"ESTs, Weakly similar to CA13_HUMAN COLLAGEN ALPHA 1(III) CHAIN PRECURSOR [H.sapiens]"
1469425	0.9610542	SRY (sex determining region Y)-box 22
814054	0.958989	KIAA0040 gene product
814528	0.9569471	"Homo sapiens cDNA: FLJ22139 fis, clone HEP20959"
68636	0.9562688	hypothetical protein MGC2477
290841	0.9562201	"H2B histone family, member A"
236034	0.9534619	"uncoupling protein 2 (mitochondrial, proton carrier)"
782428	0.9524489	KIAA0250 gene product
131094	0.9518372	"Homo sapiens cDNA: FLJ21587 fis, clone COL06946"
488202	0.9501137	"ESTs, Weakly similar to YZ28_HUMAN HYPOTHETICAL PROTEIN ZAP128 [H.sapiens]"
1435862	0.9500093	"antigen identified by monoclonal antibodies 12E7, F21 and O13"
823598	0.9452282	"proteasome (prosome, macropain) 26S subunit, non-ATPase, 12"
1492426	0.9397097	chromosome 19 open reading frame 3
795185	0.9295627	xenotropic and polytropic retrovirus receptor
810558	0.9227709	"proteasome (prosome, macropain) 26S subunit, ATPase, 4"
469686	0.9203022	"Ric (Drosophila)-like, expressed in many tissues"
280375	0.9155003	PRO2000 protein
729975	0.9149894	meningioma expressed antigen 5 (hyaluronidase)
810124	0.9124035	"platelet-activating factor acetylhydrolase, isoform Ib, gamma subunit (29kD)"
813281	0.9107661	WW domain-containing protein 1
39884	0.9091807	IMP (inosine monophosphate) dehydrogenase 1
76605	0.9028609	nesca protein
1636092	0.8997881	hypothetical protein FLJ20657

- 31 -

741474	0.8925439	glucose phosphate isomerase
1605426	0.8909868	hypothetical protein FLJ13352
788654	0.8886269	
868128	0.8873517	JM4 protein
308466	0.8803492	GTP-binding protein Sara
44292	0.8732235	Homo sapiens mRNA; cDNA DKFZp434C107 (from clone DKFZp434C107)
826256	0.8700831	transmembrane 7 superfamily member 1 (upregulated in kidney)
685516	0.8694652	putative G protein-coupled receptor
810711	0.869424	stearoyl-CoA desaturase (delta-9-desaturase)
753299	0.8679015	hypothetical protein FLJ10504
731044	0.8657953	glutaredoxin 2
824052	0.8650616	chromosome 6 open reading frame 1
843195	0.86501	phosphoserine phosphatase
292770	0.8570995	"Homo sapiens, clone IMAGE:3627860, mRNA, partial cds"
859761	0.8563964	poliovirus receptor-related 2 (herpesvirus entry mediator B)
347373	0.8555396	"transcription elongation factor B (SIII), polypeptide 1 (15kD, elongin C)"
1640821	0.8544081	"ESTs, Weakly similar to I78885 serine/threonine-specific protein kinase [H.sapiens]"
704414	0.8520508	small nuclear ribonucleoprotein polypeptides B and B1
810725	0.8510425	"ATPase, H <sup>+</sup> transporting, lysosomal (vacuolar proton pump) 21kD"
782608	0.8507815	mitochondrial ribosomal protein L9
2019223	0.849044	mitochondrial ribosomal protein L17
2110511	0.8471736	artemin
1474955	0.8460414	"TATA box binding protein (TBP)-associated factor, RNA polymerase II, N, 68kD (RNA-binding protein 56)"
810612	0.842284	S100 calcium-binding protein A11 (calgizzarin)
744417	0.8409538	carnitine acetyltransferase
969877	0.8382113	"synaptosomal-associated protein, 25kD"
272529	0.8371698	phosphomannomutase 2
768570	0.8371438	hypothetical protein FLJ11280
824879	0.8366695	hypothetical protein MGC11275

sd-71385

T032137032004

- 32 -

2014034	0.8354357	"methylene tetrahydrofolate dehydrogenase (NAD+ dependent), methenyltetrahydrofolate cyclohydrolase"
2054635	0.831532	"proteasome (prosome, macropain) subunit, alpha type, 7"
2052113	0.8307595	hypothetical protein FLJ10903
742595	0.8291965	cyclin-dependent kinase 5
2016648	0.8263671	Homo sapiens mRNA; cDNA DKFZp434N1728 (from clone DKFZp434N1728)
1573251	0.823025	peroxisomal long-chain acyl-coA thioesterase
1869201	0.8202299	hypothetical protein MGC2745
37708	0.8194577	hypothetical protein MGC3101
241348	0.8182422	prenylcysteine lyase
810063	0.8168926	"growth factor, erv1 (S. cerevisiae)-like (augmenter of liver regeneration)"
731308	0.8166873	citrate synthase
209066	0.8120797	
590774	0.8100386	mitogen-activated protein kinase 13
1435003	0.8098225	tumor suppressing subtransferable candidate 1
502774	0.8091404	hypothetical protein FLJ20623
488505	0.808712	accessory proteins BAP31/BAP29
109863	0.8066906	epithelial membrane protein 2
813419	0.8043966	"hydroxyacyl-Coenzyme A dehydrogenase, type II"
1845169	0.8033348	"RAB35, member RAS oncogene family"
742707	0.7955239	"ESTs, Weakly similar to MUC2_HUMAN MUCIN 2 PRECURSOR [H.sapiens]"
594500	0.7935958	EST
1456701	0.7934389	B-cell CLL/lymphoma 9
271472	0.7901735	C3HC4-like zinc finger protein
1473922	0.7892502	"actin related protein 2/3 complex, subunit 3 (21 kD)"
51773	0.7858986	hypothetical protein MGC3077
898032	0.7852733	KIAA0097 gene product
340558	0.7791736	"actin related protein 2/3 complex, subunit 5 (16 kD)"
773922	0.7767604	KIAA0005 gene product
768064	0.7743382	"cytochrome P450, subfamily I (aromatic compound-inducible), polypeptide 1"



- 33 -

625923	0.7725056	phosphoenolpyruvate carboxykinase 2 (mitochondrial)
150003	0.77183	hypothetical protein FLJ13187
741977	0.7715372	"B-factor, properdin"
365738	0.7709456	ESTs
814350	0.7701748	insulin-degrading enzyme
2108077	0.7700354	CGI-112 protein
209066	0.7693907	serine/threonine kinase 15
564492	0.7672548	mitochondrial carrier homolog 2
366353	0.7664569	DKFZP564C186 protein
813751	0.7659116	"sialyltransferase 4C (beta-galactosidase alpha-2,3-sialyltransferase)"
781097	0.7633668	reticulon 3
233349	0.762952	hypothetical protein FLJ10761
825327	0.7622414	"Homo sapiens cDNA FLJ14105 fis, clone MAMMA1001202"
248649	0.761865	hypothetical protein FLJ13910
589232	0.7583348	hypothetical protein FLJ11506
564847	0.757516	ESTs
754628	0.7573763	ESTs
1500162	0.7571399	ESTs
1738208	0.7565056	"a disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 4"
842994	0.7563101	cathepsin Z
42408	0.7559848	hypothetical protein MGC4604
470124	0.7539249	RAD1 (S. pombe) homolog
809357	0.7535411	Bernardinelli-Seip congenital lipodystrophy 2 (seipin)
624667	0.7524325	CGI-92 protein
49351	0.75115	SEX gene
46248	0.7510436	ADP-ribosyltransferase (NAD+; poly (ADP-ribose) polymerase)
686552	0.7506994	golgi phosphoprotein 1
1631735	0.7505843	"Homo sapiens, clone IMAGE:3604336, mRNA, partial cds"
1536006	0.7499887	ESTs
83363	0.748646	protein-L-isoaspartate (D-aspartate) O-methyltransferase
713782	0.7484589	a disintegrin and metalloproteinase domain 15 (metargidin)

- 34 -

727078	0.7472345	"Homo sapiens cDNA: FLJ23602 fis, clone LNG15735"
503889	0.7454098	hypothetical protein FLJ10656
284261	0.7445419	uncharacterized hematopoietic stem/progenitor cells protein MDS030
1473289	0.7441483	protective protein for beta-galactosidase (galactosialidosis)
49273	0.742123	"solute carrier family 27 (fatty acid transporter), member 4"
785701	0.7413837	"RAB31, member RAS oncogene family"
41569	0.7412357	hypothetical protein FLJ12650
810402	0.7407907	hypothetical protein
1456348	0.7397574	N-acetylneuraminic acid phosphate synthase; sialic acid synthase
256619	0.7395032	hydroxysteroid (17-beta) dehydrogenase 7
743589	0.7372744	"ESTs, Weakly similar to T2D3_HUMAN TRANSCRIPTION INITIATION FACTOR TFIID 135 KDA SUBUNIT [H.sapiens]"
1409509	0.7370811	"troponin T1, skeletal, slow"
122241	0.7352191	"proteasome (prosome, macropain) subunit, beta type, 2"
470099	0.7343404	HT002 protein; hypertension-related calcium-regulated gene
725454	0.7335223	CDC28 protein kinase 2
796694	0.7317383	baculoviral IAP repeat-containing 5 (survivin)
186768	0.7298378	"Homo sapiens, clone IMAGE:3604869, mRNA"
813629	0.7288266	YME1 (S.cerevisiae)-like 1
842980	0.7284256	developmentally regulated GTP-binding protein 1
769921	0.726646	ubiquitin carrier protein E2-C
2322367	0.7249854	reticulon 4
951233	0.7240615	"proteasome (prosome, macropain) subunit, beta type, 3"
842825	0.7235007	G1 to S phase transition 1
782608	0.7230962	
2043167	0.7224903	BCL2-associated athanogene 3
827171	0.7211896	ESTs
120271	0.7208543	hypothetical protein MGC4692
1518890	0.7206844	"metallothionein-like 5, testis-specific (tesmin)"
60565	0.720547	lethal giant larvae (Drosophila) homolog 2

sd-71385

TOTAL 18:12:10

- 35 -

2050827	0.7156518	"proteasome (prosome, macropain) 26S subunit, ATPase, 5"
1635681	0.7152705	"NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 2 (8kD, B8)"
731023	0.7151176	WD repeat domain 5
1518402	0.7150061	KIAA1361 protein
752631	0.7139862	"fibroblast growth factor receptor 3 (achondroplasia, thanatophoric dwarfism)"
785616	0.7131593	"signal sequence receptor, alpha (translocon-associated protein alpha)"
686172	0.7131363	KIAA0008 gene product
40173	0.7124913	KIAA0807 protein
123441	0.7124698	ribosomal protein L7a
123614	0.7117834	hypothetical protein MGC4675
811024	0.7117211	bone marrow stromal cell antigen 2
2302099	0.7117055	sialidase 3 (membrane sialidase)
756442	0.7115925	P450 (cytochrome) oxidoreductase
811585	0.7087617	huntingtin (Huntington disease)
279970	0.7040135	adenosine A2a receptor
1517171	0.7036866	"interleukin 2 receptor, alpha"
838366	0.702617	3-hydroxymethyl-3-methylglutaryl-Coenzyme A lyase (hydroxymethylglutaricaciduria)
809944	0.7021108	KIAA0310 gene product
1393018	0.701717	"general transcription factor IIIC, polypeptide 1 (alpha subunit, 220kD )"
725978	0.7010756	"ESTs, Moderately similar to ALU7_HUMAN ALU SUBFAMILY SQ SEQUENCE CONTAMINATION WARNING ENTRY [H.sapiens]"
325606	0.7008219	hypothetical protein MGC14353
795256	0.6974866	NPD007 protein
365358	0.696963	pM5 protein
67765	0.6967324	carboxypeptidase M
358267	0.6962436	"EST, Moderately similar to AF119917 63 PRO2831 [H.sapiens]"
149355	0.6943923	translocating chain-associating membrane protein
212542	0.6938079	"Homo sapiens cDNA FLJ12900 fis, clone NT2RP2004321"
79520	0.6919579	"RAB2, member RAS oncogene family"

- 36 -

739126	0.6881663	tissue specific transplantation antigen P35B
202514	0.6872212	DNA (cytosine-5-)-methyltransferase 3 alpha
1884404	0.6864546	KIAA0285 gene product
814378	0.6857148	"serine protease inhibitor, Kunitz type, 2"
2018337	0.6828744	"glucosidase, beta; acid (includes glucosylceramidase)"
701751	0.6824676	cut (Drosophila)-like 1 (CCAAT displacement protein)
2018084	0.6822251	Ste-20 related kinase
781019	0.6819736	paraoxonase 2
2244196	0.6807351	B-cell receptor-associated protein BAP29
124447	0.679866	KIAA1184 protein
789012	0.6794432	fibulin 2
490778	0.6788304	low molecular mass ubiquinone-binding protein (9.5kD)
2011515	0.6774452	DKFZP586B0923 protein
80764	0.6765616	hypothetical protein
841679	0.6762195	calcium and integrin binding protein (DNA-dependent protein kinase interacting protein)
108425	0.6761526	
878406	0.6750777	metaxin 1
487733	0.6745496	
625693	0.6744622	hypothetical protein MGC10911
1325816	0.6742078	polymerase (RNA) II (DNA directed) polypeptide L (7.6kD)
278531	0.6737637	cytochrome c oxidase subunit VIc
744374	0.6732551	putative ankyrin-repeat containing protein
203003	0.6724027	"non-metastatic cells 4, protein expressed in"
564981	0.6712134	ESTs
812994	0.671102	"retinoid X receptor, alpha"
172517	0.6701228	hippocalcin-like 1
205049	0.6682666	protein kinase H11; small stress protein-like protein HSP22
1734309	0.6672454	sperm associated antigen 4
1631699	0.6665952	valosin-containing protein
2028949	0.6662774	hypothetical protein PRO1855

- 37 -

530197	0.6659784	"ESTs, Moderately similar to ALU8_HUMAN ALU SUBFAMILY SX SEQUENCE CONTAMINATION WARNING ENTRY [H.sapiens]"
199645	0.6654138	nicastrin
704254	0.6653635	a disintegrin and metalloproteinase domain 8
1592715	0.6637095	"Homer, neuronal immediate early gene, 3"
302031	0.6626629	Ste20-related serine/threonine kinase
239568	0.661735	annexin A9
770785	0.6613519	"1,2-alpha-mannosidase IC"
785795	0.660789	hypothetical protein FLJ12910
810734	0.6603412	"polymerase (DNA-directed), delta 4"
1696757	0.6591801	hypothetical protein KIAA1165
811761	0.6587469	Nijmegen breakage syndrome 1 (nibrin)
150314	0.6584621	lysophospholipase I
489351	0.65846	hypothetical protein DKFZp566J2046
593431	0.6578268	"ESTs, Moderately similar to CEGT_HUMAN CERAMIDE GLUCOSYLTRANSFERASE [H.sapiens]"
37554	0.6574292	hypothetical protein FLJ22353
126851	0.6559601	hypothetical protein FLJ11160
265103	0.6548137	Homo sapiens mRNA; cDNA DKFZp547M123 (from clone DKFZp547M123)
51083	0.6544472	"catenin (cadherin-associated protein), delta 2 (neural plakophilin-related arm-repeat protein)"
431505	0.6534315	"ESTs, Highly similar to A31026 probable membrane receptor protein [H.sapiens]"
1420370	0.6531171	biliverdin reductase B (flavin reductase (NADPH))
1476053	0.6522237	RAD51 (S. cerevisiae) homolog (E coli RecA homolog)
882484	0.6521026	"chaperonin containing TCP1, subunit 7 (eta)"
51532	0.6504769	ADP-ribosylation factor-like 6 interacting protein
280249	0.6503028	Kruppel-like factor 7 (ubiquitous)
138788	0.6497728	prolactin receptor
1492468	0.647846	DEME-6 protein

**700809-1**

- 39 -

770845	0.6253839	hexokinase 1
813707	0.6252347	regulator of G-protein signalling 16
611443	0.6244565	myoglobin
124331	0.624455	"cleavage and polyadenylation specific factor 5, 25 kD subunit"
855800	0.6235647	prolyl endopeptidase
625234	0.62188	KDEL (Lys-Asp-Glu-Leu) endoplasmic reticulum protein retention receptor 3
753378	0.6218232	hypothetical protein FLJ22649 similar to signal peptidase SPC22/23
2055807	0.6216372	protein kinase domains containing protein similar to phosphoprotein C8FW
840878	0.6216124	seladin-1
1700436	0.6215741	ESTs
261472	0.6215341	putative nuclear protein ORF1-FL49
121251	0.6209436	hypothetical protein MGC5576
140289	0.6205892	hypothetical protein
2018821	0.6205344	ATPase inhibitor precursor
770766	0.6193342	DKFZP564C1940 protein
767798	0.6192232	"ATX1 (antioxidant protein 1, yeast) homolog 1"
845363	0.617901	"non-metastatic cells 1, protein (NM23A) expressed in"
120749	0.6172281	"ESTs, Moderately similar to KIAA1215 protein [H.sapiens]"
358162	0.616765	protein predicted by clone 23627
781342	0.6159353	hypothetical protein MGC11115
767761	0.6142782	DKFZP434B168 protein
191904	0.6141265	"BUB3 (budding uninhibited by benzimidazoles 3, yeast) homolog"
726699	0.6131491	"ESTs, Weakly similar to AAB47496 NG5 [H.sapiens]"
2326057	0.61263	MLN51 protein
784140	0.6102245	WD repeat domain 15
770355	0.6099761	"lanosterol synthase (2,3-oxidosqualene-lanosterol cyclase)"
81336	0.6099072	uteroglobin
757328	0.6097389	hypothetical protein FLJ22678
773674	0.6095732	oncogene TC21
289978	0.6090234	ubiquitin-like 4
66535	0.6088919	ornithine decarboxylase antizyme 2

- 40 -

826363	0.6087183	lysophospholipase II
1758590	0.6074571	"fatty-acid-Coenzyme A ligase, long-chain 3"
1553306	0.6066787	"proteasome (prosome, macropain) 26S subunit, non-ATPase, 11"
785766	0.6058332	hypothetical protein
725841	0.6051304	KIAA0662 gene product
2020898	0.6044163	"procollagen-lysine, 2-oxoglutarate 5-dioxygenase 3"
504308	0.6041122	hypothetical protein FLJ10540
510575	0.6040713	hypothetical protein FLJ22087
49630	0.6027878	"calcium channel, voltage-dependent, L type, alpha 1D subunit"
142586	0.602037	MCT-1 protein
725284	0.6014099	"phosphorylase kinase, gamma 2 (testis)"
429799	0.6012713	hypothetical protein FLJ21939 similar to 5-azacytidine induced gene 2
1474424	0.6006702	"Homo sapiens cDNA FLJ12758 fis, clone NT2RP2001328"
784105	0.6003082	ESTs
2018527	0.599584	dolichyl-phosphate mannosyltransferase polypeptide 3
855563	0.5987092	v-erb-b2 avian erythroblastic leukemia viral oncogene homolog 3
823574	0.5978052	endosulfine alpha
323693	0.5975785	"adaptor-related protein complex 1, sigma 1 subunit"
824962	0.595868	"karyopherin alpha 2 (RAG cohort 1, importin alpha 1)"
773426	0.5941568	KIAA0391 gene product
772925	0.5940171	HSPCO34 protein
1656062	0.593408	coagulation factor XII (Hageman factor)
825740	0.5931142	DKFZp434J1813 protein
376516	0.5906248	cell division cycle 4-like
292213	0.5883092	"guanine nucleotide binding protein (G protein), beta polypeptide 2"
1898619	0.5880791	hypothetical protein MGC15737
414992	0.587733	K562 cell-derived leucine-zipper-like protein 1
1573946	0.5875624	programmed cell death 9
739109	0.5874334	"adaptor-related protein complex 2, sigma 1 subunit"



- 41 -

430235	0.585372	"H2B histone family, member Q"
2572170	0.5852444	"ESTs, Weakly similar to T26581 hypothetical protein Y32B12A.3 - <i>Caenorhabditis elegans</i> [C.elegans]"
487733-2	0.5844776	"NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 2 (8kD, AGGG)"
1698036	0.5843382	ubiquitin-conjugating enzyme E2 variant 1
127646	0.5838788	
346696	0.5838269	TEA domain family member 4
74738	0.5835427	"Homo sapiens, clone IMAGE:3535294, mRNA, partial cds"
124781	0.5832065	squalene epoxidase
1883028	0.5828007	Homo sapiens mRNA; cDNA DKFZp434J1912 (from clone DKFZp434J1912)
183200	0.5801312	fumarylacetoacetate hydrolase (fumarylacetoacetase)
51741	0.5795151	GTP-binding protein
590759	0.5791873	sterol-C4-methyl oxidase-like
755301	0.5790795	"protein kinase C, delta"
843054	0.5776102	KIAA1533 protein
595213	0.5775511	hypothetical protein
752643	0.5774681	group XII secreted phospholipase A2
141852	0.5762886	"purinergic receptor P2Y, G-protein coupled, 2"
813631	0.5754418	seven transmembrane protein TM7SF3
1601947	0.5744444	cytochrome c oxidase subunit VIIa polypeptide 2 (liver)
51657	0.5736969	hypothetical protein ET
814209	0.5736866	ESTs
72050	0.5722845	"chloride channel, nucleotide-sensitive, 1A"
1635665	0.5714089	"Homo sapiens, RIKEN cDNA 2010100O12 gene, clone MGC:14813 IMAGE:4133274, mRNA, complete cds"
66406	0.5706229	hypothetical protein DKFZp762E1312
530310	0.5686743	KIAA0143 protein
41356	0.5683316	"protein phosphatase 2, regulatory subunit B (B56), alpha isoform"
1844765	0.5670398	Homo sapiens mRNA; cDNA DKFZp564O1763 (from clone DKFZp564O1763)

- 42 -

489823	0.5666593	"COX17 (yeast) homolog, cytochrome c oxidase assembly protein"
1517749	0.5665848	ESTs
813410	0.5664086	polymerase (RNA) II (DNA directed) polypeptide K (7.0kD)
1416782	0.5657017	"creatine kinase, brain"
1422338	0.5652946	ribonucleotide reductase M2 polypeptide
250313	0.5649247	ESTs
235986	0.5649127	"wingless-type MMTV integration site family, member 11"
66317	0.5648355	"H1 histone family, member 2"
279720	0.5645072	"ESTs, Moderately similar to A47582 B-cell growth factor precursor [H.sapiens]"
884498	0.5640535	uncharacterized hypothalamus protein HT012
503851	0.564048	nuclear receptor co-repressor/HDAC3 complex subunit
823930	0.5633774	"actin related protein 2/3 complex, subunit 1A (41 kD)"
120271	0.5626611	hypothetical protein MGC4692
782513	0.5625811	"interferon, alpha-inducible protein (clone IFI-6-16)"
246800	0.5625675	hypothetical protein FLJ10803
2309073	0.5617024	frizzled (Drosophila) homolog 5
784150	0.5598483	"RAB31, member RAS oncogene family"
139835	0.5590106	UDP-glucose dehydrogenase
1641894	0.5587185	EST
796757	0.5585876	"adaptor-related protein complex 3, sigma 1 subunit"
813616	0.5574238	FK506-binding protein like
2549634	0.5572052	activator of S phase kinase
741769	0.5565863	"polymerase (DNA directed), beta"
488642	0.5560728	"ESTs, Weakly similar to I38022 hypothetical protein [H.sapiens]"
795498	0.5557245	putative transmembrane protein
1492780	0.5555908	"Homo sapiens cDNA FLJ14459 fis, clone HEMBB1002409"
241043	0.5545302	"Human clone 137308 mRNA, partial cds"
295986	0.5544422	emopamil-binding protein (sterol isomerase)
839682	0.5542571	ubiquitin-conjugating enzyme E2N (homologous to yeast UBC13)

- 43 -

1581941	0.5537015	hypothetical protein FLJ14540
809466	0.5532437	DNA segment on chromosome 19 (unique) 1177 expressed sequence
1947381	0.5524634	hypothetical protein FLJ22329
825470	0.5523318	topoisomerase (DNA) II alpha (170kD)
1848977	0.5519913	glycerol kinase
700792	0.5493223	cyclin-dependent kinase inhibitor 3 (CDK2-associated dual specificity phosphatase)
825386	0.5487104	"ATP synthase, H <sup>+</sup> transporting, mitochondrial F1F0, subunit d"
826194	0.5486839	synaptotagmin-like 2
299815	0.5485843	hypothetical protein DC42
1753497	0.5485231	ovo (Drosophila) homolog-like 1
744944	0.5482368	myosin VI
1499828	0.5479012	"fucosyltransferase 1 (galactoside 2-alpha-L- fucosyltransferase, Bombay phenotype included)"
415102	0.5465851	cell division cycle 25C
813387	0.5460577	diaphorase (NADH/NADPH) (cytochrome b-5 reductase)
786265	0.5443881	KIAA0750 gene product
1591264	0.5440944	transaldolase 1
756595	0.5433619	"S100 calcium-binding protein A10 (annexin II ligand, calpactin I, light polypeptide (p11))"
855749	0.5432466	triosephosphate isomerase 1
470092	0.5429717	like-glycosyltransferase
244764	0.5424555	B7 homolog 3
1637282	0.542121	hexokinase 2
417801	0.5414761	mitochondrial ribosomal protein L27
826077	0.5405591	pyruvate dehydrogenase (lipoamide) beta
757489	0.5386267	"tubulin, alpha 2"
327635	0.5368033	adenylate kinase 1
343731	0.5365018	
786067	0.5361882	cell division cycle 25B
2017415	0.5353408	centromere protein A (17kD)
292996	0.5341429	"tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein, eta polypeptide"
815781	0.5340254	heat shock 105kD
346942	0.5326646	"phosphatidylinositol glycan, class Q"

- 44 -

823694	0.5321203	"Homo sapiens chromosome 19, BAC CIT-HSPC_204F22 (BC228680), complete sequence; contains bacterial insertion element"
139354	0.5318709	hypothetical protein
221499	0.5317861	KIAA0508 protein
263727	0.5313082	"DNA segment, single copy probe LNS-CAI/LNS-CAII (deleted in polyposis)"
489594	0.5308806	hypothetical protein FLJ11565
824524	0.5296138	UDP-galactose transporter related
951216	0.5291832	"NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 10 (22kD, PDSW)"
841260	0.5285587	hypothetical protein
283739	0.5276555	"Homo sapiens cDNA FLJ14028 fis, clone HEMBA1003838"
49117	0.5274035	KIAA0215 gene product
1762111	0.5272967	natriuretic peptide receptor C/guanylate cyclase C (atrionatriuretic peptide receptor C)
358609	0.5266332	NADH dehydrogenase (ubiquinone) flavoprotein 3 (10kD)
884425	0.5265114	"chaperonin containing TCP1, subunit 5 (epsilon)"
795401	0.5264278	diacylglycerol O-acyltransferase (mouse) homolog
811907	0.5246416	hypothetical protein FLJ22056
1631747	0.5245873	male-enhanced antigen
365060	0.5244569	"RAB11A, member RAS oncogene family"
743220	0.5242028	hypothetical protein FLJ12517
705064	0.5238148	"transforming, acidic coiled-coil containing protein 3"
768452	0.5237494	"Homo sapiens EST from clone 491476, full insert"
743977	0.5237158	Homo sapiens mRNA for TL132
1568825	0.5205165	Arg/Abl-interacting protein ArgBP2
772898	0.520392	ribosomal protein S15a
366834	0.5191005	envoplakin
1616253	0.5190257	breast carcinoma amplified sequence 1
322617	0.5188645	v-ral simian leukemia viral oncogene homolog B (ras related; GTP binding protein)
825296	0.5178447	low density lipoprotein receptor defect C complementing

- 45 -

491524	0.5175752	mitochondrial ribosomal protein L13
143426	0.5173212	"ras homolog gene family, member B"
73009	0.516678	"ESTs, Weakly similar to A43932 mucin 2 precursor, intestinal [H.sapiens]"
293569	0.5162069	chromosome 1 open reading frame 21
296702	0.5162041	"deiodinase, iodothyronine, type I"
773286	0.5158979	"solute carrier family 9 (sodium/hydrogen exchanger), isoform 3 regulatory factor 1"
726658	0.5157542	"non-metastatic cells 3, protein expressed in"
753897	0.5153412	autocrine motility factor receptor
293727	0.5153298	hypothetical protein MGC861
810947	0.5151346	"LIS1-interacting protein NUDE1, rat homolog"
491465	0.5150527	hypothetical protein FLJ10035
814899	0.5142481	BCL2/adenovirus E1B 19kD-interacting protein 3-like
266500	0.5135558	ESTs
1159963	0.5133656	interferon regulatory factor 7
785707	0.5133005	protein regulator of cytokinesis 1
346134	0.5132068	calcium-regulated heat-stable protein (24kD)
486110	0.5127183	profilin 2
2017403	0.5117711	regulator of G-protein signalling 3
491527	0.5116183	"Homo sapiens, Similar to CG7083 gene product, clone MGC:10534 IMAGE:3957147, mRNA, complete cds"
769600	0.5106262	uracil-DNA glycosylase 2
1558233	0.5101529	ESTs
649084	0.5099122	carbonic anhydrase XI
825822	0.5097808	hypothetical protein
1461477	0.509331	Homo sapiens mRNA; cDNA DKFZp586I0324 (from clone DKFZp586I0324)
327506	0.5092192	Homo sapiens mRNA full length insert cDNA clone EUROIMAGE 327506
769942	0.509086	kinesin-like 4
505289	0.5080495	"angiotensin II, type I receptor-associated protein"
726439	0.5080059	CGI-143 protein
79726	0.5078677	"ESTs, Highly similar to T46395 hypothetical protein DKFZp434I1120.1 [H.sapiens]"

- 46 -

810762	0.5072899	SNARE protein
785793	0.5072717	"capping protein (actin filament) muscle Z-line, alpha 1"
81599	0.5067033	ubiquitin specific protease 14 (tRNA-guanine transglycosylase)
294397	0.5066683	DKFZP586A0522 protein
1947647	0.5064941	CGI-147 protein
795805	0.5064099	KIAA0332 protein
810959	0.505619	Rho GDP dissociation inhibitor (GDI) alpha
2322223	0.5047814	small nuclear ribonucleoprotein polypeptide A
472103	0.503987	"soc-2 (suppressor of clear, C.elegans) homolog"
810609	0.5036493	hypothetical protein PP1226
897813	0.5034689	polyadenylate binding protein-interacting protein 1
1500542	0.5032679	regulator of G-protein signalling 11
839746	0.5031787	"Homo sapiens, Similar to RIKEN cDNA 5830420C20 gene, clone IMAGE:3633379, mRNA, partial cds"
343607	0.5029133	AD-015 protein
649977	0.5002695	Homo sapiens clone CDABP0014 mRNA sequence
823940	0.4997308	"transducer of ERBB2, 1"
2250839	0.4994848	androgen receptor (dihydrotestosterone receptor; testicular feminization; spinal and bulbar muscular atrophy; Kennedy disease)
731080	0.4979864	hypothetical protein FLJ12661
753400	0.496254	CGI-204 protein
869375	0.4958886	"isocitrate dehydrogenase 2 (NADP+), mitochondrial"
154610	0.4948571	dynactin 4
130835	0.4946783	"Homo sapiens, Similar to clone FLB3816, clone IMAGE:3454380, mRNA"
859228	0.4942316	"isocitrate dehydrogenase 1 (NADP+), soluble"
41698	0.4938629	progesterone binding protein
30170	0.4936252	"caspase 3, apoptosis-related cysteine protease"
266218	0.4934791	hypothetical protein FLJ11350

- 47 -

509588	0.49319	"TATA box binding protein (TBP)-associated factor, RNA polymerase II, J, 20kD"
1916575	0.4923447	BCL2-interacting killer (apoptosis-inducing)
744994	0.4916322	hypothetical protein FLJ12242
487444	0.4887966	"cyclic AMP phosphoprotein, 19 kD"
503866	0.4884193	sperm autoantigenic protein 17
814353	0.4883684	phorbol-12-myristate-13-acetate-induced protein 1
815501	0.4879053	hypothetical protein MGC2721
746163	0.4863317	"ESTs, Weakly similar to ALU1_HUMAN ALU SUBFAMILY J SEQUENCE CONTAMINATION WARNING ENTRY [H.sapiens]"
347726	0.4837701	homeo box D8
897770	0.4836347	
1854539	0.4824023	GAP-associated tyrosine phosphoprotein p62 (Sam68)
41123	0.481642	"Homo sapiens, Similar to RIKEN cDNA 2210021G21 gene, clone MGC:14859 IMAGE:3621871, mRNA, complete cds"
856447	0.4816395	"interferon, gamma-inducible protein 30"
1568561	0.4808662	BCL2-like 1
744047	0.4802334	polo (Drosophia)-like kinase
290101	0.4800598	ESTs
455275	0.4788778	hypothetical protein FLJ23469
2069602	0.4786416	melanocortin 1 receptor (alpha melanocyte stimulating hormone receptor)
592801	0.4785497	"serine palmitoyltransferase, long chain base subunit 2"
741891	0.4775456	"RAB2, member RAS oncogene family-like"
1434948	0.4771454	HIV TAT specific factor 1
430614	0.476288	"2,3-bisphosphoglycerate mutase"
454896	0.4760701	"DnaJ (Hsp40) homolog, subfamily A, member 2"
268946	0.4751692	WD40 protein Ciao1
1572710	0.4750848	hypothetical protein FLJ21213
626318	0.4746692	ubiquitin 1
377384	0.4746085	"nuclear receptor subfamily 2, group F, member 2"

- 48 -

1632252	0.474559	"complement component 1, q subcomponent, alpha polypeptide"
489444	0.4741367	hypothetical protein FLJ20211
345787	0.4734963	"highly expressed in cancer, rich in leucine heptad repeats"
767487	0.4729158	"ariadne (Drosophila) homolog, ubiquitin-conjugating enzyme E2-binding protein, 1"
empty-31	0.4726918	blank
135640	0.4708685	syntaxin 3A
754653	0.4707927	"cadherin, EGF LAG seven-pass G-type receptor 3, flamingo (Drosophila) homolog"
1557637	0.4705061	ESTs
788641	0.4703576	"adaptor-related protein complex 1, sigma 2 subunit"
810567	0.4702908	"Homo sapiens, clone MGC:3182 IMAGE:3356293, mRNA, complete cds"
366039	0.4701924	KIAA0892 protein
781047	0.4699376	budding uninhibited by benzimidazoles 1 (yeast homolog)
782688	0.4694868	"dynein, axonemal, light intermediate polypeptide"
210862	0.4692392	"acyl-Coenzyme A oxidase 1, palmitoyl"
510794	0.4691467	c-myc binding protein
121436	0.4691298	"Homo sapiens, clone MGC:4677 IMAGE:3532809, mRNA, complete cds"
325515	0.4689353	hypothetical protein FLJ10980
965223	0.4687291	"thymidine kinase 1, soluble"
1469148	0.467634	FGFR1 oncogene partner
149539	0.4671676	KIAA1700
2306987	0.4670155	secreted and transmembrane 1
122147	0.4652151	
2063982	0.4651856	"potassium channel, subfamily K, member 6 (TWIK-2)"
769890	0.4647115	nucleoside phosphorylase
1642496	0.4645092	hypothetical protein MGC11266
428582	0.4642013	hypothetical protein FLJ20296
66599	0.4639123	N-acetyltransferase 1 (arylamine N-acetyltransferase)
1565455	0.4634802	"nuclear receptor subfamily 2, group C, member 2"
810939	0.4634658	hypothetical protein FLJ22169
359723	0.4630712	mitochondrial elongation factor G



- 49 -

1027283	0.4628012	ESTs
1636844	0.4627748	ring finger protein 14
469151	0.4622121	"eukaryotic translation initiation factor 2, subunit 2 (beta, 38kD )"
813675	0.4614894	"Human D9 splice variant B mRNA, complete cds"
489657	0.4609514	tryptophan rich basic protein
469383	0.4607375	chromosome 8 open reading frame 1
40042	0.4604322	hypothetical protein FLJ10747
726637	0.45959	t-complex-associated-testis-expressed 1-like
811142	0.4595393	"phosphoinositide-3-kinase, regulatory subunit, polypeptide 2 (p85 beta)"
179212	0.4594831	"ESTs, Moderately similar to T12539 hypothetical protein DKFZp434J154.1 [H.sapiens]"
292936	0.4593712	hypothetical protein FLJ10468
2046679	0.4579596	ESTs
810497	0.4572069	"ESTs, Weakly similar to A35363 synapsin I splice form a [H.sapiens]"
1585492	0.4571478	ESTs
782689	0.4570254	"solute carrier family 6 (neurotransmitter transporter, creatine), member 8"
774446	0.4567242	adrenomedullin
366156	0.4563893	"Homo sapiens cDNA FLJ14028 fis, clone HEMBA1003838"
725649	0.4559473	"nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 4"
76362	0.4553121	"spectrin, alpha, non-erythrocytic 1 (alpha-fodrin)"
754046	0.4550413	DNA segment on chromosome X (unique) 9879 expressed sequence
345538	0.4550133	cathepsin L
782439	0.454474	"ATP synthase, H <sup>+</sup> transporting, mitochondrial F0 complex, subunit e"
1537001	0.4539586	ESTs
345423	0.4538841	DKFZP564M112 protein
1702742	0.453453	"solute carrier family 7 (cationic amino acid transporter, y <sup>+</sup> system), member 5"
825659	0.4515758	N-myc downstream regulated
1652310	0.4512689	"Homo sapiens, clone MGC:19613 IMAGE:3833049, mRNA, complete cds"

- 50 -

1523225	0.4511104	oncostatin M receptor
40773	0.4510289	"guanine nucleotide binding protein (G protein), alpha z polypeptide"
35626	0.4509825	"Homo sapiens cDNA FLJ14201 fis, clone NT2RP3002955"
358217	0.4507923	glypican 4
214996	0.45033	ESTs
177827	0.4500254	synaptotagmin VII
1075635	0.4495618	MLSN1- and TRP-related
810039	0.4491986	defender against cell death 1
772220	0.4475721	for protein disulfide isomerase-related
70606	0.4475592	ESTs
1692195	0.4471181	smg GDS-ASSOCIATED PROTEIN
841621	0.4466003	"ESTs, Weakly similar to JE0350 Anterior gradient-2 [H.sapiens]"
855707	0.4456433	ESTs
1455394	0.4455695	cytochrome c
85804	0.44449	hypothetical protein FLJ21918
809512	0.4442565	hypothetical protein FLJ10767
753457	0.4435792	NADH dehydrogenase (ubiquinone) Fe-S protein 1 (75kD) (NADH-coenzyme Q reductase)
767289	0.4432685	hypothetical protein FLJ10055
565235	0.4428854	spermine synthase
611467	0.4428543	NADH dehydrogenase (ubiquinone) Fe-S protein 6 (13kD) (NADH-coenzyme Q reductase)
812105	0.4424976	ALL1-fused gene from chromosome 1q
1558642	0.4424721	hypothetical protein MGC2771
814989	0.4419384	"protein phosphatase 1G (formerly 2C), magnesium-dependent, gamma isoform"
1899312	0.4418248	ESTs
1486533	0.441804	keratin 16 (focal non-epidermolytic palmoplantar keratoderma)
1565079	0.4415966	B-cell linker
783681	0.4412503	upstream regulatory element binding protein 1
1422723	0.4411931	interferon-induced protein 35
785368	0.4411742	PDZ-binding kinase; T-cell originated protein kinase
1517595	0.4409591	KIAA0175 gene product
713862	0.4400704	ubiquitin-protein isopeptide ligase (E3)

- 51 -

83358	0.4397161	ESTs
81203	0.4391758	paraoxonase 3
74070	0.4390933	endosulfine alpha
2306221	0.4385881	"wingless-type MMTV integration site family, member 10B"
140635	0.4384907	ESTs
150118	0.4383146	hypothetical protein DKFZp434F054
214068	0.4383007	GATA-binding protein 3
1434905	0.4382073	homeo box B7
450301	0.4380657	mutL (E. coli) homolog 3
2027952	0.437854	hypothetical protein FLJ20297
745394	0.4374406	"Homo sapiens cDNA: FLJ23249 fis, clone COL04196"
824108	0.4369933	SCAN domain-containing 1
795543	0.4364791	thioredoxin peroxidase (antioxidant enzyme)
1592837	0.4355927	interferon induced transmembrane protein 2 (1-8D)
77533	0.435582	"inositol polyphosphate-5-phosphatase, 40kD"
1583198	0.4350835	"ESTs, Weakly similar to S65824 reverse transcriptase homolog [H.sapiens]"
84295	0.4349213	interleukin 1 receptor antagonist
770992	0.4346858	
837864	0.4345283	progesterone induced protein
1909574	0.4342055	mitochondrial ribosomal protein S11
128695	0.4340119	"ESTs, Weakly similar to I38344 titin, cardiac muscle [H.sapiens]"
742581	0.433453	"Homo sapiens cDNA FLJ10366 fis, clone NT2RM2001420"
52103	0.432756	hypothetical protein FLJ23045
770675	0.4319407	"Homo sapiens cDNA: FLJ21323 fis, clone COL02374"
756373	0.4310628	Rho guanine exchange factor (GEF) 16
43977	0.4309655	KIAA0182 protein
2322079	0.4309341	
25440	0.4304195	"staufen (Drosophila, RNA-binding protein) homolog 2"
207794	0.4297389	"nuclear factor (erythroid-derived 2), 45kD"
746190	0.4291743	hypothetical protein DKFZp761B1514
74566	0.4279059	"exportin 1 (CRM1, yeast, homolog)"
76196	0.4278385	hypothetical protein FLJ20062

- 52 -

826173	0.4276784	profilin 1
1587863	0.4270229	acetyl-Coenzyme A acyltransferase 1 (peroxisomal 3-oxoacyl-Coenzyme A thiolase)
2017721	0.4268089	"ESTs, Weakly similar to 2109260A B cell growth factor [H.sapiens]"
809464	0.426456	"fibroblast growth factor receptor 2 (bacteria-expressed kinase, keratinocyte growth factor receptor, craniofacial dysostosis 1, Crouzon syndrome, Pfeiffer syndrome, Jackson-Weiss syndrome)"
41826	0.4264531	ESTs
1687976	0.4257058	"endoplasmic reticulum chaperone SIL1, homolog of yeast"
416436	0.4252319	mitochondrial ribosomal protein L24
145491	0.4250102	protocadherin 1 (cadherin-like 1)
810316	0.4245985	very long-chain acyl-CoA synthetase; lipidosin
629944	0.4242364	myosin VB
1637302	0.424164	DNAJ domain-containing
754625	0.4236648	"ATPase, Class II, type 9A"
298417	0.4235025	trefoil factor 3 (intestinal)
47853	0.4231314	"aldehyde dehydrogenase 4 family, member A1"
809727	0.4224266	unc-51 (C. elegans)-like kinase 1
774502	0.4222849	"protein tyrosine phosphatase, non-receptor type 12"
785840	0.4219115	"SEC24 (S. cerevisiae) related gene family, member D"
823907	0.4218216	hypothetical protein FLJ10511
1679942	0.4215618	KIAA1053 protein
823909	0.4215098	
1600239	0.421459	HSPC037 protein
85195	0.4207753	"growth arrest and DNA-damage-inducible, gamma"
1640282	0.4205899	KIAA1240 protein
782503	0.4205398	fatty acid desaturase 1
266312	0.4201146	"ATPase, Cu++ transporting, beta polypeptide (Wilson disease)"
884511	0.4199518	cytochrome c oxidase subunit VIIb
2051697	0.4197703	"inhibitor of kappa light polypeptide gene enhancer in B-cells, kinase beta"

- 53 -

724888	0.4196322	"cytochrome P450, subfamily IVB, polypeptide 1"
809507	0.4189998	hypothetical protein FLJ20568
151449	0.4189856	"protein tyrosine phosphatase, non-receptor type 21"
743860	0.4189656	"hypothetical protein, clone 2746033"
345833	0.4187433	heterogeneous nuclear ribonucleoprotein A/B
259950	0.4187278	hypothetical protein FLJ14991
283751	0.4179154	cortistatin
810875	0.4176896	"solute carrier family 26, member 6"
26883	0.4176151	"protein kinase (cAMP-dependent, catalytic) inhibitor beta"
38356	0.4173012	"cytochrome P450, subfamily 46 (cholesterol 24-hydroxylase)"
35191	0.4165847	stromal cell-derived factor 2
75859	-2.4150177	N-myc downstream-regulated gene 2
1569187	-1.9090486	heparan sulfate (glucosamine) 3-O-sulfotransferase 4
345670	-1.889471	"ESTs, Moderately similar to I59348 CCAAT binding transcription factor CBF subunit C - rat [R.norvegicus]"
760299	-1.8654555	dickkopf (Xenopus laevis) homolog 3
753071	-1.8417236	"Homo sapiens cDNA: FLJ22528 fis, clone HRC12825"
344720	-1.7951928	glycophorin C (Gerbich blood group)
810002	-1.7256692	"Homo sapiens, clone MGC:19762 IMAGE:3636045, mRNA, complete cds"
200814	-1.7140538	"membrane metallo-endopeptidase (neutral endopeptidase, enkephalinase, CALLA, CD10)"
131839	-1.7087069	folate receptor 1 (adult)
300632	-1.6969092	hypothetical protein FLJ21044 similar to Rbig1
811920	-1.6924158	"interleukin 11 receptor, alpha"
486683	-1.6799604	Homo sapiens mRNA; cDNA DKFZp564J0323 (from clone DKFZp564J0323)
1558675	-1.6577507	SRY (sex determining region Y)-box 10
840266	-1.6530726	"Homo sapiens cDNA: FLJ22667 fis, clone HSI08385"
285377	-1.6514168	pellino (Drosophila) homolog 2
1469377	-1.6473741	lipoma HMGIC fusion partner-like 2

- 54 -

45099	-1.6325981	regucalcin (senescence marker protein-30)
1161564	-1.5917197	desmuslin
298122	-1.5678888	frizzled (Drosophila) homolog 7
796542	-1.5553964	ets variant gene 5 (ets-related molecule)
289760	-1.5534789	"ESTs, Highly similar to T00391 hypothetical protein KIAA0612 [H.sapiens]"
22917	-1.5156635	Homo sapiens mRNA; cDNA DKFZp761M0111 (from clone DKFZp761M0111)
838478	-1.5059801	neurocalcin delta
788234	-1.4824055	"inhibitor of DNA binding 4, dominant negative helix-loop-helix protein"
377461	-1.4787671	"caveolin 1, caveolae protein, 22kD"
67741	-1.4786722	PP2135 protein
839736	-1.4745405	"crystallin, alpha B"
712139	-1.4736508	ADP-ribosylation factor-like 7
160192	-1.4695435	"ESTs, Weakly similar to 2004399A chromosomal protein [H.sapiens]"
742685	-1.4440371	disabled (Drosophila) homolog 2 (mitogen-responsive phosphoprotein)
1556433	-1.4191955	GRO3 oncogene
1554167	-1.4126653	hypothetical protein FLJ14529
877621	-1.3987023	nGAP-like protein
1882697	-1.3873266	peanut (Drosophila)-like 2
72778	-1.3873177	"caspase 7, apoptosis-related cysteine protease"
666879	-1.3624072	annexin A8
69002	-1.3581187	PPAR(gamma) angiopoietin related protein
811848	-1.3549757	hypothetical protein
490023	-1.3516302	hypothetical protein MGC2648
781017	-1.3232108	early growth response 2 (Krox-20 (Drosophila) homolog)
767202	-1.3217103	latent transforming growth factor beta binding protein 2
291478	-1.3155899	runt-related transcription factor 3
757191	-1.2902603	ESTs
130201	-1.2885369	intercellular adhesion molecule 2
1635320	-1.2878152	"amiloride-sensitive cation channel 2, neuronal"

sd-71385

- 56 -

1917449 -1.1785893 "serum amyloid A4, constitutive"  
1493218 -1.164886 hypothetical protein FLJ22297  
843070 -1.1647509 nucleoporin 88kD  
753038 -1.1629978 kinesin family member C3  
811088 -1.1625002 ephrin-B3  
2106144 -1.1619925 regulated in glioma  
796181 -1.1584587 growth arrest-specific 6  
1257131 -1.1566479 ESTs  
1473274 -1.1557062 "myosin regulatory light chain 2, smooth muscle isoform"  
  
208718 -1.1498532 annexin A1  
811837 -1.1468554 eukaryotic translation elongation factor 1 alpha 1  
  
768571 -1.1461886 SRY (sex determining region Y)-box 8  
1474900 -1.1347123 keratin 15  
756708 -1.1346161 "potassium intermediate/small conductance calcium-  
activated channel, subfamily N, member 4"  
  
1555924 -1.1342985 CSR1 protein  
811837 -1.1269309  
1878409 -1.1144369 catechol-O-methyltransferase  
786069 -1.1143906 beta-site APP-cleaving enzyme  
815737 -1.1116223 "ATP synthase, H<sup>+</sup> transporting, mitochondrial F1 complex,  
alpha subunit, isoform 1, cardiac muscle"  
  
132857 -1.0989746 Homo sapiens mRNA; cDNA DKFZp586N1323 (from clone  
DKFZp586N1323)  
  
1871116 -1.0975491 Homo sapiens mRNA; cDNA DKFZp434C1714 (from clone  
DKFZp434C1714); partial cds  
  
133236 -1.0945253 "RNA binding motif protein, X chromosome"  
  
283023 -1.0825166 chemokine (C-X3-C) receptor 1  
2016775 -1.0761731 "G protein-coupled receptor, family C, group 5, member B"  
  
897963 -1.0751021 phosphatidic acid phosphatase type 2A  
814316 -1.0654718 ribosomal protein L13  
47043 -1.0559861 tensin  
625399 -1.0523138 hypothetical protein similar to beta-transducin family  
  
416959 -1.051868 nuclear factor I/B  
366100 -1.0467604 matrilin 2



- 57 -

753162 -1.0416438 KIAA0603 gene product  
491403 -1.0411657 "tumor necrosis factor receptor superfamily, member 1B"  
  
813256 -1.0407652 "ATP-binding cassette, sub-family B (MDR/TAP), member  
1"  
740620 -1.0395569 tropomyosin 2 (beta)  
1577736 -1.035115 epidermal growth factor (beta-urogastrone)  
70245 -1.0348655 Homo sapiens mRNA full length insert cDNA clone  
EUROIMAGE 50374  
1626996 -1.0328113 c-fos induced growth factor (vascular endothelial growth  
factor D)  
360254 -1.0325653 "cysteine-rich, angiogenic inducer, 61"  
80344 -1.0323882 interleukin 7 receptor  
530958 -1.0316423 smoothened (Drosophila) homolog  
34150 -1.0302576 ESTs  
809784 -1.030253 "kallikrein 6 (neurosin, zyme)"  
990881 -1.0289558  
416434 -1.0284229 SCN Circadian Oscillatory Protein (SCOP)  
162308 -1.023071  
796475 -1.0217348 four and a half LIM domains 3  
32489 -1.0196444 hypothetical protein DKFZp566A1524  
2504881 -1.0194516 signal transducer and activator of transcription 5A  
  
188335 -1.0161259 "egf-like module containing, mucin-like, hormone receptor-  
like sequence 2"  
593023 -1.0135883 "dystrobrevin, beta"  
139660 -1.0134677 ESTs  
772913 -1.0122192 calreticulin  
270826 -1.0112851 "Homo sapiens cDNA FLJ13329 fis, clone  
OVARC1001795"  
1864302 -1.0099902 E74-like factor 5 (ets domain transcription factor)  
  
1587710 -1.0092611 period (Drosophila) homolog 1  
1584540 -1.0077508 Homo sapiens mRNA; cDNA DKFZp586M2022 (from clone  
DKFZp586M2022)  
  
788136 -1.0073983 "phosphodiesterase 4B, cAMP-specific (dunce  
(Drosophila)-homolog phosphodiesterase E4)"  
  
40027 -1.0057005 ESTs  
1856063 -1.0036618 tweety (Drosophila) homolog 1  
209537 -0.9988461 zinc finger protein 221

- 58 -

628955 -0.997687 forkhead box O1A (rhabdomyosarcoma)  
2095066 -0.9972552 "alcohol dehydrogenase 1C (class I), gamma polypeptide"  
  
1636166 -0.9969091 KIAA0668 protein  
882248 -0.9965858 transgelin  
415816 -0.9954308 ESTs  
897731 -0.9927981 latrophilin  
868396 -0.9916319 "tyrosine 3-monooxygenase/tryptophan 5-monooxygenase  
activation protein, eta polypeptide"  
  
343760 -0.9907562 SH3 domain binding glutamic acid-rich protein like 2  
1572298 -0.9906095 "CD3Z antigen, zeta polypeptide (TiT3 complex)"  
  
1474337 -0.986117 "phosphorylase, glycogen; brain"  
1609665 -0.9851713 BarH-like homeobox 2  
2243051 -0.9835274 kallikrein 8 (neuropsin/ovasin)  
27544 -0.9834143 prominin (mouse)-like 1  
281190 -0.9830533 ESTs  
161456 -0.9820939 serum amyloid A1  
505243 -0.980381 "inositol 1,4,5-triphosphate receptor, type 2"

Example IV: Genes for discriminating between normal and ADH

As shown in Table 3 below, 600 genes were identified as being able to discriminate between normal and ADH.

Table 3

ClonID	Weight	Description
1323448	1.3237504	cysteine-rich protein 1 (intestinal)
594500	1.158305	EST
824052	1.034408	chromosome 6 open reading frame 1
149539	0.9878366	KIAA1700
788654	0.9585887	growth factor receptor-bound protein 2
1517749	0.947675	ESTs
1492238	0.9155471	HSPC003 protein

sd-71385

- 59 -

366353	0.9154768	DKFZP564C186 protein
1609836	0.898716	glutamate-ammonia ligase (glutamine synthase)
138189	0.8950816	Wolfram syndrome 1 (wolframin)
344959	0.8945717	gene for serine/threonine protein kinase
1435862	0.8934643	antigen identified by monoclonal antibodies 12E7, F21 and O13
2014373	0.8820247	HNK-1 sulfotransferase
23776	0.879215	quinoid dihydropteridine reductase
293569	0.8708985	chromosome 1 open reading frame 21
130835	0.8659688	Homo sapiens, Similar to clone FLB3816, clone IMAGE:3454380, mRNA
236034	0.8541082	uncoupling protein 2 (mitochondrial, proton carrier)
488025	0.8525751	Kruppel-associated box protein
178805	0.8524069	Human DNA sequence from clone RP5-850E9 on chromosome 20. Contains part of the gene for a novel C2H2 type zinc finger protein similar to Drosophila Scratch (Scrt), Slug and Xenopus Snail, a novel gene similar to Drosophila CG6762, STSs, GSSs and five CpG
745606	0.8514931	hypothetical protein PP591
1492463	0.8378401	selenoprotein X, 1
530197	0.8206765	ESTs, Moderately similar to ALU8_HUMAN ALU SUBFAMILY SX SEQUENCE CONTAMINATION WARNING ENTRY [H.sapiens]
898222	0.8092364	Homo sapiens clone 24418 mRNA sequence
782608	0.8041049	mitochondrial ribosomal protein L9
450301	0.8030667	mutL (E. coli) homolog 3
196189	0.7992668	cytochrome b-5
363144	0.7924672	transcription factor AP-2 beta (activating enhancer-binding protein 2 beta)
68636	0.7904354	hypothetical protein MGC2477
795185	0.7765346	xenotropic and polytropic retrovirus receptor
255754	0.7720883	LUNIX protein; PLUNC (palate lung and nasal epithelium clone); tracheal epithelium enriched protein
767798	0.7682106	ATX1 (antioxidant protein 1, yeast) homolog 1

- 60 -

1558233	0.7654043	ESTs
769600	0.763743	uracil-DNA glycosylase 2
1156538	0.7512287	potassium inwardly-rectifying channel, subfamily J, member 11
1587863	0.7507482	acetyl-Coenzyme A acyltransferase 1 (peroxisomal 3-oxoacyl-Coenzyme A thiolase)
1709791	0.740832	BAI1-associated protein 1
741891	0.7360507	RAB2, member RAS oncogene family-like
788654	0.7336359	
811774	0.7223819	CGI-49 protein
1738208	0.7166687	a disintegrin-like and metalloprotease (repolysin type) with thrombospondin type 1 motif, 4
120138	0.7096046	J domain containing protein 1
2248488	0.7033736	ems1 sequence (mammary tumor and squamous cell carcinoma-associated (p80/85 src substrate)
265103	0.702883	Homo sapiens mRNA; cDNA DKFZp547M123 (from clone DKFZp547M123)
2054122	0.698129	solute carrier family 11 (proton-coupled divalent metal ion transporters), member 3
701751	0.6961443	cut (Drosophila)-like 1 (CCAAT displacement protein)
811582	0.690137	golgi phosphoprotein 2
266312	0.6856592	ATPase, Cu++ transporting, beta polypeptide (Wilson disease)
35147	0.6842172	ESTs, Weakly similar to unnamed protein product [H.sapiens]
1325816	0.6829962	polymerase (RNA) II (DNA directed) polypeptide L (7.6kD)
358217	0.6801945	glypican 4
233349	0.6764709	hypothetical protein FLJ10761
741977	0.6758385	B-factor, properdin
145132	0.6725186	mannose-P-dolichol utilization defect 1
33267	0.6679827	
279720	0.6607182	ESTs, Moderately similar to A47582 B-cell growth factor precursor [H.sapiens]
565319	0.6603668	Homo sapiens mRNA; cDNA DKFZp564B1264 (from clone DKFZp564B1264)
771173	0.6594128	mitochondrial ribosomal protein S21

[illegible]

- 62 -

814350	0.6173648	insulin-degrading enzyme
611443	0.6112713	myoglobin
782608	0.6105794	
22778	0.6100322	sulfortranferase family 4A, member 1
128695	0.608873	ESTs, Weakly similar to I38344 titin, cardiac muscle [H.sapiens]
47853	0.6077928	aldehyde dehydrogenase 4 family, member A1
814209	0.6076904	ESTs
590774	0.6071935	mitogen-activated protein kinase 13
969877	0.6064355	synaptosomal-associated protein, 25kD
1469425	0.6055267	SRY (sex determining region Y)-box 22
301122	0.6041028	extracellular matrix protein 1
199403	0.6027193	lectin, galactoside-binding, soluble, 8 (galectin 8)
813845	0.6018909	RNA, U transporter 1
810331	0.6014164	quiescin Q6
770766	0.600234	DKFZP564C1940 protein
279970	0.5992634	adenosine A2a receptor
490778	0.5957445	low molecular mass ubiquinone-binding protein (9.5kD)
288999	0.5955672	small protein effector 1 of Cdc42
504959	0.5953946	Homo sapiens mRNA; cDNA DKFZp586G0321 (from clone DKFZp586G0321)
42408	0.5930231	hypothetical protein MGC4604
263727	0.5926363	DNA segment, single copy probe LNS-CAI/LNS-CAII (deleted in polyposis)
854763	0.5918867	Homo sapiens cDNA FLJ11341 fis, clone PLACE1010786
729975	0.5908615	meningioma expressed antigen 5 (hyaluronidase)
1733262	0.5891897	BLu protein
730313	0.5887579	ESTs, Weakly similar to A43932 mucin 2 precursor, intestinal [H.sapiens]
1500000	0.5871787	H2B histone family, member B
595037	0.5870709	retinoic acid induced 3
1635681	0.5859113	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 2 (8kD, B8)
2325804	0.5840656	95 kDa retinoblastoma protein binding protein
2014856	0.5837779	major histocompatibility complex, class I-like sequence

- 63 -

40173	0.5771304	KIAA0807 protein
796723	0.576028	Homo sapiens clone CDABP0014 mRNA sequence
239568	0.5750382	annexin A9
2018527	0.5732092	dolichyl-phosphate mannosyltransferase polypeptide 3
1649374	0.5731982	homogentisate 1,2-dioxygenase (homogentisate oxidase)
809357	0.571929	Bernardinelli-Seip congenital lipodystrophy 2 (seipin)
366132	0.5711651	succinate dehydrogenase complex, subunit C, integral membrane protein, 15kD
83358	0.5709886	ESTs
1762111	0.5703701	natriuretic peptide receptor C/guanylate cyclase C (atrionatriuretic peptide receptor C)
244801	0.5693836	Rho guanine exchange factor (GEF) 11
127646	0.5663395	
39884	0.5659552	IMP (inosine monophosphate) dehydrogenase 1
202577	0.5629317	histamine N-methyltransferase
588262	0.5614929	Homo sapiens, Similar to RIKEN cDNA 2600001A11 gene, clone MGC:9907 IMAGE:3870073, mRNA, complete cds
813419	0.5602324	hydroxyacyl-Coenzyme A dehydrogenase, type II
1911343	0.559091	RAB26, member RAS oncogene family
859761	0.5582503	poliovirus receptor-related 2 (herpesvirus entry mediator B)
342181	0.557997	B-cell CLL/lymphoma 2
1558108	0.5577768	ATP-binding cassette, sub-family C (CFTR/MRP), member 8
2017403	0.5568495	regulator of G-protein signalling 3
122394	0.5554851	guanine nucleotide binding protein (G protein), alpha 15 (Gq class)
824879	0.5548666	hypothetical protein MGC11275
1708055	0.554593	glioblastoma overexpressed
1456701	0.5545861	B-cell CLL/lymphoma 9
186301	0.5537115	Homo sapiens cDNA FLJ12924 fis, clone NT2RP2004709
126851	0.5535041	hypothetical protein FLJ11160
826622	0.5532463	KIAA0430 gene product

- 64 -

151449	0.5503739	protein tyrosine phosphatase, non-receptor type 21
826668	0.550158	KIAA0274 gene product
76605	0.5482232	nesca protein
786072	0.5450231	myosin IC
2244196	0.5432772	B-cell receptor-associated protein BAP29
704414	0.5431136	small nuclear ribonucleoprotein polypeptides B and B1
739578	0.5402953	GPI-anchored metastasis-associated protein homolog
1404774	0.5399447	parathyroid hormone-like hormone
66599	0.5399379	N-acetyltransferase 1 (arylamine N-acetyltransferase)
359250	0.5382989	carbonic anhydrase IV
488202	0.5378254	ESTs, Weakly similar to YZ28_HUMAN HYPOTHETICAL PROTEIN ZAP128 [H.sapiens]
235986	0.5371486	wingless-type MMTV integration site family, member 11
214068	0.5357344	GATA-binding protein 3
1500542	0.5341753	regulator of G-protein signalling 11
470092	0.5319336	like-glycosyltransferase
589115	0.531314	matrix metalloproteinase 1 (interstitial collagenase)
810734	0.5308572	polymerase (DNA-directed), delta 4
685516	0.5304998	putative G protein-coupled receptor
171912	0.5297738	Homo sapiens cDNA FLJ10960 fis, clone PLACE1000564
1858892	0.5295858	hypothetical protein MGC4825
414992	0.5293379	K562 cell-derived leucine-zipper-like protein 1
37708	0.5283276	hypothetical protein MGC3101
713782	0.5273402	a disintegrin and metalloproteinase domain 15 (metargidin)
471568	0.5255851	hematological and neurological expressed 1
812238	0.5237287	hypothetical protein MGC4692
1631735	0.5228778	Homo sapiens, clone IMAGE:3604336, mRNA, partial cds
74070	0.5225238	endosulfine alpha
782497	0.5218113	Homo sapiens, clone IMAGE:3010666, mRNA, partial cds
502774	0.5204735	hypothetical protein FLJ20623



- 65 -

1517171	0.5204242	interleukin 2 receptor, alpha
50562	0.5195335	chromosome 8 open reading frame 4
60565	0.5192032	lethal giant larvae (Drosophila) homolog 2
1641894	0.5187462	EST
51083	0.5170561	catenin (cadherin-associated protein), delta 2 (neural plakophilin-related arm-repeat protein)
296702	0.5164562	deiodinase, iodothyronine, type I
2018337	0.5163293	glucosidase, beta; acid (includes glucosylceramidase)
1605426	0.5150013	hypothetical protein FLJ13352
1474149	0.5138729	poliovirus receptor-related 1 (herpesvirus entry mediator C; nectin)
26294	0.5137265	RNB6
810017	0.5135433	plasminogen activator, urokinase receptor
825822	0.5134094	hypothetical protein
810725	0.51329	ATPase, H <sup>+</sup> transporting, lysosomal (vacuolar proton pump) 21kD
781019	0.5132896	paraoxonase 2
726658	0.5132765	non-metastatic cells 3, protein expressed in
1916575	0.5127478	BCL2-interacting killer (apoptosis-inducing)
1469148	0.5125351	FGFR1 oncogene partner
726703	0.5118991	Homo sapiens clone 23736 mRNA sequence
725978	0.5108817	ESTs, Moderately similar to ALU7_HUMAN ALU SUBFAMILY SQ SEQUENCE CONTAMINATION WARNING ENTRY [H.sapiens]
1573946	0.509585	programmed cell death 9
51741	0.5094912	GTP-binding protein
156363	0.5086284	hypothetical protein FLJ12934
177827	0.5085582	synaptotagmin VII
294397	0.505999	DKFZP586A0522 protein
826077	0.503983	pyruvate dehydrogenase (lipoamide) beta
949938	0.5011384	cystatin C (amyloid angiopathy and cerebral hemorrhage)
1957136	0.5006732	phenylethanolamine N-methyltransferase
2018084	0.4999242	Ste-20 related kinase
784140	0.4984039	WD repeat domain 15
810981	0.498146	hypothetical protein FLJ20699

- 66 -

627248	0.4972176	SBBI31 protein
2302099	0.4967769	sialidase 3 (membrane sialidase)
302031	0.49677	Ste20-related serine/threonine kinase
183440	0.4967506	arylsulfatase A
1609748	0.4955206	hypothetical protein MGC10882
183200	0.4955056	fumarylacetoacetate hydrolase (fumarylacetoacetase)
2306697	0.4953741	neuromedin B
292770	0.4944242	Homo sapiens, clone IMAGE:3627860, mRNA, partial cds
856447	0.494183	interferon, gamma-inducible protein 30
855563	0.4941473	v-erb-b2 avian erythroblastic leukemia viral oncogene homolog 3
839081	0.4939384	homolog of yeast long chain polyunsaturated fatty acid elongation enzyme 2
49273	0.4937997	solute carrier family 27 (fatty acid transporter), member 4
289505	0.4933344	ESTs, Moderately similar to S65657 alpha-1C-adrenergic receptor splice form 2 [H.sapiens]
782689	0.4930071	solute carrier family 6 (neurotransmitter transporter, creatine), member 8
767495	0.4922026	GLI-Kruppel family member GLI3 (Greig cephalopolysyndactyly syndrome)
782537	0.490998	Homo sapiens cDNA: FLJ22562 fis, clone HSI01814
855029	0.4904632	Ac-like transposable element
705274	0.4904221	diacylglycerol kinase, delta (130kD)
202514	0.4902228	DNA (cytosine-5-)-methyltransferase 3 alpha
725649	0.4878638	nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 4
1884404	0.4878341	KIAA0285 gene product
825296	0.4858551	low density lipoprotein receptor defect C complementing
839580	0.4826315	ESTs
289857	0.4823282	phenylethanolamine N-methyltransferase
491465	0.4821169	hypothetical protein FLJ10035
360778	0.4820033	
261472	0.4814013	putative nuclear protein ORF1-FL49
840768	0.4812559	ATPase, H+ transporting, lysosomal (vacuolar proton pump) 16kD



- 68 -

810402	0.4583168	hypothetical protein
767761	0.4565666	DKFZP434B168 protein
1473289	0.4540707	protective protein for beta-galactosidase (galactosialidosis)
773142	0.4530754	hypothetical protein MGC2827
1652310	0.4525789	Homo sapiens, clone MGC:19613 IMAGE:3833049, mRNA, complete cds
812143	0.4519186	fibronectin leucine rich transmembrane protein 3
2326057	0.4506179	MLN51 protein
1075635	0.4502629	MLSN1- and TRP-related
825327	0.4497029	Homo sapiens cDNA FLJ14105 fis, clone MAMMA1001202
809838	0.4466933	
786545	0.4453714	protein phosphatase 1, regulatory (inhibitor) subunit 3B
183556	0.4452493	gap junction protein, alpha 4, 37kD (connexin 37)
503889	0.4433034	hypothetical protein FLJ10656
725321	0.44288	estrogen receptor 1
731459	0.4427978	Homo sapiens cDNA: FLJ22296 fis, clone HRC04468
752631	0.4426961	fibroblast growth factor receptor 3 (achondroplasia, thanatophoric dwarfism)
1558642	0.4426424	hypothetical protein MGC2771
120271	0.4425879	hypothetical protein MGC4692
487932	0.4419815	synaptotagmin-like 2
1592715	0.4417898	Homer, neuronal immediate early gene, 3
308466	0.4417728	GTP-binding protein Sara
840882	0.441569	nucleotide binding protein
823727	0.4406546	Homo sapiens, clone IMAGE:2905978, mRNA, partial cds
1636092	0.4404649	hypothetical protein FLJ20657
1505038	0.43988	hypothetical protein FLJ20171
490965	0.4389746	ESTs
346696	0.437234	TEA domain family member 4
345032	0.4360992	ESTs
487733	0.4356439	
810612	0.4352602	S100 calcium-binding protein A11 (calgizzarin)
122077	0.434495	putative membrane protein
417801	0.4325892	mitochondrial ribosomal protein L27



sd-71385

- 71 -

753301	-0.8393651	carcinoembryonic antigen-related cell adhesion molecule 1 (biliary glycoprotein)
51078	-0.8348835	ribosomal protein L44
3172883	-0.8293392	ESTs, Weakly similar to 1709359A dopamine D4 receptor [H.sapiens]
877621	-0.8289253	nGAP-like protein
811920	-0.8286379	interleukin 11 receptor, alpha
156211	-0.8269263	ATPase, H <sup>+</sup> transporting, lysosomal (vacuolar proton pump), beta polypeptide, 56/58kD, isoform 1 (Renal tubular acidosis with deafness)
760299	-0.8251086	dickkopf (Xenopus laevis) homolog 3
285377	-0.8214706	pellino (Drosophila) homolog 2
345034	-0.8161772	small inducible cytokine subfamily B (Cys-X-Cys), member 14 (BRAK)
298122	-0.8158372	frizzled (Drosophila) homolog 7
842896	-0.8123068	hypothetical protein DKFZp762L0311
843070	-0.8071984	nucleoporin 88kD
1257131	-0.8069059	ESTs
796539	-0.803124	KRAB-associated protein 1
139660	-0.79833	ESTs
666879	-0.7950485	annexin A8
200814	-0.7920988	membrane metallo-endopeptidase (neutral endopeptidase, enkephalinase, CALLA, CD10)
2016775	-0.7914834	G protein-coupled receptor, family C, group 5, member B
1946448	-0.790849	caveolin 2
1473471	-0.7906678	KIAA0194 protein
1556433	-0.7877792	GRO3 oncogene
1636156	-0.7827089	hypothetical protein FLJ21709
40299	-0.78007	growth differentiation factor 10
1636166	-0.7788972	KIAA0668 protein
1609665	-0.7765472	BarH-like homeobox 2
250797	-0.7739669	hypothetical protein FLJ20038
220395	-0.7687195	hypothetical protein FLJ23293 similar to ARL-6 interacting protein-2
208718	-0.7660278	annexin A1
41869	-0.7646494	hypothetical protein FLJ11017
490023	-0.7621736	hypothetical protein MGC2648
1607229	-0.7562938	tumor protein D52-like 1
133236	-0.7561129	RNA binding motif protein, X chromosome

- 72 -

785733	-0.7554682	hypothetical protein FLJ12892
32489	-0.7521767	hypothetical protein DKFZp566A1524
70201	-0.7513314	mitochondrial solute carrier
611481	-0.7507264	HMG-box transcription factor TCF-3
1554167	-0.7503289	hypothetical protein FLJ14529
768571	-0.7476813	SRY (sex determining region Y)-box 8
783698	-0.7411614	lipin 1
341759	-0.7393558	lung type-I cell membrane-associated glycoprotein
665384	-0.7388404	KIAA1609 protein
142259	-0.7380151	tumor necrosis factor alpha-inducible cellular protein containing leucine zipper domains; Huntingtin interacting protein L; transcription factor IIIA-interacting protein
810057	-0.7363941	vasoactive intestinal peptide receptor 1
27544	-0.7342174	prominin (mouse)-like 1
739193	-0.7338774	cellular retinoic acid-binding protein 1
813256	-0.7334866	ATP-binding cassette, sub-family B (MDR/TAP), member 1
293964	-0.7275302	butyrophilin, subfamily 3, member A1
272706	-0.7274769	alpha2,3-sialyltransferase
209537	-0.7245778	zinc finger protein 221
67741	-0.7229251	PP2135 protein
823714	-0.7215045	nuclear receptor co-repressor/HDAC3 complex subunit
471196	-0.7192752	integral membrane protein 3
813265	-0.7192289	Homo sapiens mRNA; cDNA DKFZp564H1916 (from clone DKFZp564H1916)
42681	-0.718704	NY-REN-25 antigen
1601845	-0.7178673	Ca <sup>2+</sup> -promoted Ras inactivator
725152	-0.7173869	hypothetical protein DKFZp762A227
22917	-0.7173328	Homo sapiens mRNA; cDNA DKFZp761M0111 (from clone DKFZp761M0111)
703541	-0.7118124	KIAA1858 protein
529843	-0.7117107	ESTs, Moderately similar to JC5238 galactosylceramide-like protein, GCP [H.sapiens]
377461	-0.7093974	caveolin 1, caveolae protein, 22kD
1569187	-0.7091259	heparan sulfate (glucosamine) 3-O-sulfotransferase 4



- 73 -

1456118 -0.7074166 proteasome (prosome, macropain) subunit, beta type, 9 (large multifunctional protease 2)

289760 -0.7058375 ESTs, Highly similar to T00391 hypothetical protein KIAA0612 [H.sapiens]

341774 -0.7044768 major histocompatibility complex, class I, A

75078 -0.7028415 ESTs

505864 -0.7020337 RalGDS-like gene

415191 -0.700303 KIAA0161 gene product

188335 -0.7002579 egf-like module containing, mucin-like, hormone receptor-like sequence 2

132711 -0.6985643 Kruppel-like factor 5 (intestinal)

1864302 -0.6943806 E74-like factor 5 (ets domain transcription factor)

344854 -0.692296 ankyrin repeat domain 3

454970 -0.6853932 DKFZP434G032 protein

79629 -0.6849525

767068 -0.6819032 DKFZP586G1517 protein

378813 -0.679848 secretory leukocyte protease inhibitor (antileukoproteinase)

725076 -0.6798005 5'-nucleotidase (purine), cytosolic type B

491403 -0.6760814 tumor necrosis factor receptor superfamily, member 1B

43764 -0.6743049 hypothetical protein FLJ14033 similar to hypoxia inducible factor 3, alpha subunit

725622 -0.6731681 ESTs

431231 -0.6692022 EGF-containing fibulin-like extracellular matrix protein 2

488956 -0.6682156 CUG triplet repeat, RNA-binding protein 2

72778 -0.6614808 caspase 7, apoptosis-related cysteine protease

360254 -0.6612871 cysteine-rich, angiogenic inducer, 61

76182 -0.6603322 hypothetical protein DKFZp761F241

773487 -0.6598647 hypothetical protein MGC3032

162308 -0.6580605 ESTs

283023 -0.6577611 chemokine (C-X3-C) receptor 1

196992 -0.6573113 aldo-keto reductase family 1, member C1 (dihydrodiol dehydrogenase 1; 20-alpha (3-alpha)-hydroxysteroid dehydrogenase)

303109 -0.6571713 purinergic receptor (family A group 5)

897731 -0.6548064 latrophilin

- 74 -

289428 -0.6542789 neurotrophic tyrosine kinase, receptor, type 2

196435 -0.653913 ESTs

504791 -0.6537859 glutathione S-transferase A4

812975 -0.6527728 KIAA0172 protein

488404 -0.6521658 Homo sapiens clone TUA8 Cri-du-chat region mRNA

345056 -0.6490368 KIAA1404 protein

786069 -0.6484842 beta-site APP-cleaving enzyme

48518 -0.6455041 ATP-binding cassette, sub-family A (ABC1), member 5

812048 -0.6438062 prion protein (p27-30) (Creutzfeld-Jakob disease, Gerstmann-Strausler-Scheinker syndrome, fatal familial insomnia)

868396 -0.6426678 tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein, eta polypeptide

70245 -0.6422928 Homo sapiens mRNA full length insert cDNA clone EUROIMAGE 50374

40027 -0.6418674 ESTs

840944 -0.6399619 early growth response 1

809784 -0.6348991 kallikrein 6 (neurosin, zyme)

1881774 -0.6332545 KIAA1678

52419 -0.6326767 Friedreich ataxia region gene X123

120106 -0.631375 caspase 1, apoptosis-related cysteine protease (interleukin 1, beta, convertase)

725680 -0.6295778 transcription factor AP-2 gamma (activating enhancer-binding protein 2 gamma)

1493218 -0.6292715 hypothetical protein FLJ22297

2018423 -0.6275903 death-associated protein kinase 2

1486082 -0.6241275 heparin-binding growth factor binding protein

868169 -0.6224058 lipoprotein lipase

209137 -0.6161588 gamma-aminobutyric acid (GABA) A receptor, epsilon

248886 -0.6161134 rab3 GTPase-activating protein, non-catalytic subunit (150kD)

73252 -0.6115602 3-hydroxy-3-methylglutaryl-Coenzyme A synthase 1 (soluble)

612274 -0.6108452 tubulin, alpha 1 (testis specific)

346545 -0.6098367 laminin, beta 1

84464 -0.6091345 hypothetical protein FLJ12806

- 75 -

628955 -0.6088522 forkhead box O1A (rhabdomyosarcoma)  
490414 -0.6078858 chromosome 2 open reading frame 2  
752837 -0.6056177 Homo sapiens mRNA for FLJ00074 protein, partial cds  
  
796904 -0.604755 pleiomorphic adenoma gene-like 1  
418159 -0.6038708 synaptogyrin 1  
323238 -0.6032171 GRO1 oncogene (melanoma growth stimulating activity, alpha)  
  
1570427 -0.601608 hypothetical protein MGC4309  
1500815 -0.6008126 Homo sapiens cDNA: FLJ22130 fis, clone HEP19632  
  
416959 -0.5999728 nuclear factor I/B  
782460 -0.5988226 Homo sapiens cDNA FLJ10500 fis, clone NT2RP2000369  
  
824602 -0.5965816 interferon, gamma-inducible protein 16  
878798 -0.5956349 beta-2-microglobulin  
32299 -0.5945347 inositol(myo)-1(or 4)-monophosphatase 2  
277571 -0.59335 KIAA1706 protein  
703964 -0.5918604 inositol polyphosphate phosphatase-like 1  
796498 -0.5900328 hypothetical protein FLJ14007  
504927 -0.5894078 epithelial protein up-regulated in carcinoma, membrane associated protein 17  
  
813520 -0.5889772 EphB3  
341763 -0.5868818 caspase 5, apoptosis-related cysteine protease  
  
1856063 -0.5866417 tweety (Drosophila) homolog 1  
825356 -0.5850287 Homo sapiens cDNA FLJ11997 fis, clone HEMBB1001458  
  
745011 -0.5836864 Homo sapiens mRNA for KIAA1750 protein, partial cds  
  
812256 -0.5832398 Homo sapiens cDNA: FLJ21693 fis, clone COL09609  
  
2017960 -0.5822336 PP1201 protein  
713129 -0.5822078 granzyme A (granzyme 1, cytotoxic T-lymphocyte-associated serine esterase 3)  
815142 -0.5782827 spastic ataxia of Charlevoix-Saguenay (sacsin)  
  
80727 -0.5766546 receptor tyrosine kinase-like orphan receptor 1  
  
45099 -0.5753698 regucalcin (senescence marker protein-30)  
  
415816 -0.5750184 ESTs

- 76 -

306798 -0.5742664 NGFI-A binding protein 1 (EGR1 binding protein 1)

563634 -0.5737239 dendritic cell protein

757191 -0.5735888 ESTs

740620 -0.5730915 tropomyosin 2 (beta)

1470048 -0.5729078 lymphocyte antigen 6 complex, locus E

788518 -0.5724497 peroxisomal membrane protein 3 (35kD, Zellweger syndrome)

177665 -0.5718286

221846 -0.5714352 checkpoint suppressor 1

37671 -0.5712258 hypothetical protein FLJ21610

2072862 -0.5694163 v-akt murine thymoma viral oncogene homolog 2

132637 -0.5694152 grancalcin, EF-hand calcium-binding protein

1474900 -0.5693091 keratin 15

34093 -0.5692081 EST

1881689 -0.568518 hypothetical protein FLJ20281

1883327 -0.5684295 ESTs

1902764 -0.5683052 6-phosphofructo-2-kinase/fructose-2,6-biphosphatase 3

781017 -0.5676944 early growth response 2 (Krox-20 (Drosophila) homolog)

34150 -0.5673709 ESTs

137602 -0.5667198 Homo sapiens mRNA; cDNA DKFZp434G0972 (from clone DKFZp434G0972)

460126 -0.5652362 KIAA0819 protein

470128 -0.5652125 myosin IE

1880885 -0.5638973 ESTs

811848 -0.5637994 hypothetical protein

1417886 -0.563564 hypothetical protein FLJ23239

756847 -0.5632582 deformed epidermal autoregulatory factor 1 (Drosophila)

505243 -0.5616577 inositol 1,4,5-triphosphate receptor, type 2

840677 -0.5611985 immunoglobulin kappa constant

1911663 -0.5610314 ESTs

66491 -0.5579872 plasmolipin

1420842 -0.5574834 mucosa associated lymphoid tissue lymphoma translocation gene 1

1626996 -0.5573507 c-fos induced growth factor (vascular endothelial growth factor D)

- 77 -

281190	-0.5572026	ESTs
796475	-0.5563457	four and a half LIM domains 3
1870305	-0.5558155	heat shock 27kD protein 2
767456	-0.5555493	hypothetical protein FLJ22167
292388	-0.554297	
795178	-0.5532479	lactate dehydrogenase C
809998	-0.5519459	amylase, alpha 2A; pancreatic
1734754	-0.5512835	hypothetical protein B
27769	-0.5501268	ESTs
950355	-0.550114	ESTs, Weakly similar to S13495 pregnancy zone protein [H.sapiens]
144924	-0.5498123	chromosome 6 open reading frame 5
502436	-0.5489864	insulin receptor substrate 2
138672	-0.5483104	ESTs
160192	-0.5482009	ESTs, Weakly similar to 2004399A chromosomal protein [H.sapiens]
768007	-0.547374	hypothetical protein
130201	-0.5468031	intercellular adhesion molecule 2
1587847	-0.5466442	minichromosome maintenance deficient (mis5, S. pombe) 6
1558212	-0.5458261	ESTs
1470278	-0.5449844	hypothetical protein FLJ21841
811088	-0.5445374	ephrin-B3
197056	-0.5440514	ESTs
868630	-0.5426831	transforming growth factor beta-stimulated protein TSC-22
530958	-0.5423532	smoothened (Drosophila) homolog
810097	-0.5423072	Homo sapiens cDNA: FLJ21721 fis, clone COLF0381
207735	-0.5422739	serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1
79254	-0.541673	MHC class I region ORF
85634	-0.5410684	complement component 1, s subcomponent
531319	-0.5404527	serine/threonine kinase 12
754028	-0.5396001	KIAA0469 gene product
470148	-0.5393193	ESTs
35828	-0.5391332	diphtheria toxin receptor (heparin-binding epidermal growth factor-like growth factor)

- 78 -

970590 -0.5383645 Homo sapiens mRNA; cDNA DKFZp434A115 (from clone DKFZp434A115)

753162 -0.5377767 KIAA0603 gene product

241066 -0.5369493 Homo sapiens cDNA: FLJ21028 fis, clone CAE07155

1909935 -0.5368632 ESTs

153760 -0.5361176 EphB1

247616 -0.5358709 lipoma HMGIC fusion partner

1031640 -0.5354869 ESTs

24958 -0.5349519 Homo sapiens mRNA; cDNA DKFZp434C2016 (from clone DKFZp434C2016)

1686766 -0.5342051 Rag D protein

2009491 -0.5334821 cyclin-E binding protein 1

611532 -0.5320107 troponin I, skeletal, fast

811149 -0.5319742 chromosome 9 open reading frame 3

844703 -0.5309356 Sam68-like phosphotyrosine protein, T-STAR

70749 -0.5295091 Homo sapiens cDNA: FLJ21874 fis, clone HEP02488

324927 -0.5284875 KIAA0375 gene product

491519 -0.5282605 Homo sapiens clone 24775 mRNA sequence

754040 -0.5282144 NS1-associated protein 1

767765 -0.5282008 GTP-binding protein overexpressed in skeletal muscle

2028722 -0.5277252 osteoblast specific factor 2 (fascin I-like)

509458 -0.5266016 hypothetical protein from clone 643

360518 -0.5253745 guanine nucleotide binding protein (G protein), gamma transducing activity polypeptide 1

40021 -0.5240809 hairless protein (putative single zinc finger transcription factor protein, responsible for autosomal recessive universal congenital alopecia, HR gene)

1901310 -0.5229967 KIAA1209 protein

33949 -0.5222135 phosphoribosyl pyrophosphate synthetase-associated protein 1

814316 -0.521831 ribosomal protein L13

488130 -0.5201845 Homo sapiens cDNA FLJ20767 fis, clone COL06986

462939 -0.5200097 ESTs

- 79 -

770848 -0.5188717 ESTs, Weakly similar to ALU1\_HUMAN ALU SUBFAMILY J  
SEQUENCE CONTAMINATION WARNING ENTRY [H.sapiens]

245296 -0.5184238 RAD52 (S. cerevisiae) homolog  
753034 -0.5182019 nuclear factor I/X (CCAAT-binding transcription factor)

712829 -0.5181236 LIM domain only 2 (rhombotin-like 1)  
1577920 -0.5161377 ESTs  
814526 -0.5156162 seb4D  
773301 -0.5147295 cadherin 3, type 1, P-cadherin (placental)  
77972 -0.5144916 complement component 3  
1704155 -0.5133015 hypothetical protein SBBI48  
1572298 -0.5126426 CD3Z antigen, zeta polypeptide (TiT3 complex)

755689 -0.5122586 retinoic acid receptor, gamma  
1876217 -0.5108977 DnaJ (Hsp40) homolog, subfamily A, member 2

712401 -0.5107426 phosphoinositide-3-kinase, catalytic, delta polypeptide

713839 -0.5088558 transcription factor AP-4 (activating enhancer-binding protein 4)

180803 -0.5083354 inositol polyphosphate-1-phosphatase  
814443 -0.5075293 hypothetical protein MGC3232

Example V: Genes for discriminating between normal and DCIS

As shown in Table 4 below, 1300 genes were identified as being able to discriminate between normal and DCIS.

**Table 4**

<b>CloneID</b>	<b>Weight</b>	<b>Description</b>
1323448	1.397391	cysteine-rich protein 1 (intestinal)
788654	1.2615322	growth factor receptor-bound protein 2
1505038	1.1904802	hypothetical protein FLJ20171
745606	1.1825585	hypothetical protein PP591

- 80 -

565319	1.0969429	Homo sapiens mRNA; cDNA DKFZp564B1264 (from clone DKFZp564B1264)
35147	1.0886084	ESTs, Weakly similar to unnamed protein product [H.sapiens]
178805	1.0274897	Human DNA sequence from clone RP5-850E9 on chromosome 20. Contains part of the gene for a novel C2H2 type zinc finger protein similar to Drosophila Scratch (Scrt), Slug and Xenopus Snail, a novel gene similar to Drosophila CG6762, STSs, GSSs and five CpG
796469	1.0269115	HSPC150 protein similar to ubiquitin-conjugating enzyme
1609836	1.0164283	glutamate-ammonia ligase (glutamine synthase)
366132	1.013995	succinate dehydrogenase complex, subunit C, integral membrane protein, 15kD
1500000	1.0118809	H2B histone family, member B
812238	1.0043787	hypothetical protein MGC4692
595037	0.990461	retinoic acid induced 3
1554549	0.9764206	hydroxyacyl glutathione hydrolase
488964	0.9569176	H2A histone family, member O
138189	0.9386802	Wolfram syndrome 1 (wolframin)
283919	0.937383	H2A histone family, member L
823598	0.9153521	proteasome (prosome, macropain) 26S subunit, non-ATPase, 12
810711	0.9128832	stearoyl-CoA desaturase (delta-9-desaturase)
1911343	0.9116745	RAB26, member RAS oncogene family
503215	0.9108688	pilin-like transcription factor
811774	0.9073627	CGI-49 protein
122077	0.9021901	putative membrane protein
1492238	0.9019233	HSPC003 protein
469686	0.8919992	Ric (Drosophila)-like, expressed in many tissues
685516	0.8762873	putative G protein-coupled receptor
359887	0.8692353	translocase of inner mitochondrial membrane 17 (yeast) homolog A
131094	0.8675719	Homo sapiens cDNA: FLJ21587 fis, clone COL06946
1492463	0.8634521	selenoprotein X, 1



- 81 -

843195	0.8603524	phosphoserine phosphatase
796723	0.8573155	Homo sapiens clone CDABP0014 mRNA sequence
488202	0.8463961	ESTs, Weakly similar to YZ28_HUMAN HYPOTHETICAL PROTEIN ZAP128 [H.sapiens]
1492426	0.8445255	chromosome 19 open reading frame 3
280375	0.8392836	PRO2000 protein
290841	0.8346933	H2B histone family, member A
1917941	0.8340565	purine-rich element binding protein B
1469425	0.8296141	SRY (sex determining region Y)-box 22
2016908	0.8282806	ESTs, Weakly similar to CA13_HUMAN COLLAGEN ALPHA 1(III) CHAIN PRECURSOR [H.sapiens]
471568	0.8264546	hematological and neurological expressed 1
150003	0.8251408	hypothetical protein FLJ13187
2029173	0.8213721	ESTs, Weakly similar to N-WASP [H.sapiens]
731044	0.8169506	glutaredoxin 2
244801	0.8066357	Rho guanine exchange factor (GEF) 11
2054635	0.7914755	proteasome (prosome, macropain) subunit, alpha type, 7
1605426	0.787298	hypothetical protein FLJ13352
1709791	0.7837549	BAI1-associated protein 1
470061	0.77977	seven in absentia (Drosophila) homolog 2
1640821	0.7790286	ESTs, Weakly similar to I78885 serine/threonine-specific protein kinase [H.sapiens]
199403	0.7754748	lectin, galactoside-binding, soluble, 8 (galectin 8)
741474	0.7732047	glucose phosphate isomerase
1435862	0.7697423	antigen identified by monoclonal antibodies 12E7, F21 and O13
239568	0.767277	annexin A9
841679	0.7666686	calcium and integrin binding protein (DNA-dependent protein kinase interacting protein)
729975	0.7649717	meningioma expressed antigen 5 (hyaluronidase)
2016648	0.7633048	Homo sapiens mRNA; cDNA DKFZp434N1728 (from clone DKFZp434N1728)
2052113	0.7627076	hypothetical protein FLJ10903

sd-71385

T0231-BF032001

- 82 -

347373	0.7563599	transcription elongation factor B (SIII), polypeptide 1 (15kD, elongin C)
814054	0.7539407	KIAA0040 gene product
741977	0.7538359	B-factor, properdin
272529	0.7486269	phosphomannomutase 2
824879	0.7432221	hypothetical protein MGC11275
122241	0.7394284	proteasome (prosome, macropain) subunit, beta type, 2
744417	0.7356767	carnitine acetyltransferase
868128	0.7347171	JM4 protein
241348	0.7324115	prenylcysteine lyase
509823	0.7304507	carcinoembryonic antigen-related cell adhesion molecule 6 (non-specific cross reacting antigen)
1636092	0.7270017	hypothetical protein FLJ20657
42408	0.725109	hypothetical protein MGC4604
686552	0.7231319	golgi phosphoprotein 1
898032	0.7216	KIAA0097 gene product
754628	0.7195265	ESTs
826256	0.7190206	transmembrane 7 superfamily member 1 (upregulated in kidney)
2043167	0.7167431	BCL2-associated athanogene 3
827171	0.7154964	ESTs
502774	0.7148695	hypothetical protein FLJ20623
781097	0.7126945	reticulon 3
810725	0.7115051	ATPase, H <sup>+</sup> transporting, lysosomal (vacuolar proton pump) 21kD
768570	0.7102463	hypothetical protein FLJ11280
813281	0.7092607	WW domain-containing protein 1
39884	0.7075527	IMP (inosine monophosphate) dehydrogenase 1
969877	0.707469	synaptosomal-associated protein, 25kD
704254	0.7061648	a disintegrin and metalloproteinase domain 8
236034	0.7039329	uncoupling protein 2 (mitochondrial, proton carrier)
782428	0.701077	KIAA0250 gene product
489351	0.7003616	hypothetical protein DKFZp566J2046
37708	0.6986007	hypothetical protein MGC3101
271472	0.6978431	C3HC4-like zinc finger protein
782608	0.6978285	mitochondrial ribosomal protein L9

- 83 -

704414	0.6940045	small nuclear ribonucleoprotein polypeptides B and B1
288999	0.6938843	small protein effector 1 of Cdc42
51773	0.6925666	hypothetical protein MGC3077
124447	0.6918546	KIAA1184 protein
470099	0.6915952	HT002 protein; hypertension-related calcium-regulated gene
429799	0.6915155	hypothetical protein FLJ21939 similar to 5-azacytidine induced gene 2
1858892	0.6904662	hypothetical protein MGC4825
796694	0.6893307	baculoviral IAP repeat-containing 5 (survivin)
284261	0.6890531	uncharacterized hematopoietic stem/progenitor cells protein MDS030
814528	0.6878182	Homo sapiens cDNA: FLJ22139 fis, clone HEP20959
788654	0.6866898	
68636	0.6838604	hypothetical protein MGC2477
308466	0.6800103	GTP-binding protein Sara
1574058	0.6796357	1-acylglycerol-3-phosphate O-acyltransferase 2 (lysophosphatidic acid acyltransferase, beta)
79520	0.6769129	RAB2, member RAS oncogene family
1473289	0.674396	protective protein for beta-galactosidase (galactosialidosis)
365738	0.6740773	ESTs
768064	0.673441	cytochrome P450, subfamily I (aromatic compound-inducible), polypeptide 1
773922	0.6730903	KIAA0005 gene product
1869201	0.6727191	hypothetical protein MGC2745
686172	0.6724684	KIAA0008 gene product
810558	0.6698194	proteasome (prosome, macropain) 26S subunit, ATPase, 4
753299	0.6694573	hypothetical protein FLJ10504
725454	0.6669973	CDC28 protein kinase 2
470124	0.6641768	RAD1 (S. pombe) homolog
2014034	0.6620303	methylene tetrahydrofolate dehydrogenase (NAD+ dependent), methenyltetrahydrofolate cyclohydrolase
209066	0.6611038	
859761	0.6573769	poliovirus receptor-related 2 (herpesvirus entry mediator B)
589232	0.6551447	hypothetical protein FLJ11506

- 84 -

340558	0.6550804	actin related protein 2/3 complex, subunit 5 (16 kD)
186768	0.6550741	Homo sapiens, clone IMAGE:3604869, mRNA
813629	0.6549767	YME1 (S.cerevisiae)-like 1
1492780	0.6544659	Homo sapiens cDNA FLJ14459 fis, clone HEMBB1002409
809944	0.6530236	KIAA0310 gene product
83363	0.6526164	protein-L-isoaspartate (D-aspartate) O-methyltransferase
149355	0.6520333	translocating chain-associating membrane protein
752631	0.650399	fibroblast growth factor receptor 3 (achondroplasia, thanatophoric dwarfism)
785616	0.6484744	signal sequence receptor, alpha (translocon-associated protein alpha)
76605	0.6484522	nesca protein
742707	0.6476434	ESTs, Weakly similar to MUC2_HUMAN MUCIN 2 PRECURSOR [H.sapiens]
325606	0.647638	hypothetical protein MGC14353
769921	0.6464245	ubiquitin carrier protein E2-C
1435003	0.6457486	tumor suppressing subtransferable candidate 1
1473922	0.6450042	actin related protein 2/3 complex, subunit 3 (21 kD)
1456348	0.6417276	N-acetylneuraminic acid phosphate synthase; sialic acid synthase
1500162	0.6416738	ESTs
1573251	0.6410435	peroxisomal long-chain acyl-coA thioesterase
46248	0.640944	ADP-ribosyltransferase (NAD+; poly (ADP-ribose) polymerase)
49351	0.6386132	SEX gene
824052	0.6375787	chromosome 6 open reading frame 1
1420370	0.6344209	biliverdin reductase B (flavin reductase (NADPH))
842994	0.6343858	cathepsin Z
810612	0.6341098	S100 calcium-binding protein A11 (calgizzarin)
123614	0.633308	hypothetical protein MGC4675
124781	0.6324418	squalene epoxidase
814378	0.6317382	serine protease inhibitor, Kunitz type, 2
2108077	0.6312583	CGI-112 protein

sd-71385

10026018-100401

- 85 -

289978	0.6308861	ubiquitin-like 4
67765	0.6300886	carboxypeptidase M
742595	0.6265846	cyclin-dependent kinase 5
727078	0.6263882	Homo sapiens cDNA: FLJ23602 fis, clone LNG15735
487733	0.6262778	
366067	0.625758	cerebellar degeneration-related protein (62kD)
292770	0.6231261	Homo sapiens, clone IMAGE:3627860, mRNA, partial cds
366353	0.6203012	DKFZP564C186 protein
810063	0.620086	growth factor, erv1 (S. cerevisiae)-like (augmenter of liver regeneration)
810124	0.6162712	platelet-activating factor acetylhydrolase, isoform Ib, gamma subunit (29kD)
713782	0.6149877	a disintegrin and metalloproteinase domain 15 (metargidin)
782608	0.6137782	
530197	0.6122721	ESTs, Moderately similar to ALU8_HUMAN ALU SUBFAMILY SX SEQUENCE CONTAMINATION WARNING ENTRY [H.sapiens]
44292	0.6122043	Homo sapiens mRNA; cDNA DKFZp434C107 (from clone DKFZp434C107)
2018084	0.6119069	Ste-20 related kinase
789012	0.6104002	fibulin 2
256619	0.6102239	hydroxysteroid (17-beta) dehydrogenase 7
825327	0.6094796	Homo sapiens cDNA FLJ14105 fis, clone MAMMA1001202
842980	0.6081528	developmentally regulated GTP-binding protein 1
811761	0.6045809	Nijmegen breakage syndrome 1 (nibrin)
1601947	0.6045149	cytochrome c oxidase subunit VIIa polypeptide 2 (liver)
1474955	0.6045103	TATA box binding protein (TBP)-associated factor, RNA polymerase II, N, 68kD (RNA-binding protein 56)
1758590	0.6040962	fatty-acid-Coenzyme A ligase, long-chain 3
1456701	0.6038789	B-cell CLL/lymphoma 9
207288	0.603346	insulin induced gene 1
1631699	0.6027911	valosin-containing protein

sd-71385

"1008018" 1008018

- 86 -

1635681	0.6022549	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 2 (8kD, B8)
768452	0.6018854	Homo sapiens EST from clone 491476, full insert
897806	0.6015789	hypoxia-inducible factor 1, alpha subunit (basic helix-loop-helix transcription factor)
108425	0.6008772	
770785	0.600311	1,2-alpha-mannosidase IC
2110511	0.5987863	artemin
785795	0.598504	hypothetical protein FLJ12910
120271	0.5982507	hypothetical protein MGC4692
248649	0.5975539	hypothetical protein FLJ13910
2244196	0.5972111	B-cell receptor-associated protein BAP29
66406	0.5966701	hypothetical protein DKFZp762E1312
1537001	0.59528	ESTs
2028949	0.5946445	hypothetical protein PRO1855
40173	0.5940582	KIAA0807 protein
490778	0.5932928	low molecular mass ubiquinone-binding protein (9.5kD)
503889	0.5930018	hypothetical protein FLJ10656
781019	0.5918462	paraoxonase 2
196189	0.5916252	cytochrome b-5
1734309	0.5894492	sperm associated antigen 4
814350	0.588877	insulin-degrading enzyme
2018821	0.5887114	ATPase inhibitor precursor
365358	0.5883524	pM5 protein
430235	0.5874504	H2B histone family, member Q
1422338	0.5874097	ribonucleotide reductase M2 polypeptide
770845	0.5870877	hexokinase 1
172517	0.5869599	hippocalcin-like 1
1884404	0.5849803	KIAA0285 gene product
625923	0.5839066	phosphoenolpyruvate carboxykinase 2 (mitochondrial)
2322367	0.5832711	reticulon 4
624667	0.5828923	CGI-92 protein
826363	0.5823873	lysophospholipase II
859228	0.5817344	isocitrate dehydrogenase 1 (NADP+), soluble
2302099	0.5813788	sialidase 3 (membrane sialidase)
209066	0.5794211	serine/threonine kinase 15

- 87 -

1844765	0.57783	Homo sapiens mRNA; cDNA DKFZp564O1763 (from clone DKFZp564O1763)
191904	0.5770884	BUB3 (budding uninhibited by benzimidazoles 3, yeast) homolog
951233	0.5767874	proteasome (prosome, macropain) subunit, beta type, 3
809357	0.5767382	Bernardinelli-Seip congenital lipodystrophy 2 (seipin)
825740	0.5762093	DKFZp434J1813 protein
51532	0.5758765	ADP-ribosylation factor-like 6 interacting protein
203003	0.5743416	non-metastatic cells 4, protein expressed in
2015517	0.5742448	hypothetical protein FLJ22237
205049	0.5734444	protein kinase H11; small stress protein-like protein HSP22
752643	0.5732323	group XII secreted phospholipase A2
753400	0.5732322	CGI-204 protein
564847	0.573202	ESTs
785766	0.5731938	hypothetical protein
825585	0.5722829	tubulin-specific chaperone e
37554	0.5706008	hypothetical protein FLJ22353
144880	0.5704416	hypothetical protein from EUROIMAGE 1759349
49273	0.5691181	solute carrier family 27 (fatty acid transporter), member 4
743589	0.568516	ESTs, Weakly similar to T2D3_HUMAN TRANSCRIPTION INITIATION FACTOR TFIID 135 KDA SUBUNIT [H.sapiens]
1616253	0.567837	breast carcinoma amplified sequence 1
1581941	0.5676045	hypothetical protein FLJ14540
431505	0.5670119	ESTs, Highly similar to A31026 probable membrane receptor protein [H.sapiens]
2019223	0.5663073	mitochondrial ribosomal protein L17
840878	0.5648009	seladin-1
811024	0.5646461	bone marrow stromal cell antigen 2
564492	0.5626905	mitochondrial carrier homolog 2
725841	0.561197	KIAA0662 gene product
701751	0.561139	cut (Drosophila)-like 1 (CCAAT displacement protein)
767798	0.560181	ATX1 (antioxidant protein 1, yeast) homolog 1

- 88 -

109863	0.559767	epithelial membrane protein 2
1536006	0.5590182	ESTs
1845169	0.5588842	RAB35, member RAS oncogene family
343607	0.5588222	AD-015 protein
212542	0.5587394	Homo sapiens cDNA FLJ12900 fis, clone NT2RP2004321
1700436	0.5569949	ESTs
795256	0.5567908	NPD007 protein
2011515	0.5566484	DKFZP586B0923 protein
504308	0.5564295	hypothetical protein FLJ10540
772925	0.5562164	HSPCO34 protein
469383	0.5558732	chromosome 8 open reading frame 1
810402	0.5555749	hypothetical protein
1409509	0.554703	troponin T1, skeletal, slow
2050827	0.5544641	proteasome (prosome, macropain) 26S subunit, ATPase, 5
770355	0.5518948	lanosterol synthase (2,3-oxidosqualene-lanosterol cyclase)
813410	0.5513568	polymerase (RNA) II (DNA directed) polypeptide K (7.0kD)
80764	0.5510626	hypothetical protein
1631735	0.5502891	Homo sapiens, clone IMAGE:3604336, mRNA, partial cds
625234	0.550211	KDEL (Lys-Asp-Glu-Leu) endoplasmic reticulum protein retention receptor 3
279970	0.5502099	adenosine A2a receptor
1518402	0.5497179	KIAA1361 protein
753378	0.548977	hypothetical protein FLJ22649 similar to signal peptidase SPC22/23
594500	0.5474088	EST
590774	0.5460342	mitogen-activated protein kinase 13
2055807	0.5460123	protein kinase domains containing protein similar to phosphoprotein C8FW
773188	0.5454398	nuclear receptor subfamily 1, group D, member 2
126851	0.5454014	hypothetical protein FLJ11160
244764	0.5450377	B7 homolog 3
813419	0.5447006	hydroxyacyl-Coenzyme A dehydrogenase, type II
233349	0.5446353	hypothetical protein FLJ10761
785701	0.5446211	RAB31, member RAS oncogene family
268946	0.5440003	WD40 protein Ciao1



- 89 -

611443 0.5432426 myoglobin  
510575 0.5430211 hypothetical protein FLJ22087  
491465 0.5420241 hypothetical protein FLJ10035  
1393018 0.5404243 general transcription factor IIIC, polypeptide 1 (alpha subunit,  
220kD )  
785840 0.5401209 SEC24 (S. cerevisiae) related gene family, member D  
1460110 0.5392177 proteasome (prosome, macropain) subunit, beta type, 5  
731308 0.5371204 citrate synthase  
301122 0.5368073 extracellular matrix protein 1  
66535 0.5363248 ornithine decarboxylase antizyme 2  
265103 0.5361177 Homo sapiens mRNA; cDNA DKFZp547M123 (from clone  
DKFZp547M123)  
788334 0.5347204 mitochondrial ribosomal protein L23  
813631 0.534683 seven transmembrane protein TM7SF3  
1553306 0.5344506 proteasome (prosome, macropain) 26S subunit, non-ATPase,  
11  
1461477 0.5343834 Homo sapiens mRNA; cDNA DKFZp586I0324 (from clone  
DKFZp586I0324)  
199645 0.5334951 nicastrin  
884425 0.5327862 chaperonin containing TCP1, subunit 5 (epsilon)  
1518890 0.5314577 metallothionein-like 5, testis-specific (tesmin)  
756442 0.5313744 P450 (cytochrome) oxidoreductase  
2326057 0.5310995 MLN51 protein  
138788 0.5302936 prolactin receptor  
1698036 0.5298397 ubiquitin-conjugating enzyme E2 variant 1  
811585 0.5288102 huntingtin (Huntington disease)  
795185 0.5276861 xenotropic and polytropic retrovirus receptor  
629944 0.5273447 myosin VB  
784105 0.5259815 ESTs  
1696757 0.5254296 hypothetical protein KIAA1165  
139835 0.5253386 UDP-glucose dehydrogenase  
1738208 0.5240868 a disintegrin-like and metalloprotease (repolysin type) with  
thrombospondin type 1 motif, 4

- 90 -

593431	0.5235073	ESTs, Moderately similar to CEGT_HUMAN CERAMIDE GLUCOSYLTRANSFERASE [H.sapiens]
2309073	0.5226599	frizzled (Drosophila) homolog 5
825470	0.5218731	topoisomerase (DNA) II alpha (170kD)
839682	0.5208954	ubiquitin-conjugating enzyme E2N (homologous to yeast UBC13)
1517171	0.5195255	interleukin 2 receptor, alpha
376516	0.5195127	cell division cycle 4-like
774446	0.5181672	adrenomedullin
1476053	0.5179015	RAD51 (S. cerevisiae) homolog (E coli RecA homolog)
489594	0.5178551	hypothetical protein FLJ11565
1573946	0.5173691	programmed cell death 9
503851	0.5170168	nuclear receptor co-repressor/HDAC3 complex subunit
882484	0.5165756	chaperonin containing TCP1, subunit 7 (eta)
51083	0.5162686	catenin (cadherin-associated protein), delta 2 (neural plakophilin-related arm-repeat protein)
221295	0.5157877	inhibitor of DNA binding 2, dominant negative helix-loop-helix protein
725978	0.5147267	ESTs, Moderately similar to ALU7_HUMAN ALU SUBFAMILY SQ SEQUENCE CONTAMINATION WARNING ENTRY [H.sapiens]
796757	0.5122331	adaptor-related protein complex 3, sigma 1 subunit
150314	0.5114988	lysophospholipase I
302031	0.5111067	Ste20-related serine/threonine kinase
293569	0.5103638	chromosome 1 open reading frame 21
838366	0.5072623	3-hydroxymethyl-3-methylglutaryl-Coenzyme A lyase (hydroxymethylglutaricaciduria)
122147	0.5067128	
120749	0.5060444	ESTs, Moderately similar to KIAA1215 protein [H.sapiens]
855563	0.5059683	v-erb-b2 avian erythroblastic leukemia viral oncogene homolog 3
487733	0.5057167	NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 2 (8kD, AGGG)
625693	0.5055986	hypothetical protein MGC10911

sd-71385

1003019 BT03201

- 91 -

813751	0.5051993	sialyltransferase 4C (beta-galactosidase alpha-2,3-sialyltransferase)
842825	0.5047729	G1 to S phase transition 1
323693	0.5047237	adaptor-related protein complex 1, sigma 1 subunit
202514	0.5045411	DNA (cytosine-5-)-methyltransferase 3 alpha
1517749	0.5045134	ESTs
124331	0.5025464	cleavage and polyadenylation specific factor 5, 25 kD subunit
1474424	0.5021308	Homo sapiens cDNA FLJ12758 fis, clone NT2RP2001328
1848977	0.5015371	glycerol kinase
454896	0.5013449	DnaJ (Hsp40) homolog, subfamily A, member 2
1702742	0.5011725	solute carrier family 7 (cationic amino acid transporter, y+ system), member 5
746163	0.5009715	ESTs, Weakly similar to ALU1_HUMAN ALU SUBFAMILY J SEQUENCE CONTAMINATION WARNING ENTRY [H.sapiens]
810734	0.4998276	polymerase (DNA-directed), delta 4
725284	0.4992842	phosphorylase kinase, gamma 2 (testis)
52103	0.4988158	hypothetical protein FLJ23045
739109	0.4986821	adaptor-related protein complex 2, sigma 1 subunit
488505	0.4981852	accessory proteins BAP31/BAP29
365060	0.4976091	RAB11A, member RAS oncogene family
41569	0.4974126	hypothetical protein FLJ12650
81336	0.4955353	uteroglobin
700792	0.4949149	cyclin-dependent kinase inhibitor 3 (CDK2-associated dual specificity phosphatase)
1592715	0.4944665	Homer, neuronal immediate early gene, 3
142586	0.4942961	MCT-1 protein
261472	0.4941066	putative nuclear protein ORF1-FL49
246800	0.4927231	hypothetical protein FLJ10803
124298	0.4922937	microsomal glutathione S-transferase 3
1898619	0.4916939	hypothetical protein MGC15737
739126	0.4905859	tissue specific transplantation antigen P35B
376875	0.490328	flavin containing monooxygenase 1
358162	0.4901786	protein predicted by clone 23627
590759	0.4895021	sterol-C4-methyl oxidase-like

sd-71385

1002304-12101  
"BTF032001"

- 92 -

1947647	0.4887766	CGI-147 protein
120271	0.4886584	hypothetical protein MGC4692
488642	0.4881613	ESTs, Weakly similar to I38022 hypothetical protein [H.sapiens]
1492468	0.4878466	DEME-6 protein
345538	0.4875796	cathepsin L
344091	0.4871044	ESTs
123441	0.4862068	ribosomal protein L7a
202901	0.48608	vav 2 oncogene
757489	0.4859039	tubulin, alpha 2
49117	0.4856399	KIAA0215 gene product
812994	0.4854155	retinoid X receptor, alpha
731023	0.4853217	WD repeat domain 5
814899	0.4840969	BCL2/adenovirus E1B 19kD-interacting protein 3-like
756666	0.4835569	protein phosphatase 1, catalytic subunit, alpha isoform
2018337	0.4833207	glucosidase, beta; acid (includes glucosylceramidase)
1568825	0.4829928	Arg/Abl-interacting protein ArgBP2
869375	0.4827093	isocitrate dehydrogenase 2 (NADP+), mitochondrial
595213	0.4824381	hypothetical protein
744374	0.4824256	putative ankyrin-repeat containing protein
73009	0.4822587	ESTs, Weakly similar to A43932 mucin 2 precursor, intestinal [H.sapiens]
70606	0.481356	ESTs
358456	0.4812268	Sec61 gamma
810762	0.4806851	SNARE protein
823930	0.4800704	actin related protein 2/3 complex, subunit 1A (41 kD)
2020898	0.4797032	procollagen-lysine, 2-oxoglutarate 5-dioxygenase 3
564981	0.4782207	ESTs
1637282	0.4777562	hexokinase 2
855749	0.476968	triosephosphate isomerase 1
744944	0.4769116	myosin VI
263727	0.4767486	DNA segment, single copy probe LNS-CAI/LNS-CAII (deleted in polyposis)
470092	0.476018	like-glycosyltransferase
782513	0.4759571	interferon, alpha-inducible protein (clone IFI-6-16)



- 94 -

345787 0.4647372 highly expressed in cancer, rich in leucine heptad repeats  
825296 0.4646099 low density lipoprotein receptor defect C complementing  
2248488 0.4645687 ems1 sequence (mammary tumor and squamous cell  
carcinoma-associated (p80/85 src substrate)  
824962 0.4638758 karyopherin alpha 2 (RAG cohort 1, importin alpha 1)  
843054 0.4637278 KIAA1533 protein  
140289 0.4624364 hypothetical protein  
1558233 0.4622449 ESTs  
265592 0.4615543 activated RNA polymerase II transcription cofactor 4  
767761 0.4608709 DKFZP434B168 protein  
770766 0.4603527 DKFZP564C1940 protein  
814209 0.4602081 ESTs  
773674 0.4601667 oncogene TC21  
703707 0.4599434 aspartate beta-hydroxylase  
951216 0.4594172 NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 10  
(22kD, PDSW)  
139354 0.4582974 hypothetical protein  
79710 0.4579008 KIAA0174 gene product  
823574 0.457224 endosulfine alpha  
845363 0.4570873 non-metastatic cells 1, protein (NM23A) expressed in  
283751 0.4567022 cortistatin  
810156 0.4561539 deoxythymidylate kinase (thymidylate kinase)  
1416782 0.4558415 creatine kinase, brain  
810609 0.4551167 hypothetical protein PP1226  
770675 0.454729 Homo sapiens cDNA: FLJ21323 fis, clone COL02374  
626318 0.4529715 ubinuclein 1  
38356 0.4513099 cytochrome P450, subfamily 46 (cholesterol 24-hydroxylase)  
705064 0.4497555 transforming, acidic coiled-coil containing protein 3  
127646 0.4493747  
299815 0.4482635 hypothetical protein DC42  
815026 0.4472842 cleavage stimulation factor, 3' pre-RNA, subunit 3, 77kD

- 95 -

489823	0.4468978	COX17 (yeast) homolog, cytochrome c oxidase assembly protein
469151	0.4462659	eukaryotic translation initiation factor 2, subunit 2 (beta, 38kD )
41826	0.4456463	ESTs
47853	0.445149	aldehyde dehydrogenase 4 family, member A1
144740	0.4449229	similar to phosphatidylcholine transfer protein 2
726658	0.4446657	non-metastatic cells 3, protein expressed in
325515	0.444486	hypothetical protein FLJ10980
815781	0.4441245	heat shock 105kD
746190	0.4430243	hypothetical protein DKFZp761B1514
2016194	0.4417536	glia maturation factor, beta
1584287	0.4413331	KIAA0241 protein
147834	0.4412795	zinc finger protein 217
154610	0.4410751	dynactin 4
754653	0.4405873	cadherin, EGF LAG seven-pass G-type receptor 3, flamingo (Drosophila) homolog
742581	0.440367	Homo sapiens cDNA FLJ10366 fis, clone NT2RM2001420
2306987	0.4402912	secreted and transmembrane 1
878406	0.4396935	metaxin 1
430614	0.438809	2,3-bisphosphoglycerate mutase
415102	0.4387892	cell division cycle 25C
358217	0.4385343	glypican 4
1636669	0.4384707	hypothetical protein FLJ21939 similar to 5-azacytidine induced gene 2
83358	0.4380365	ESTs
1325816	0.4379227	polymerase (RNA) II (DNA directed) polypeptide L (7.6kD)
824524	0.4375695	UDP-galactose transporter related
795805	0.4370764	KIAA0332 protein
743977	0.4368155	Homo sapiens mRNA for TL132
785793	0.4367093	capping protein (actin filament) muscle Z-line, alpha 1
266218	0.4362517	hypothetical protein FLJ11350
769600	0.4362014	uracil-DNA glycosylase 2
1656062	0.4355127	coagulation factor XII (Hageman factor)
2017415	0.43546	centromere protein A (17kD)
772220	0.4352917	for protein disulfide isomerase-related

- 96 -

811907	0.4340842	hypothetical protein FLJ22056
789376	0.4339171	thioredoxin reductase 1
782503	0.4334388	fatty acid desaturase 1
753897	0.4324066	autocrine motility factor receptor
767289	0.4323485	hypothetical protein FLJ10055
1159963	0.4323117	interferon regulatory factor 7
1652310	0.4321289	Homo sapiens, clone MGC:19613 IMAGE:3833049, mRNA, complete cds
358936	0.4314082	ESTs, Weakly similar to T2D3_HUMAN TRANSCRIPTION INITIATION FACTOR TFIID 135 KDA SUBUNIT [H.sapiens]
41698	0.4310061	progesterone binding protein
1635665	0.430087	Homo sapiens, RIKEN cDNA 2010100O12 gene, clone MGC:14813 IMAGE:4133274, mRNA, complete cds
1558642	0.4298376	hypothetical protein MGC2771
1733262	0.4289711	BLu protein
823694	0.4280476	Homo sapiens chromosome 19, BAC CIT-HSPC_204F22 (BC228680), complete sequence; contains bacterial insertion element
1641894	0.4277815	EST
2572170	0.4277651	ESTs, Weakly similar to T26581 hypothetical protein Y32B12A.3 - Caenorhabditis elegans [C.elegans]
731080	0.4276128	hypothetical protein FLJ12661
649084	0.4275738	carbonic anhydrase XI
795498	0.4270889	putative transmembrane protein
897770	0.4270685	
2250839	0.4266248	androgen receptor (dihydrotestosterone receptor; testicular feminization; spinal and bulbar muscular atrophy; Kennedy disease)
510794	0.4264564	c-myc binding protein
825822	0.4255725	hypothetical protein
825659	0.424655	N-myc downstream regulated
826077	0.4242762	pyruvate dehydrogenase (lipoamide) beta
293727	0.4242722	hypothetical protein MGC861
503866	0.4230404	sperm autoantigenic protein 17
491527	0.4219324	Homo sapiens, Similar to CG7083 gene product, clone MGC:10534 IMAGE:3957147, mRNA, complete cds
292936	0.4217115	hypothetical protein FLJ10468

sd-71385

FILED OCT 3 2001



- 97 -

823940	0.4214277	transducer of ERBB2, 1
414992	0.420994	K562 cell-derived leucine-zipper-like protein 1
346134	0.4209894	calcium-regulated heat-stable protein (24kD)
51657	0.4208387	hypothetical protein ET
2009779	0.420743	rabaptin-5
1523225	0.4195302	oncostatin M receptor
826194	0.4177687	synaptotagmin-like 2
85804	0.4171445	hypothetical protein FLJ21918
725223	0.4167519	KIAA0077 protein
79726	0.4162264	ESTs, Highly similar to T46395 hypothetical protein DKFZp434I1120.1 [H.sapiens]
1762111	0.4160469	natriuretic peptide receptor C/guanylate cyclase C (atrionatriuretic peptide receptor C)
509588	0.4157059	TATA box binding protein (TBP)-associated factor, RNA polymerase II, J, 20kD
327506	0.4152106	Homo sapiens mRNA full length insert cDNA clone EUROIMAGE 327506
744047	0.4151137	polo (Drosophia)-like kinase
221499	0.4149929	KIAA0508 protein
745394	0.4147627	Homo sapiens cDNA: FLJ23249 fis, clone COL04196
149539	0.4138942	KIAA1700
823907	0.4135955	hypothetical protein FLJ10511
66317	0.4132405	H1 histone family, member 2
295986	0.413087	emopamil-binding protein (sterol isomerase)
565235	0.4124667	spermine synthase
825386	0.4121622	ATP synthase, H <sup>+</sup> transporting, mitochondrial F1F0, subunit d
250313	0.4121296	ESTs
72050	0.4114235	chloride channel, nucleotide-sensitive, 1A
417801	0.4112944	mitochondrial ribosomal protein L27
841501	0.4111707	KIAA0102 gene product
813707	0.410787	regulator of G-protein signalling 16
745083	0.4107735	ubiquitin specific protease 18
244974	0.4105439	hypothetical protein FLJ22875
26883	0.4103824	protein kinase (cAMP-dependent, catalytic) inhibitor beta

280249	0.4096162	Kruppel-like factor 7 (ubiquitous)
74738	0.409593	Homo sapiens, clone IMAGE:3535294, mRNA, partial cds
1753497	0.409432	ovo (Drosophila) homolog-like 1
204299	0.4073087	replication protein A3 (14kD)
358609	0.4071867	NADH dehydrogenase (ubiquinone) flavoprotein 3 (10kD)
290101	0.4071477	ESTs
2306221	0.4068933	wingless-type MMTV integration site family, member 10B
726439	0.4067151	CGI-143 protein
431805	0.405447	granulin
111362	0.4045961	
472103	0.4043572	soc-2 (suppressor of clear, C.elegans) homolog
788444	0.4042598	KIAA0033 protein
1632252	0.404193	complement component 1, q subcomponent, alpha polypeptide
150118	0.4039789	hypothetical protein DKFZp434F054
594226	0.403676	Homo sapiens cDNA FLJ14459 fis, clone HEMBB1002409
1499828	0.4030396	fucosyltransferase 1 (galactoside 2-alpha-L-fucosyltransferase, Bombay phenotype included)
810939	0.40256	hypothetical protein FLJ22169
841621	0.4020253	ESTs, Weakly similar to JE0350 Anterior gradient-2 [H.sapiens]
200144	0.4019827	ring-box 1
486110	0.4017126	profilin 2
292213	0.4014458	guanine nucleotide binding protein (G protein), beta polypeptide 2
1631132	0.401216	peptide transporter 3
669379	0.4011648	Homo sapiens BAC clone RP11-505D17 from 7p22-p21
773685	0.4009493	phosphodiesterase 4D interacting protein (myomegalin)
810411	0.400096	hypothetical protein FLJ13222
1583198	0.39981	ESTs, Weakly similar to S65824 reverse transcriptase homolog [H.sapiens]
84295	0.3989864	interleukin 1 receptor antagonist
757328	0.3985331	hypothetical protein FLJ22678
35626	0.39822	Homo sapiens cDNA FLJ14201 fis, clone NT2RP3002955

- 99 -

1587863	0.3980812	acetyl-Coenzyme A acyltransferase 1 (peroxisomal 3-oxoacyl-Coenzyme A thiolase)
713862	0.3965573	ubiquitin-protein isopeptide ligase (E3)
343731	0.3962504	
43977	0.3961848	KIAA0182 protein
78869	0.3954821	cell membrane glycoprotein, 110000M(r) (surface antigen)
756595	0.395121	S100 calcium-binding protein A10 (annexin II ligand, calpactin I, light polypeptide (p11))
701115	0.3936994	hypothetical protein PRO2013
81599	0.3932769	ubiquitin specific protease 14 (tRNA-guanine transglycosylase)
1636844	0.3926235	ring finger protein 14
703739	0.392298	nuclear cap binding protein subunit 1, 80kD
279633	0.3907503	ESTs, Highly similar to I38759 zinc finger/leucine zipper protein [H.sapiens]
122091	0.3903482	casein kinase 2, alpha 1 polypeptide
786067	0.390151	cell division cycle 25B
1909574	0.3900357	mitochondrial ribosomal protein S11
263013	0.3894536	procollagen-lysine, 2-oxoglutarate 5-dioxygenase (lysine hydroxylase) 2
491001	0.3884144	glyoxalase I
455275	0.3869874	hypothetical protein FLJ23469
85450	0.3868399	acyl-Coenzyme A oxidase 2, branched chain
773286	0.3861543	solute carrier family 9 (sodium/hydrogen exchanger), isoform 3 regulatory factor 1
366156	0.3861394	Homo sapiens cDNA FLJ14028 fis, clone HEMBA1003838
49630	0.3858335	calcium channel, voltage-dependent, L type, alpha 1D subunit
121251	0.385792	hypothetical protein MGC5576
2322223	0.3853156	small nuclear ribonucleoprotein polypeptide A
276915	0.3851081	DNA (cytosine-5-)-methyltransferase 3 beta
1591264	0.3849595	transaldolase 1
1434948	0.3849593	HIV TAT specific factor 1
489657	0.3847527	tryptophan rich basic protein

- 100 -

810947	0.3833137	LIS1-interacting protein NUDE1, rat homolog
298417	0.3833014	trefoil factor 3 (intestinal)
154707	0.3831976	MpV17 transgene, murine homolog, glomerulosclerosis
79032	0.3816428	CGI-82 protein
1916575	0.3814206	BCL2-interacting killer (apoptosis-inducing)
1585492	0.3812472	ESTs
1455394	0.3811391	cytochrome c
769942	0.3810371	kinesin-like 4
429387	0.3810004	chimerin (chimaerin) 2
768377	0.3809674	activity-dependent neuroprotective protein
214996	0.380548	ESTs
1557637	0.3805285	ESTs
1517595	0.3804132	KIAA0175 gene product
841260	0.3801713	hypothetical protein
491524	0.3785047	mitochondrial ribosomal protein L13
30170	0.3782337	caspase 3, apoptosis-related cysteine protease
76196	0.3781541	hypothetical protein FLJ20062
2017721	0.3781067	ESTs, Weakly similar to 2109260A B cell growth factor [H.sapiens]
134918	0.3779855	ESTs
347726	0.377767	homeo box D8
1469148	0.3776558	FGFR1 oncogene partner
1649374	0.3771383	homogentisate 1,2-dioxygenase (homogentisate oxidase)
241043	0.3770901	Human clone 137308 mRNA, partial cds
815835	0.3764754	hypothetical protein R33729_1
143426	0.3764075	ras homolog gene family, member B
1412245	0.3762847	carboxypeptidase A2 (pancreatic)
786265	0.3760397	KIAA0750 gene product
266500	0.3758559	ESTs
755301	0.3749416	protein kinase C, delta
294397	0.3744208	DKFZP586A0522 protein
155806	0.3741516	
41123	0.3739475	Homo sapiens, Similar to RIKEN cDNA 2210021G21 gene, clone MGC:14859 IMAGE:3621871, mRNA, complete cds
454339	0.3738929	thiopurine S-methyltransferase

- 101 -

810316	0.3732635	very long-chain acyl-CoA synthetase; lipidosis
812105	0.3731726	ALL1-fused gene from chromosome 1q
773426	0.3727398	KIAA0391 gene product
1572710	0.3725734	hypothetical protein FLJ21213
772898	0.372064	ribosomal protein S15a
283739	0.3714571	Homo sapiens cDNA FLJ14028 fis, clone HEMBA1003838
289505	0.3714361	ESTs, Moderately similar to S65657 alpha-1C-adrenergic receptor splice form 2 [H.sapiens]
897177	0.3712716	phosphoglycerate mutase 1 (brain)
782547	0.3708058	ESTs
647866	0.3706822	Homo sapiens cDNA FLJ13975 fis, clone Y79AA1001585
1679942	0.3706798	KIAA1053 protein
566443	0.3705134	Homo sapiens cDNA FLJ12793 fis, clone NT2RP2002033
302996	0.3698028	chloride intracellular channel 3
726699	0.3695061	ESTs, Weakly similar to AAB47496 NG5 [H.sapiens]
2094232	0.3693257	chromosome 1 open reading frame 12
2549634	0.3692338	activator of S phase kinase
177827	0.3691459	synaptotagmin VII
852829	0.3691079	karyopherin alpha 3 (importin alpha 4)
207794	0.3679245	nuclear factor (erythroid-derived 2), 45kD
823909	0.3675927	
743810	0.3667087	hypothetical protein MGC2577
810039	0.3662086	defender against cell death 1
53039	0.3659853	carbohydrate (keratan sulfate Gal-6) sulfotransferase 1
809727	0.365892	unc-51 (C. elegans)-like kinase 1
26171	0.3656147	KIAA0856 protein
1570663	0.3656056	FK506-binding protein 4 (59kD)
179212	0.3649976	ESTs, Moderately similar to T12539 hypothetical protein DKFZp434J154.1 [H.sapiens]
292996	0.36439	tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein, eta polypeptide
259950	0.364359	hypothetical protein FLJ14991
815794	0.3643266	nucleobindin 2
965223	0.3642603	thymidine kinase 1, soluble

- 102 -

592801	0.3638731	serine palmitoyltransferase, long chain base subunit 2
42831	0.3638165	N-terminal kinase-like
788641	0.3638059	adaptor-related protein complex 1, sigma 2 subunit
813845	0.3637519	RNA, U transporter 1
52076	0.3633002	neuroblastoma (nerve tissue) protein
154493	0.3629466	interferon-induced protein 75, 52kD
1519013	0.3628651	Homo sapiens, clone IMAGE:3537447, mRNA, partial cds
1434905	0.3624537	homeo box B7
825282	0.3624112	DKFZP586L0724 protein
322617	0.3623149	v-ral simian leukemia viral oncogene homolog B (ras related; GTP binding protein)
814353	0.3622592	phorbol-12-myristate-13-acetate-induced protein 1
2017917	0.360911	Homo sapiens cDNA: FLJ23371 fis, clone HEP16068, highly similar to HSTFIISH Homo sapiens mRNA for transcription elongation factor TFIIS
813616	0.3608471	FK506-binding protein like
1846982	0.3602851	inhibin, beta C
505289	0.3595549	angiotensin II, type I receptor-associated protein
1565079	0.358728	B-cell linker
795401	0.3573481	diacylglycerol O-acyltransferase (mouse) homolog
781047	0.357207	budding uninhibited by benzimidazoles 1 (yeast homolog)
346696	0.3570344	TEA domain family member 4
809466	0.3568464	DNA segment on chromosome 19 (unique) 1177 expressed sequence
743220	0.3566593	hypothetical protein FLJ12517
131268	0.3564734	growth factor receptor-bound protein 14
119133	0.3564449	Homo sapiens mRNA; cDNA DKFZp434B231 (from clone DKFZp434B231)
242706	0.3562469	Homo sapiens, Similar to RIKEN cDNA 5730494N06 gene, clone MGC:13349 IMAGE:4249231, mRNA, complete cds
66599	0.3561699	N-acetyltransferase 1 (arylamine N-acetyltransferase)
46716	0.3557556	ESTs

sd-71385

Patent 4,000,000

- 103 -

271899	0.355689	Ser-Thr protein kinase related to the myotonic dystrophy protein kinase
25440	0.3554359	staufen (Drosophila, RNA-binding protein) homolog 2
1899312	0.3551595	ESTs
783681	0.354881	upstream regulatory element binding protein 1
131091	0.3548154	Not56 (D. melanogaster)-like protein
300099	0.3547857	ESTs
588436	0.3544892	butyrate-induced transcript 1
2069602	0.3539588	melanocortin 1 receptor (alpha melanocyte stimulating hormone receptor)
810959	0.3534647	Rho GDP dissociation inhibitor (GDI) alpha
649977	0.3534567	Homo sapiens clone CDABP0014 mRNA sequence
345069	0.3532976	nuclear factor (erythroid-derived 2)-like 3
767487	0.3530592	ariadne (Drosophila) homolog, ubiquitin-conjugating enzyme E2-binding protein, 1
824943	0.3527414	hypothetical protein
744994	0.3526118	hypothetical protein FLJ12242
773381	0.3525581	N-ethylmaleimide-sensitive factor attachment protein, alpha
1568561	0.3518952	BCL2-like 1
2028916	0.3516982	Homo sapiens mRNA for Hmob33 protein, 3' untranslated region
753457	0.3515361	NADH dehydrogenase (ubiquinone) Fe-S protein 1 (75kD) (NADH-coenzyme Q reductase)
2161427	0.3509243	progesterone receptor
1947381	0.3509235	hypothetical protein FLJ22329
82421	0.3506778	zinc finger protein 6 (CMPX1)
504461	0.3506753	opsin 3 (encephalopsin)
754625	0.3503938	ATPase, Class II, type 9A
795543	0.3500184	thioredoxin peroxidase (antioxidant enzyme)
1910316	0.3498986	KIAA0535 gene product
172785	0.3493683	NAG-5 protein
1910078	0.3491481	ESTs, Weakly similar to YK54_YEAST HYPOTHETICAL 18.4 KD PROTEIN IN SIS2-MTD1 INTERGENIC REGION [S.cerevisiae]
321354	0.3489273	mitochondrial ribosomal protein L15

sd-71385

FOOTNOTES

- 104 -

878815	0.3478601	ADP-ribosylation factor 3
741769	0.3475255	polymerase (DNA directed), beta
487831	0.3464379	Homo sapiens cDNA FLJ14059 fis, clone HEMBB1000573
754046	0.3464097	DNA segment on chromosome X (unique) 9879 expressed sequence
344988	0.3464047	
530093	0.3463461	myelin protein zero-like 1
823615	0.3462131	Homo sapiens cDNA: FLJ21245 fis, clone COL01184
144881	0.3461767	calumenin
128695	0.3455381	ESTs, Weakly similar to I38344 titin, cardiac muscle [H.sapiens]
884511	0.3440809	cytochrome c oxidase subunit VIIb
128711	0.3437491	anillin (Drosophila Scraps homolog), actin binding protein
470035	0.3436575	prokineticin 1 precursor
1568126	0.3434836	solute carrier family 37 (glycerol-3-phosphate transporter), member 1
588911	0.3427563	2',5'-oligoadenylate synthetase 1 (40-46 kD)
35191	0.3427094	stromal cell-derived factor 2
814989	0.3426862	protein phosphatase 1G (formerly 2C), magnesium-dependent, gamma isoform
73638	0.3425644	protein tyrosine phosphatase type IVA, member 2
1605178	0.3416196	amino acid transporter system A1
214068	0.3414185	GATA-binding protein 3
755599	0.3413351	interferon induced transmembrane protein 1 (9-27)
782439	0.3413092	ATP synthase, H <sup>+</sup> transporting, mitochondrial F0 complex, subunit e
595070	0.3408109	stress-associated endoplasmic reticulum protein 1; ribosome associated membrane protein 4
183440	0.3406314	arylsulfatase A
774502	0.3402891	protein tyrosine phosphatase, non-receptor type 12
235986	0.338857	wingless-type MMTV integration site family, member 11
289936	0.3382708	solute carrier family 7, (cationic amino acid transporter, y <sup>+</sup> system) member 11
824108	0.3381616	SCAN domain-containing 1

sd-71385

F0222T 8T08200T



- 105 -

26617	0.3371923	activated leucocyte cell adhesion molecule
810497	0.3367408	ESTs, Weakly similar to A35363 synapsin I splice form a [H.sapiens]
40042	0.3362557	hypothetical protein FLJ10747
811142	0.3361932	phosphoinositide-3-kinase, regulatory subunit, polypeptide 2 (p85 beta)
2046679	0.3358043	ESTs
487444	0.3357395	cyclic AMP phosphoprotein, 19 kD
78736	0.3354693	Homo sapiens clone 24877 mRNA sequence
1637302	0.3353374	DNAJ domain-containing
825606	0.335057	kinesin-like 1
76362	0.3349202	spectrin, alpha, non-erythrocytic 1 (alpha-fodrin)
1883028	0.3345488	Homo sapiens mRNA; cDNA DKFZp434J1912 (from clone DKFZp434J1912)
789383	0.3341667	cAMP responsive element modulator
302549	0.3340582	basic transcription element binding protein 1
309288	0.334016	replication factor C (activator 1) 4 (37kD)
712314	0.3338146	myosin regulatory light chain interacting protein
241677	0.3332521	Homo sapiens, clone MGC:18110 IMAGE:4152745, mRNA, complete cds
898253	0.3331924	reticulocalbin 2, EF-hand calcium binding domain
770992	0.3331315	
81203	0.3328573	paraoxonase 3
416436	0.3325571	mitochondrial ribosomal protein L24
1660666	0.332029	carbonic anhydrase VB, mitochondrial
811582	0.3319191	golgi phosphoprotein 2
2054122	0.3318413	solute carrier family 11 (proton-coupled divalent metal ion transporters), member 3
837864	0.3314764	progesterone induced protein
1632247	0.3312131	hypothetical protein FLJ23436
298143	0.3307258	ESTs
2027952	0.3307056	hypothetical protein FLJ20297
1600281	0.3304323	nucleolar protein 3 (apoptosis repressor with CARD domain)
814270	0.3300091	polymyositis/scleroderma autoantigen 1 (75kD)

[illegible]

- 107 -

1631194	0.3219004	t-complex-associated-testis-expressed 1-like 1
366834	0.3217743	envoplakin
839580	0.3216628	ESTs
245485	0.3215041	hypothetical protein MGC5178
815501	0.3214849	hypothetical protein MGC2721
2017403	0.3211799	regulator of G-protein signalling 3
281476	0.3210873	aspartylglucosaminidase
130835	0.3209345	Homo sapiens, Similar to clone FLB3816, clone IMAGE:3454380, mRNA
77533	0.3205713	inositol polyphosphate-5-phosphatase, 40kD
42076	0.3194462	TRK-fused gene
810998	0.3179036	DKFZP434C131 protein
75859	-1.8328232	N-myc downstream-regulated gene 2
760299	-1.7010462	dickkopf (Xenopus laevis) homolog 3
344720	-1.5196431	glycophorin C (Gerbich blood group)
345670	-1.4445337	ESTs, Moderately similar to I59348 CCAAT binding transcription factor CBF subunit C - rat [R.norvegicus]
200814	-1.4325283	membrane metallo-endopeptidase (neutral endopeptidase, enkephalinase, CALLA, CD10)
1882697	-1.4318896	peanut (Drosophila)-like 2
1161564	-1.3891189	desmuslin
788234	-1.3761312	inhibitor of DNA binding 4, dominant negative helix-loop-helix protein
1558675	-1.3749762	SRY (sex determining region Y)-box 10
285377	-1.3699153	pellino (Drosophila) homolog 2
45099	-1.3515907	regucalcin (senescence marker protein-30)
611532	-1.3158379	troponin I, skeletal, fast
377275	-1.3102234	ataxia-telangiectasia group D-associated protein
810002	-1.3038896	Homo sapiens, clone MGC:19762 IMAGE:3636045, mRNA, complete cds
1569187	-1.2971252	heparan sulfate (glucosamine) 3-O-sulfotransferase 4
377461	-1.2798035	caveolin 1, caveolae protein, 22kD
712139	-1.2738321	ADP-ribosylation factor-like 7
742685	-1.272916	disabled (Drosophila) homolog 2 (mitogen-responsive phosphoprotein)

- 108 -

811920 -1.2690713 interleukin 11 receptor, alpha  
796542 -1.2625621 ets variant gene 5 (ets-related molecule)  
486683 -1.2614406 Homo sapiens mRNA; cDNA DKFZp564J0323 (from clone  
DKFZp564J0323)  
  
753071 -1.2529315 Homo sapiens cDNA: FLJ22528 fis, clone HRC12825  
  
131839 -1.2372478 folate receptor 1 (adult)  
300632 -1.2368481 hypothetical protein FLJ21044 similar to Rbig1  
  
839736 -1.2271618 crystallin, alpha B  
813265 -1.2230435 Homo sapiens mRNA; cDNA DKFZp564H1916 (from clone  
DKFZp564H1916)  
  
1469377 -1.2127781 lipoma HMGIC fusion partner-like 2  
490023 -1.2114805 hypothetical protein MGC2648  
298122 -1.1962478 frizzled (Drosophila) homolog 7  
72778 -1.1752838 caspase 7, apoptosis-related cysteine protease  
  
160192 -1.1751869 ESTs, Weakly similar to 2004399A chromosomal protein  
[H.sapiens]  
22917 -1.1590598 Homo sapiens mRNA; cDNA DKFZp761M0111 (from clone  
DKFZp761M0111)  
  
877621 -1.1501074 nGAP-like protein  
840266 -1.1476896 Homo sapiens cDNA: FLJ22667 fis, clone HSI08385  
  
289760 -1.1390686 ESTs, Highly similar to T00391 hypothetical protein KIAA0612  
[H.sapiens]  
811848 -1.1350073 hypothetical protein  
1556433 -1.1222978 GRO3 oncogene  
838478 -1.1175047 neurocalcin delta  
1473274 -1.1172693 myosin regulatory light chain 2, smooth muscle isoform  
  
1917449 -1.0894686 serum amyloid A4, constitutive  
811088 -1.0851546 ephrin-B3  
1554167 -1.0818834 hypothetical protein FLJ14529  
67741 -1.0705249 PP2135 protein  
2106144 -1.067765 regulated in glioma  
2056139 -1.0673271 LIM domain protein  
767202 -1.0654145 latent transforming growth factor beta binding protein 2

- 109 -

345034 -1.0600859 small inducible cytokine subfamily B (Cys-X-Cys), member 14  
(BRAK)  
666879 -1.0463245 annexin A8  
130201 -1.0401997 intercellular adhesion molecule 2  
878836 -1.037352 secretory granule, neuroendocrine protein 1 (7B2 protein)  
  
52419 -1.0368509 Friedreich ataxia region gene X123  
291478 -1.0212528 runt-related transcription factor 3  
781014 -1.0161379 suppression of tumorigenicity 5  
132857 -1.0084069 Homo sapiens mRNA; cDNA DKFZp586N1323 (from clone  
DKFZp586N1323)  
  
529843 -1.0036824 ESTs, Moderately similar to JC5238 galactosylceramide-like  
protein, GCP [H.sapiens]  
  
2504881 -1.000863 signal transducer and activator of transcription 5A  
  
69002 -0.9960916 PPAR(gamma) angiopoietin related protein  
  
841308 -0.9953716 myosin, light polypeptide kinase  
488404 -0.9880176 Homo sapiens clone TUA8 Cri-du-chat region mRNA  
  
277571 -0.9841522 KIAA1706 protein  
1871116 -0.9821709 Homo sapiens mRNA; cDNA DKFZp434C1714 (from clone  
DKFZp434C1714); partial cds  
  
897963 -0.9775616 phosphatidic acid phosphatase type 2A  
1584540 -0.9754693 Homo sapiens mRNA; cDNA DKFZp586M2022 (from clone  
DKFZp586M2022)  
  
196435 -0.9665947 ESTs  
140574 -0.9652218 small inducible cytokine subfamily D (Cys-X3-Cys), member 1  
(fractalkine, neurotactin)  
  
740620 -0.96496 tropomyosin 2 (beta)  
1473471 -0.9623482 KIAA0194 protein  
1635320 -0.9586226 amiloride-sensitive cation channel 2, neuronal  
  
416959 -0.957248 nuclear factor I/B  
814798 -0.9559895 aldehyde dehydrogenase 1 family, member A3  
  
823871 -0.9429443 SPARC-like 1 (mast9, hevin)  
303109 -0.9414602 purinergic receptor (family A group 5)

- 110 -

270826 -0.9389651 Homo sapiens cDNA FLJ13329 fis, clone OVARC1001795

360254 -0.9286811 cysteine-rich, angiogenic inducer, 61  
66491 -0.9280245 plasmolipin  
814316 -0.9276916 ribosomal protein L13  
153760 -0.9227865 EphB1  
811149 -0.9216006 chromosome 9 open reading frame 3  
786069 -0.9203158 beta-site APP-cleaving enzyme  
208718 -0.9190244 annexin A1  
416676 -0.9140494 pellino (Drosophila) homolog 1  
248631 -0.9106505 aminomethyltransferase (glycine cleavage system protein T)

76182 -0.9106484 hypothetical protein DKFZp761F241  
756708 -0.910505 potassium intermediate/small conductance calcium-activated  
channel, subfamily N, member 4

712401 -0.9068788 phosphoinositide-3-kinase, catalytic, delta polypeptide

625399 -0.9066615 hypothetical protein similar to beta-transducin family

753038 -0.893161 kinesin family member C3  
162308 -0.8910356 ESTs  
796181 -0.8888496 growth arrest-specific 6  
133236 -0.8888316 RNA binding motif protein, X chromosome  
415816 -0.8887425 ESTs  
323780 -0.8879854 Homo sapiens cDNA FLJ11177 fis, clone PLACE1007402

1493218 -0.884795 hypothetical protein FLJ22297  
1636166 -0.8845211 KIAA0668 protein  
1856063 -0.8833091 tweety (Drosophila) homolog 1  
34093 -0.8800259 EST  
2028722 -0.8800075 osteoblast specific factor 2 (fasciclin I-like)  
811837 -0.8764231 eukaryotic translation elongation factor 1 alpha 1

628955 -0.8761473 forkhead box O1A (rhabdomyosarcoma)  
884462 -0.8743987 Down syndrome critical region gene 1  
306798 -0.8718483 NGFI-A binding protein 1 (EGR1 binding protein 1)

882248 -0.8700731 transgelin  
770935 -0.8700104 hypothetical protein FLJ13511  
41208 -0.865227 bone morphogenetic protein 1  
768571 -0.8572018 SRY (sex determining region Y)-box 8

- 111 -

781017 -0.8526926 early growth response 2 (Krox-20 (Drosophila) homolog)  
811837 -0.848137  
815737 -0.8444322 ATP synthase, H<sup>+</sup> transporting, mitochondrial F1 complex,  
alpha subunit, isoform 1, cardiac muscle  
814826 -0.8443265 ESTs  
767164 -0.842198 ESTs, Weakly similar to MUC2\_HUMAN MUCIN 2  
PRECURSOR [H.sapiens]  
71087 -0.8416951 v-maf musculoaponeurotic fibrosarcoma (avian) oncogene  
family, protein F  
283023 -0.8413949 chemokine (C-X3-C) receptor 1  
160609 -0.8405669 ESTs  
814443 -0.8385965 hypothetical protein MGC3232  
142259 -0.8376566 tumor necrosis factor alpha-inducible cellular protein containing  
leucine zipper domains; Huntingtin interacting protein L;  
transcription factor IIIA-interacting protein  
990881 -0.8335353  
970590 -0.8333968 Homo sapiens mRNA; cDNA DKFZp434A115 (from clone  
DKFZp434A115)  
470393 -0.830751 matrix metalloproteinase 7 (matrilysin, uterine)  
772913 -0.8306579 calreticulin  
741139 -0.8299902 eyes absent (Drosophila) homolog 2  
27769 -0.8298821 ESTs  
544639 -0.8275232 ESTs  
1587710 -0.8252065 period (Drosophila) homolog 1  
757191 -0.8247438 ESTs  
80344 -0.8243418 interleukin 7 receptor  
1635062 -0.8243024 DKFZP586A011 protein  
119290 -0.8224756 cortic al thymocyte receptor (X. laevis CTX) like  
161456 -0.8215502 serum amyloid A1  
1909935 -0.8169271 ESTs  
1257131 -0.8163443 ESTs  
868396 -0.808677 tyrosine 3-monooxygenase/tryptophan 5-monooxygenase  
activation protein, eta polypeptide  
1609665 -0.8086161 BarH-like homeobox 2  
2016775 -0.8063264 G protein-coupled receptor, family C, group 5, member B

796475	-0.8041136	four and a half LIM domains 3
1555924	-0.8025163	CSR1 protein
416434	-0.8006527	SCN Circadian Oscillatory Protein (SCOP)
322561	-0.7970805	ribosomal protein L31
132711	-0.7962294	Kruppel-like factor 5 (intestinal)
70245	-0.795375	Homo sapiens mRNA full length insert cDNA clone EUROIMAGE 50374
283124	-0.7933772	ESTs, Moderately similar to LONG-CHAIN FATTY ACID TRANSPORT PROTEIN [M.musculus]
281190	-0.7927001	ESTs
250883	-0.7925068	ubiquitin-activating enzyme E1-like
346545	-0.7887371	laminin, beta 1
1878409	-0.7869766	catechol-O-methyltransferase
530958	-0.7809384	smoothened (Drosophila) homolog
40027	-0.7779473	ESTs
788136	-0.7778375	phosphodiesterase 4B, cAMP-specific (dunce (Drosophila)- homolog phosphodiesterase E4)
1626304	-0.7769866	spinal cord-derived growth factor-B
1474337	-0.7766266	phosphorylase, glycogen; brain
85840	-0.7754435	nicotinamide N-methyltransferase
290378	-0.7751825	podocalyxin-like
2243051	-0.7736287	kallikrein 8 (neurotrophin/ovastin)
813256	-0.7731864	ATP-binding cassette, sub-family B (MDR/TAP), member 1
47043	-0.7729643	tensin
343695	-0.7706809	hypothetical protein FLJ10875
491403	-0.7700599	tumor necrosis factor receptor superfamily, member 1B
767765	-0.7691523	GTP-binding protein overexpressed in skeletal muscle
681992	-0.7691299	Homo sapiens cDNA FLJ13384 fis, clone PLACE1001062, highly similar to Homo sapiens mRNA for lysine-ketoglutarate reductase/saccharopine dehydrogenase
505864	-0.7677235	RaIGDS-like gene
139660	-0.7677153	ESTs
869450	-0.7659922	ribosomal protein L11
162308	-0.7654139	
209537	-0.763749	zinc finger protein 221
753162	-0.7617972	KIAA0603 gene product



- 113 -

810728 -0.7607171 hypothetical gene ZD52F10  
366591 -0.7595643 T-cell lymphoma invasion and metastasis 2  
  
488258 -0.7574184 collagen, type XVI, alpha 1  
32489 -0.7571259 hypothetical protein DKFZp566A1524  
1519147 -0.7548183  
418159 -0.7545888 synaptogyrin 1  
80727 -0.7538606 receptor tyrosine kinase-like orphan receptor 1  
  
593023 -0.7516846 dystrobrevin, beta  
586803 -0.7423407 placental growth factor, vascular endothelial growth factor-  
related protein  
188335 -0.7399294 egf-like module containing, mucin-like, hormone receptor-like  
sequence 2  
1161775 -0.7380353 villin 1  
490668 -0.7364434 Homo sapiens mRNA; cDNA DKFZp586B211 (from clone  
DKFZp586B211)  
  
1474900 -0.736252 keratin 15  
378813 -0.7346408 secretory leukocyte protease inhibitor (antileukoproteinase)  
  
1630990 -0.7312981 ribosomal protein L29  
280907 -0.7307519 Kruppel-type zinc finger protein  
742562 -0.730047 chromosome 16 open reading frame 5  
73252 -0.7283898 3-hydroxy-3-methylglutaryl-Coenzyme A synthase 1 (soluble)  
  
26566 -0.7278186 protein-O-mannosyltransferase 1  
1471829 -0.7275378 uncharacterized hypothalamus protein HSMNP1  
  
289428 -0.7270494 neurotrophic tyrosine kinase, receptor, type 2  
  
898092 -0.7267434 connective tissue growth factor  
34150 -0.7263513 ESTs  
24958 -0.7225195 Homo sapiens mRNA; cDNA DKFZp434C2016 (from clone  
DKFZp434C2016)  
  
725680 -0.7197488 transcription factor AP-2 gamma (activating enhancer-binding  
protein 2 gamma)  
  
77972 -0.7194713 complement component 3  
1686766 -0.7192745 Rag D protein  
471196 -0.7192268 integral membrane protein 3

- 114 -

431231 -0.7182897 EGF-containing fibulin-like extracellular matrix protein 2  
1908834 -0.7177051 ubiquitin A-52 residue ribosomal protein fusion product 1  
207735 -0.7167416 serine (or cysteine) proteinase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1  
754157 -0.7165332 Homo sapiens mRNA; cDNA DKFZp434K2172 (from clone DKFZp434K2172)  
35828 -0.7163689 diphtheria toxin receptor (heparin-binding epidermal growth factor-like growth factor)  
1946448 -0.7150027 caveolin 2  
470148 -0.712917 ESTs  
840944 -0.7099353 early growth response 1  
809784 -0.7095414 kallikrein 6 (neurosin, zyme)  
877827 -0.7093855 ribosomal protein S27a  
32493 -0.7093777 integrin, alpha 6  
813520 -0.7092422 EphB3  
782501 -0.7086066 hypothetical protein PP1665  
27544 -0.708383 prominin (mouse)-like 1  
184022 -0.7070266 amyloid beta (A4) precursor protein-binding, family B, member 1 (Fe65)  
323238 -0.7042807 GRO1 oncogene (melanoma growth stimulating activity, alpha)  
488956 -0.70192 CUG triplet repeat, RNA-binding protein 2  
1926246 -0.701769 ESTs, Moderately similar to T46371 hypothetical protein DKFZp434P1018.1 [H.sapiens]  
325365 -0.7012106 HIV-1 rev binding protein 2  
221846 -0.6982802 checkpoint suppressor 1  
1870305 -0.6958213 heat shock 27kD protein 2  
277044 -0.6947855 KIAA1183 protein  
503335 -0.6944661 hypothetical protein FLJ11196  
757873 -0.6940266 cyclin-dependent kinase 5, regulatory subunit 1 (p35)  
2306682 -0.6934667 A kinase (PRKA) anchor protein (yotiao) 9  
1864302 -0.6932327 E74-like factor 5 (ets domain transcription factor)  
843070 -0.6925728 nucleoporin 88kD  
812975 -0.6921755 KIAA0172 protein

- 115 -

46129	-0.6919169	ESTs, Highly similar to T17245 hypothetical protein DKFZp586J0917.1 [H.sapiens]
270917	-0.6918383	secreted frizzled-related protein 1
789382	-0.6898974	Notch (Drosophila) homolog 4
611481	-0.6894079	HMG-box transcription factor TCF-3
35300	-0.688788	KIAA0869 protein
50892	-0.6887497	Homo sapiens, clone MGC:9913 IMAGE:3870821, mRNA, complete cds
366100	-0.6885372	matrilin 2
345056	-0.6863529	KIAA1404 protein
2783721	-0.6854649	cholinergic receptor, nicotinic, beta polypeptide 1 (muscle)
178137	-0.6839022	ribosomal protein L34
1626996	-0.682449	c-fos induced growth factor (vascular endothelial growth factor D)
188388	-0.6824191	integrin, alpha 10
133518	-0.6817662	microtubule-associated protein, RP/EB family, member 2
839796	-0.6807912	candidate tumor suppressor p33 ING1 homolog
726582	-0.6807781	actin-related protein 3-beta
50586	-0.6794282	KIAA1545 protein
491763	-0.679034	interleukin 1, beta
52096	-0.6789228	platelet-derived growth factor receptor, alpha polypeptide
263716	-0.6767498	collagen, type VI, alpha 1
2095066	-0.676021	alcohol dehydrogenase 1C (class I), gamma polypeptide
1475738	-0.6753091	ribosomal protein S25
41869	-0.6747514	hypothetical protein FLJ11017
1350439	-0.6740833	KIAA0015 gene product
502689	-0.6739476	Homo sapiens clone HH409 unknown mRNA
773373	-0.6730112	hypothetical protein MGC14258
308539	-0.6726187	Homo sapiens cDNA FLJ12777 fis, clone NT2RP2001720
343760	-0.6724144	SH3 domain binding glutamic acid-rich protein like 2
769959	-0.6720196	collagen, type IV, alpha 2
2252954	-0.6687686	enhancer of filamentation 1 (cas-like docking; Crk-associated substrate related)
70201	-0.6675012	mitochondrial solute carrier

sd-71385

485772004300

505243	-0.6671218	inositol 1,4,5-triphosphate receptor, type 2
897731	-0.6670507	latrophilin
296880	-0.6655878	membrane protein, palmitoylated 1 (55kD)
1486082	-0.6635259	heparin-binding growth factor binding protein
1470278	-0.6633411	hypothetical protein FLJ21841
177665	-0.6624986	
725152	-0.6591675	hypothetical protein DKFZp762A227
247616	-0.6585351	lipoma HMGIC fusion partner
752668	-0.65723	KIAA0440 protein
837870	-0.657186	proline arginine-rich end leucine-rich repeat protein
744918	-0.6552361	Homo sapiens mRNA; cDNA DKFZp761I0911 (from clone DKFZp761I0911)
39600	-0.6540275	adenylate kinase 5
795288	-0.6528792	ubiquitin specific protease 4 (proto-oncogene)
1665444	-0.651994	tumor endothelial marker 1 precursor
774078	-0.6508021	leiomodulin 1 (smooth muscle)
415233	-0.6506593	ribosomal protein L37a
307029	-0.6500872	ribosomal protein L26
78946	-0.648366	Homo sapiens pyruvate dehydrogenase kinase 4 mRNA, 3' untranslated region, partial sequence
785733	-0.6476493	hypothetical protein FLJ12892
415415	-0.646492	EST, Moderately similar to RL1X_HUMAN 60S RIBOSOMAL PROTEIN L18A [H.sapiens]
858167	-0.6453859	fatty-acid-Coenzyme A ligase, long-chain 4
823714	-0.642903	nuclear receptor co-repressor/HDAC3 complex subunit
193913	-0.6425581	v-yes-1 Yamaguchi sarcoma viral related oncogene homolog
220395	-0.6418175	hypothetical protein FLJ23293 similar to ARL-6 interacting protein-2
840942	-0.6383764	major histocompatibility complex, class II, DP beta 1
725308	-0.6371261	ras inhibitor
1492147	-0.636656	ribosomal protein S4, X-linked
1466893	-0.6364598	ESTs
75078	-0.6362307	ESTs

- 117 -

796152 -0.6356106 Homo sapiens cDNA FLJ11685 fis, clone HEMBA1004934  
190059 -0.634814 guanine nucleotide binding protein (G protein), gamma 7  
840511 -0.6341661 vimentin  
81409 -0.6338327 GABA(A) receptor-associated protein like 1  
745490 -0.6329354 hypothetical protein FLJ20607  
530036 -0.6324794 hypothetical protein FLJ13078  
755855 -0.6298333 solute carrier family 25 (mitochondrial carrier; adenine  
nucleotide translocator), member 6  
785538 -0.6289951 Homo sapiens mRNA; cDNA DKFZp434N2116 (from clone  
DKFZp434N2116)  
2284803 -0.6265431 secretory carrier membrane protein 1  
1623016 -0.6263561 EST  
154472 -0.6253276 fibroblast growth factor receptor 1 (fms-related tyrosine kinase  
2, Pfeiffer syndrome)  
774754 -0.6243946 catenin (cadherin-associated protein), beta 1 (88kD)  
1572298 -0.6228686 CD3Z antigen, zeta polypeptide (TiT3 complex)  
2252417 -0.6222381 ribosomal protein S10  
180803 -0.621787 inositol polyphosphate-1-phosphatase  
683059 -0.6216483 Cdc42 effector protein 3  
1577736 -0.6215965 epidermal growth factor (beta-urogastrone)  
241489 -0.6212938 adrenergic, beta-2-, receptor, surface  
2316441 -0.6206597 ras homolog gene family, member C  
1712825 -0.6204409 ESTs  
52724 -0.6199204 hypothetical protein FLJ20241  
150897 -0.6188264 UDP-GlcNAc:betaGal beta-1,3-N-  
acetylglucosaminyltransferase 3  
823851 -0.6182341 AE-binding protein 1  
810057 -0.6178096 vasoactive intestinal peptide receptor 1  
81316 -0.6176265 ESTs  
298134 -0.6152731 frizzled (Drosophila) homolog 1  
2028238 -0.6141182 KIAA0300 protein  
1898758 -0.6126872 BCE-1 protein  
250797 -0.6124667 hypothetical protein FLJ20038  
754040 -0.610807 NS1-associated protein 1  
1584628 -0.6102937 alpha-actinin-2-associated LIM protein

sd-71385

FILED IN 12-18-2004

230910	-0.6096107	
172783	-0.6088873	hypothetical protein FLJ10390
824799	-0.6054867	ART-4 protein
785967	-0.6049171	erythrocyte membrane protein band 4.1-like 2
342349	-0.603487	mitogen-activated protein kinase kinase kinase 14
67067	-0.6003961	Homo sapiens mRNA; cDNA DKFZp564O222 (from clone DKFZp564O222)
1533710	-0.5988282	ortholog of mouse integral membrane glycoprotein LIG-1
768007	-0.5983563	hypothetical protein
1707637	-0.5964339	ESTs
1634832	-0.5931096	class I cytokine receptor
1609625	-0.5920312	selectin P ligand
1592530	-0.5918818	mammalian inositol hexakisphosphate kinase 2
272706	-0.5914377	alpha2,3-sialyltransferase
897720	-0.5904719	trophinin
1472735	-0.5888169	metallothionein 1E (functional)
767641	-0.5881449	mitogen-activated protein kinase 8 interacting protein 2
1651906	-0.5863631	hypothetical protein FLJ23544
564801	-0.5824549	Homo sapiens cDNA: FLJ21409 fis, clone COL03924
248886	-0.5823315	rab3 GTPase-activating protein, non-catalytic subunit (150kD)
250654	-0.5815576	secreted protein, acidic, cysteine-rich (osteonectin)
1968422	-0.5802058	Homo sapiens mRNA full length insert cDNA clone EUROIMAGE 1968422
712829	-0.5797871	LIM domain only 2 (rhombotin-like 1)
752837	-0.5794612	Homo sapiens mRNA for FLJ00074 protein, partial cds
346902	-0.5794362	ESTs, Weakly similar to ALU2_HUMAN ALU SUBFAMILY SB SEQUENCE CONTAMINATION WARNING ENTRY [H.sapiens]
782193	-0.5788147	thioredoxin
358531	-0.5784457	v-jun avian sarcoma virus 17 oncogene homolog
2271240	-0.5782432	hypothetical protein

- 119 -

66532 -0.5773248 endothelin 3  
120106 -0.5770131 caspase 1, apoptosis-related cysteine protease (interleukin 1,  
beta, convertase)  
810552 -0.5768989 B-cell associated protein  
770848 -0.5767154 ESTs, Weakly similar to ALU1\_HUMAN ALU SUBFAMILY J  
SEQUENCE CONTAMINATION WARNING ENTRY  
[H.sapiens]  
1032831 -0.5764161 glycosyltransferase  
460403 -0.5739604 laminin, gamma 2 (nicein (100kD), kalinin (105kD), BM600  
(100kD), Herlitz junctional epidermolysis bullosa))  
489755 -0.5730453 a disintegrin and metalloproteinase domain 12 (meltrin alpha)  
502753 -0.5724623 angiopoietin 2  
68534 -0.5724394 Homo sapiens cDNA: FLJ22290 fis, clone HRC04405  
854645 -0.5723062 CDC-like kinase 3  
322024 -0.5714933 ESTs, Highly similar to T12495 hypothetical protein  
DKFZp434H071.1 [H.sapiens]  
140071 -0.5708303 frizzled-related protein  
788518 -0.5697316 peroxisomal membrane protein 3 (35kD, Zellweger syndrome)  
240752 -0.56913 hypothetical protein MGC14797  
773487 -0.5686324 hypothetical protein MGC3032  
1404774 -0.5685596 parathyroid hormone-like hormone  
814815 -0.5675649 plakophilin 4  
462939 -0.5662456 ESTs  
240248 -0.5660209 transferrin  
208001 -0.5640857 CD59 antigen p18-20 (antigen identified by monoclonal  
antibodies 16.3A5, EJ16, EJ30, EL32 and G344)  
1909433 -0.5638242 DKFZP727C091 protein  
1636360 -0.5629882 hypothetical protein FLJ14957  
138242 -0.5621088 ESTs, Moderately similar to MAS2\_HUMAN MANNAN-  
BINDING LECTIN SERINE PROTEASE 2 PRECURSOR  
[H.sapiens]  
703964 -0.5620699 inositol polyphosphate phosphatase-like 1  
1522734 -0.5618289 ESTs  
843094 -0.5617478 ubiquitin-like 1 (sentrin)

- 120 -

491519 -0.561401 Homo sapiens clone 24775 mRNA sequence

377314 -0.561312 casein kinase 2, alpha prime polypeptide

208969 -0.561267 EST

788558 -0.5600825 KIAA1479 protein

785530 -0.5600048 integrin, alpha 1

1473131 -0.5594101 transducin-like enhancer of split 2, homolog of Drosophila  
E(sp1)

812959 -0.5591846 KIAA1638 protein

868169 -0.5584677 lipoprotein lipase

322233 -0.5578912 ribosomal protein, large, P0

50519 -0.5577761 aryl hydrocarbon receptor nuclear translocator-like

299720 -0.5565137

215000 -0.555725 vasoactive intestinal peptide receptor 1

51981 -0.5554908 ribosomal protein L7a

703541 -0.5534921 KIAA1858 protein

238907 -0.5534598 hypothetical protein, clone  
Telethon(Italy\_B41)\_Strait02270\_FL142

415613 -0.5528761 DHHC1 protein

282404 -0.5528532 Homo sapiens mRNA for KIAA1671 protein, partial cds

344588 -0.55146 kallikrein 5

746373 -0.5513143 RNA polymerase I transcription factor RRN3

491186 -0.5507413 Homo sapiens cDNA: FLJ23131 fis, clone LNG08502

75059 -0.5504962 poly(A)-binding protein, cytoplasmic 1-like

345764 -0.5504113 special AT-rich sequence binding protein 1 (binds to nuclear  
matrix/scaffold-associating DNA's)

1577920 -0.5496944 ESTs

288748 -0.5493871 Homo sapiens, clone MGC:5352 IMAGE:3048106, mRNA,  
complete cds

810097 -0.5487174 Homo sapiens cDNA: FLJ21721 fis, clone COLF0381

755881 -0.5473964 aquaporin 5

1568391 -0.5467858 plastin 3 (T isoform)

290866 -0.5467613 v-raf-1 murine leukemia viral oncogene homolog 1

503671 -0.5461871 Homo sapiens cDNA FLJ14368 fis, clone HEMBA1001122

2388571 -0.5459823 A kinase (PRKA) anchor protein 8

sd-71385

10028018-120101



739193	-0.5457998	cellular retinoic acid-binding protein 1
756372	-0.5447652	retinoic acid receptor responder (tazarotene induced) 2
53081	-0.5446617	ESTs
486179	-0.5443525	Homo sapiens cDNA FLJ10205 fis, clone HEMBA1004954
84464	-0.5426397	hypothetical protein FLJ12806
2213824	-0.5412089	protease inhibitor 3, skin-derived (SKALP)
1597813	-0.5406089	hypothetical protein PP1044
810358	-0.5403705	acyl-Coenzyme A dehydrogenase, very long chain
341759	-0.5403406	lung type-I cell membrane-associated glycoprotein
309449	-0.5401954	ribosomal protein S4, Y-linked
42681	-0.5388424	NY-REN-25 antigen
299720	-0.5388238	eukaryotic translation elongation factor 1 alpha 1
162775	-0.5385174	DEAD/H (Asp-Glu-Ala-Asp/His) box polypeptide 5 (RNA helicase, 68kD)
490102	-0.5383385	TBP-associated factor 172
877835	-0.5382125	ribosomal protein L35
299720	-0.5378801	eukaryotic translation elongation factor 1 alpha 1
1534493	-0.5376323	ESTs
595637	-0.5373697	Homo sapiens mRNA; cDNA DKFZp586N012 (from clone DKFZp586N012)
725622	-0.5370309	ESTs
233679	-0.5356625	hypothetical protein FLJ22362
785933	-0.5340762	sushi-repeat-containing protein, X chromosome
825356	-0.533771	Homo sapiens cDNA FLJ11997 fis, clone HEMBB1001458
767982	-0.533231	Homo sapiens, clone IMAGE:4134852, mRNA, partial cds
277627	-0.5329107	Human SH3 domain-containing protein SH3P18 mRNA, complete cds
460126	-0.5320146	KIAA0819 protein
346688	-0.5315035	melanoma inhibitory activity
712023	-0.5302372	AT-binding transcription factor 1
24729	-0.5302207	cholinergic receptor, muscarinic 1
1590021	-0.5302114	ets variant gene 6 (TEL oncogene)
809998	-0.5297152	

- 122 -

811779	-0.5291718	phosphoinositide-3-kinase, regulatory subunit, polypeptide 1 (p85 alpha)
488130	-0.5289721	Homo sapiens cDNA FLJ20767 fis, clone COL06986
299720	-0.5284407	
159462	-0.528299	serum constituent protein
1570427	-0.5280955	hypothetical protein MGC4309
2308346	-0.5268833	cyclin-dependent kinase 2
1881774	-0.5265891	KIAA1678
591055	-0.5258289	protein kinase C and casein kinase substrate in neurons 2
470128	-0.5253542	myosin IE
687990	-0.5253354	Rac/Cdc42 guanine exchange factor (GEF) 6
753620	-0.5248489	insulin-like growth factor binding protein 6
80374	-0.5236475	pyruvate dehydrogenase (lipoamide) alpha 1
878421	-0.5233873	transgelin 2
343079	-0.5224833	Homo sapiens mRNA; cDNA DKFZp761P0114 (from clone DKFZp761P0114)
284592	-0.5205573	PRO1659 protein
855061	-0.5203046	vascular endothelial growth factor B
183704	-0.5184644	Homo sapiens, clone MGC:13446 IMAGE:4275731, mRNA, complete cds
155896	-0.5182247	ORF
842939	-0.5172329	adenosine deaminase, RNA-specific, B1 (homolog of rat RED1)
685185	-0.5172158	reticulon 2
753301	-0.5171758	carcinoembryonic antigen-related cell adhesion molecule 1 (biliary glycoprotein)
795178	-0.5169102	lactate dehydrogenase C
1911663	-0.5168962	ESTs
345023	-0.5166203	ESTs, Highly similar to T08701 hypothetical protein DKFZp564N123.1 [H.sapiens]
1631682	-0.5162406	peptidylprolyl isomerase E (cyclophilin E)
868308	-0.5157133	ribosomal protein S23
809998	-0.5144681	amylase, alpha 2A; pancreatic
771004	-0.5144276	KIAA1201 protein
842896	-0.5137089	hypothetical protein DKFZp762L0311
714213	-0.5129593	tumor necrosis factor receptor superfamily, member 6

- 123 -

197056 -0.5128128 ESTs  
415191 -0.512457 KIAA0161 gene product  
2306752 -0.5123647 stathmin-like 2  
469306 -0.5113377 gastrin-releasing peptide  
824602 -0.5107682 interferon, gamma-inducible protein 16  
307687 -0.5106552 protease, serine, 16 (thymus)  
1637296 -0.5105214 ribosomal protein S24  
1358393 -0.5090372 mitogen-activated protein kinase kinase 3  
504940 -0.508056  
713839 -0.5080365 transcription factor AP-4 (activating enhancer-binding protein 4)  
  
220293 -0.5078291 Homo sapiens cDNA: FLJ21800 fis, clone HEP00618  
  
125342 -0.5078237 SUMO-1-specific protease  
1601845 -0.5074894 Ca<sup>2+</sup>-promoted Ras inactivator  
240620 -0.5065007 vascular Rab-GAP/TBC-containing  
43679 -0.5063146 ESTs  
825013 -0.505654 acidic protein rich in leucines  
2116188 -0.5046972 histone deacetylase 5  
161998 -0.5044691 hypothetical protein FLJ23138  
83999 -0.5044612 Human DNA sequence from clone 889N15 on chromosome  
Xq22.1-22.3. Contains part of the gene for a novel protein  
similar to X. laevis Cortical Thymocyte Marker CTX, the  
possibly alternatively spliced gene for 26S Proteasome subunit  
p28 (Ankyrin repeat prote  
  
593840 -0.5033966 DKFZP564K1964 protein  
51448 -0.5031992 activating transcription factor 3  
383501 -0.502319 regulator of G-protein signalling 9  
2018423 -0.501705 death-associated protein kinase 2  
796984 -0.5010588 cytochrome b-245, beta polypeptide (chronic granulomatous  
disease)  
1160723 -0.5005952 LIM domain kinase 2  
414999 -0.5002542 ets variant gene 4 (E1A enhancer-binding protein, E1AF)  
  
770212 -0.4981295 chitinase 3-like 1 (cartilage glycoprotein-39)  
  
48518 -0.4966991 ATP-binding cassette, sub-family A (ABC1), member 5  
  
85634 -0.4961026 complement component 1, s subcomponent

- 124 -

341763 -0.4953701 caspase 5, apoptosis-related cysteine protease

812161 -0.4948424 ESTs

1417886 -0.4941304 hypothetical protein FLJ23239

1500815 -0.4934665 Homo sapiens cDNA: FLJ22130 fis, clone HEP19632

868400 -0.4928479 glutaminyl-tRNA synthetase

809894 -0.491862 acetyl-CoA synthetase

753745 -0.4917192 ESTs, Weakly similar to S57447 HPBR11-7 protein [H.sapiens]

85394 -0.4913773 phosphatidic acid phosphatase type 2B

134192 -0.4908497 ESTs

813698 -0.4898644 sprouty (Drosophila) homolog 2

253884 -0.4897811 Human BAC clone GS1-99H8

131012 -0.4897744 hypothetical protein FLJ10633

855755 -0.4897236 fibrillarin

1711456 -0.4896574 H factor (complement)-like 1

815142 -0.4881454 spastic ataxia of Charlevoix-Saguenay (sacsin)

2413337 -0.4879268 sortilin-related receptor, L(DLR class) A repeats-containing

327432 -0.4870045 semaphorin Y

1897944 -0.4855156 ESTs, Weakly similar to T00366 hypothetical protein KIAA0669 [H.sapiens]

Example VI: Genes for discriminating between normal and ADH (non-malignant) versus DCIS and IDC (malignant)

As shown in Table 5 below, 400 genes were identified as being able to discriminate between normal and ADH (non-malignant) versus DCIS and IDC (malignant).

**Table 5**

CloneID	Weight	Description
796469	1.4207633	HSPC150 protein similar to ubiquitin-conjugating enzyme
488964	1.3447179	H2A histone family, member O

sd-71385

- 125 -

1505038	1.3277637	hypothetical protein FLJ20171
1500000	1.2926116	H2B histone family, member B
1554549	1.2787033	hydroxyacyl glutathione hydrolase
812238	1.2664748	hypothetical protein MGC4692
35147	1.2581066	ESTs, Weakly similar to unnamed protein product [H.sapiens]
122077	1.2576139	putative membrane protein
788654	1.2573483	growth factor receptor-bound protein 2
595037	1.2486446	retinoic acid induced 3
565319	1.2155833	Homo sapiens mRNA; cDNA DKFZp564B1264 (from clone DKFZp564B1264)
283919	1.2112507	H2A histone family, member L
1917941	1.1872008	purine-rich element binding protein B
359887	1.1837896	translocase of inner mitochondrial membrane 17 (yeast) homolog A
471568	1.1673113	hematological and neurological expressed 1
290841	1.1670252	H2B histone family, member A
796694	1.1580364	baculoviral IAP repeat-containing 5 (survivin)
366132	1.1440486	succinate dehydrogenase complex, subunit C, integral membrane protein, 15kD
823598	1.1425746	proteasome (prosome, macropain) 26S subunit, non-ATPase, 12
1323448	1.1309009	cysteine-rich protein 1 (intestinal)
810711	1.1074523	stearoyl-CoA desaturase (delta-9-desaturase)
741474	1.0870449	glucose phosphate isomerase
745606	1.0663136	hypothetical protein PP591
2054635	1.0613961	proteasome (prosome, macropain) subunit, alpha type, 7
178805	1.0513873	Human DNA sequence from clone RP5-850E9 on chromosome 20. Contains part of the gene for a novel C2H2 type zinc finger protein similar to Drosophila Scratch (Scrt), Slug and Xenopus Snail, a novel gene similar to Drosophila CG6762, STSs, GSSs and five CpG
347373	1.0454939	transcription elongation factor B (SIII), polypeptide 1 (15kD, elongin C)

- 126 -

199403	1.0142329	lectin, galactoside-binding, soluble, 8 (galectin 8)
814054	1.0130821	KIAA0040 gene product
2029173	1.0094499	ESTs, Weakly similar to N-WASP [H.sapiens]
209066	1.0066096	
782428	1.0008279	KIAA0250 gene product
769921	0.9944462	ubiquitin carrier protein E2-C
470061	0.9920108	seven in absentia (Drosophila) homolog 2
796723	0.9915885	Homo sapiens clone CDABP0014 mRNA sequence
1609836	0.9886168	glutamate-ammonia ligase (glutamine synthase)
2322367	0.9859632	reticulon 4
280375	0.9793036	PRO2000 protein
2016908	0.969649	ESTs, Weakly similar to CA13_HUMAN COLLAGEN ALPHA 1(III) CHAIN PRECURSOR [H.sapiens]
1858892	0.9669022	hypothetical protein MGC4825
46248	0.9628117	ADP-ribosyltransferase (NAD <sup>+</sup> ; poly (ADP-ribose) polymerase)
2014034	0.9619198	methylene tetrahydrofolate dehydrogenase (NAD <sup>+</sup> dependent), methenyltetrahydrofolate cyclohydrolase
898032	0.9547022	KIAA0097 gene product
725454	0.9476507	CDC28 protein kinase 2
79520	0.9457391	RAB2, member RAS oncogene family
810558	0.943041	proteasome (prosome, macropain) 26S subunit, ATPase, 4
272529	0.9423688	phosphomannomutase 2
122241	0.939204	proteasome (prosome, macropain) subunit, beta type, 2
469686	0.9381847	Ric (Drosophila)-like, expressed in many tissues
624667	0.9336899	CGI-92 protein
488202	0.932881	ESTs, Weakly similar to YZ28_HUMAN HYPOTHETICAL PROTEIN ZAP128 [H.sapiens]
825470	0.9274271	topoisomerase (DNA) II alpha (170kD)
1640821	0.920874	ESTs, Weakly similar to I78885 serine/threonine-specific protein kinase [H.sapiens]

- 127 -

686552	0.9182272	golgi phosphoprotein 1
2016648	0.917373	Homo sapiens mRNA; cDNA DKFZp434N1728 (from clone DKFZp434N1728)
1911343	0.9110591	RAB26, member RAS oncogene family
781097	0.9100493	reticulon 3
244801	0.908481	Rho guanine exchange factor (GEF) 11
754628	0.9061145	ESTs
1574058	0.9058211	1-acylglycerol-3-phosphate O-acyltransferase 2 (lysophosphatidic acid acyltransferase, beta)
753299	0.9019921	hypothetical protein FLJ10504
811774	0.9014801	CGI-49 protein
595213	0.8978322	hypothetical protein
868128	0.8970146	JM4 protein
810124	0.8891272	platelet-activating factor acetylhydrolase, isoform 1b, gamma subunit (29kD)
66406	0.8888639	hypothetical protein DKFZp762E1312
1636092	0.8841788	hypothetical protein FLJ20657
1869201	0.8825788	hypothetical protein MGC2745
625923	0.8821749	phosphoenolpyruvate carboxykinase 2 (mitochondrial)
1492238	0.8804306	HSPC003 protein
731044	0.8716644	glutaredoxin 2
839682	0.8696528	ubiquitin-conjugating enzyme E2N (homologous to yeast UBC13)
951233	0.8614727	proteasome (prosome, macropain) subunit, beta type, 3
843195	0.8606568	phosphoserine phosphatase
288999	0.8593924	small protein effector 1 of Cdc42
51773	0.8588635	hypothetical protein MGC3077
209066	0.8582298	serine/threonine kinase 15
1474955	0.8573467	TATA box binding protein (TBP)-associated factor, RNA polymerase II, N, 68kD (RNA-binding protein 56)
2043167	0.8551193	BCL2-associated athanogene 3
742707	0.8515067	ESTs, Weakly similar to MUC2_HUMAN MUCIN 2 PRECURSOR [H.sapiens]
743589	0.8514377	ESTs, Weakly similar to T2D3_HUMAN TRANSCRIPTION INITIATION FACTOR TFIID 135 KDA SUBUNIT [H.sapiens]

- 128 -

704414	0.8498631	small nuclear ribonucleoprotein polypeptides B and B1
2309073	0.8484971	frizzled (Drosophila) homolog 5
2052113	0.8477245	hypothetical protein FLJ10903
686172	0.846207	KIAA0008 gene product
150003	0.8447372	hypothetical protein FLJ13187
705064	0.8401441	transforming, acidic coiled-coil containing protein 3
1709791	0.8397779	BAI1-associated protein 1
1469425	0.8391993	SRY (sex determining region Y)-box 22
429799	0.8386406	hypothetical protein FLJ21939 similar to 5-azacytidine induced gene 2
729975	0.8385313	meningioma expressed antigen 5 (hyaluronidase)
1393018	0.8376676	general transcription factor IIIC, polypeptide 1 (alpha subunit, 220kD )
1492463	0.8360771	selenoprotein X, 1
2028949	0.8358024	hypothetical protein PRO1855
789012	0.8351735	fibulin 2
470124	0.8347241	RAD1 (S. pombe) homolog
1409509	0.8339967	troponin T1, skeletal, slow
1605426	0.8317254	hypothetical protein FLJ13352
44292	0.831599	Homo sapiens mRNA; cDNA DKFZp434C107 (from clone DKFZp434C107)
1435003	0.8311727	tumor suppressing subtransferable candidate 1
503215	0.8286483	pilin-like transcription factor
504308	0.8199799	hypothetical protein FLJ10540
785707	0.8176557	protein regulator of cytokinesis 1
1500162	0.8155519	ESTs
149355	0.8134342	translocating chain-associating membrane protein
1845169	0.8131362	RAB35, member RAS oncogene family
869375	0.808792	isocitrate dehydrogenase 2 (NADP+), mitochondrial
1492426	0.8032542	chromosome 19 open reading frame 3
782513	0.8030042	interferon, alpha-inducible protein (clone IFI-6-16)
813281	0.8016742	WW domain-containing protein 1
814378	0.8015236	serine protease inhibitor, Kunitz type, 2



- 129 -

700792	0.8005506	cyclin-dependent kinase inhibitor 3 (CDK2-associated dual specificity phosphatase)
340558	0.7990715	actin related protein 2/3 complex, subunit 5 (16 kD)
842994	0.7983201	cathepsin Z
138189	0.7953361	Wolfram syndrome 1 (wolframin)
289978	0.7931469	ubiquitin-like 4
2019223	0.7893602	mitochondrial ribosomal protein L17
2110511	0.7863117	artemin
814528	0.7845635	Homo sapiens cDNA: FLJ22139 fis, clone HEP20959
1573251	0.7816689	peroxisomal long-chain acyl-coA thioesterase
773922	0.7799164	KIAA0005 gene product
172517	0.7779159	hippocalcin-like 1
564981	0.7745626	ESTs
1420370	0.7726877	biliverdin reductase B (flavin reductase (NADPH))
308466	0.771216	GTP-binding protein Sara
199645	0.769591	nicastrin
1422338	0.7690604	ribonucleotide reductase M2 polypeptide
1474424	0.7689082	Homo sapiens cDNA FLJ12758 fis, clone NT2RP2001328
813751	0.7665427	sialyltransferase 4C (beta-galactosidase alpha-2,3-sialyltransferase)
131094	0.7626373	Homo sapiens cDNA: FLJ21587 fis, clone COL06946
842980	0.762412	developmentally regulated GTP-binding protein 1
212542	0.7585053	Homo sapiens cDNA FLJ12900 fis, clone NT2RP2004321
108425	0.7581954	
811761	0.7578696	Nijmegen breakage syndrome 1 (nibrin)
241348	0.757138	prenylcysteine lyase
810725	0.7561061	ATPase, H <sup>+</sup> transporting, lysosomal (vacuolar proton pump) 21kD
1460110	0.7559869	proteasome (prosome, macropain) subunit, beta type, 5
120749	0.7545697	ESTs, Moderately similar to KIAA1215 protein [H.sapiens]
827171	0.7543905	ESTs

- 130 -

39884	0.7526496	IMP (inosine monophosphate) dehydrogenase 1
124298	0.7507816	microsomal glutathione S-transferase 3
753378	0.7492212	hypothetical protein FLJ22649 similar to signal peptidase SPC22/23
210862	0.7483198	acyl-Coenzyme A oxidase 1, palmitoyl
785766	0.7476331	hypothetical protein
1553306	0.7471557	proteasome (prosome, macropain) 26S subunit, non-ATPase, 11
470099	0.7443747	HT002 protein; hypertension-related calcium- regulated gene
83363	0.7435058	protein-L-isoaspartate (D-aspartate) O- methyltransferase
489351	0.7422879	hypothetical protein DKFZp566J2046
810612	0.7419597	S100 calcium-binding protein A11 (calgizzarin)
825585	0.741258	tubulin-specific chaperone e
1456348	0.7401571	N-acetylneuraminic acid phosphate synthase; sialic acid synthase
1473922	0.7399948	actin related protein 2/3 complex, subunit 3 (21 kD)
150314	0.739762	lysophospholipase I
897806	0.7388301	hypoxia-inducible factor 1, alpha subunit (basic helix-loop-helix transcription factor)
811585	0.738212	huntingtin (Huntington disease)
685516	0.7378926	putative G protein-coupled receptor
76605	0.7376829	nesca protein
1476053	0.7367106	RAD51 (S. cerevisiae) homolog (E coli RecA homolog)
221295	0.7342275	inhibitor of DNA binding 2, dominant negative helix-loop-helix protein
824879	0.7333071	hypothetical protein MGC11275
768570	0.732628	hypothetical protein FLJ11280
73009	0.731401	ESTs, Weakly similar to A43932 mucin 2 precursor, intestinal [H.sapiens]
564492	0.7301895	mitochondrial carrier homolog 2
2017415	0.7281714	centromere protein A (17kD)
788654	0.7244749	
488505	0.7242619	accessory proteins BAP31/BAP29
824962	0.7238234	karyopherin alpha 2 (RAG cohort 1, importin alpha 1)

[illegible]

- 132 -

744417	0.6975761	carnitine acetyltransferase
814306	0.6963874	tumor protein D52
41356	0.6961169	protein phosphatase 2, regulatory subunit B (B56), alpha isoform
813419	0.6951349	hydroxyacyl-Coenzyme A dehydrogenase, type II
629944	0.6950339	myosin VB
327506	0.6943362	Homo sapiens mRNA full length insert cDNA clone EUROIMAGE 327506
768064	0.6936336	cytochrome P450, subfamily I (aromatic compound-inducible), polypeptide 1
770992	0.693225	
469383	0.6925975	chromosome 8 open reading frame 1
1492780	0.6923749	Homo sapiens cDNA FLJ14459 fis, clone HEMBB1002409
41569	0.691145	hypothetical protein FLJ12650
509588	0.6903363	TATA box binding protein (TBP)-associated factor, RNA polymerase II, J, 20kD
144880	0.6902434	hypothetical protein from EUROIMAGE 1759349
789376	0.6883473	thioredoxin reductase 1
268946	0.6881593	WD40 protein Ciao1
810156	0.6879064	deoxythymidylate kinase (thymidylate kinase)
84295	0.6876167	interleukin 1 receptor antagonist
246800	0.6867481	hypothetical protein FLJ10803
589232	0.6865999	hypothetical protein FLJ11506
859761	0.6846134	poliovirus receptor-related 2 (herpesvirus entry mediator B)
431505	0.6840493	ESTs, Highly similar to A31026 probable membrane receptor protein [H.sapiens]
109863	0.683291	epithelial membrane protein 2
770355	0.6829507	lanosterol synthase (2,3-oxidosqualene-lanosterol cyclase)
344091	0.6818771	ESTs
813707	0.6814476	regulator of G-protein signalling 16
124781	0.6809199	squalene epoxidase
502774	0.6807524	hypothetical protein FLJ20623
825740	0.6805157	DKFZp434J1813 protein
1536006	0.6805004	ESTs

- 133 -

203003	0.6802818	non-metastatic cells 4, protein expressed in
785616	0.6800658	signal sequence receptor, alpha (translocon-associated protein alpha)
785840	0.6800437	SEC24 ( <i>S. cerevisiae</i> ) related gene family, member D
256619	0.6797698	hydroxysteroid (17-beta) dehydrogenase 7
2108077	0.6796871	CGI-112 protein
742595	0.6784804	cyclin-dependent kinase 5
1476065	0.6783519	stathmin 1/oncoprotein 18
68636	0.6778817	hypothetical protein MGC2477
564847	0.6770455	ESTs
744047	0.6748419	polo ( <i>Drosophila</i> )-like kinase
897813	0.6741253	polyadenylate binding protein-interacting protein 1
823930	0.6738266	actin related protein 2/3 complex, subunit 1A (41 kD)
236034	0.673506	uncoupling protein 2 (mitochondrial, proton carrier)
743977	0.6729967	Homo sapiens mRNA for TL132
293727	0.6728542	hypothetical protein MGC861
590759	0.672734	sterol-C4-methyl oxidase-like
811024	0.6706275	bone marrow stromal cell antigen 2
785795	0.6683919	hypothetical protein FLJ12910
530310	0.6682042	KIAA0143 protein
1734309	0.666164	sperm associated antigen 4
1898619	0.6651268	hypothetical protein MGC15737
358267	0.6648183	EST, Moderately similar to AF119917 63 PRO2831 [ <i>H.sapiens</i> ]
753400	0.6633015	CGI-204 protein
292770	0.6609782	Homo sapiens, clone IMAGE:3627860, mRNA, partial cds
884425	0.6608337	chaperonin containing TCP1, subunit 5 (epsilon)
1696757	0.6599512	hypothetical protein KIAA1165
121251	0.6596925	hypothetical protein MGC5576
770785	0.658751	1,2-alpha-mannosidase IC
590774	0.657447	mitogen-activated protein kinase 13
969877	0.6564571	synaptosomal-associated protein, 25kD
415102	0.6562436	cell division cycle 25C
120271	0.6562017	hypothetical protein MGC4692

- 134 -

713782	0.6552297	a disintegrin and metalloproteinase domain 15 (metargidin)
49351	0.6551453	SEX gene
488642	0.6541915	ESTs, Weakly similar to I38022 hypothetical protein [H.sapiens]
343607	0.6510034	AD-015 protein
2306987	0.6501071	secreted and transmembrane 1
280249	0.64759	Kruppel-like factor 7 (ubiquitous)
752631	0.6473517	fibroblast growth factor receptor 3 (achondroplasia, thanatophoric dwarfism)
489594	0.645574	hypothetical protein FLJ11565
773188	0.6448143	nuclear receptor subfamily 1, group D, member 2
2015517	0.6441737	hypothetical protein FLJ22237
343731	0.6424907	
491524	0.6424448	mitochondrial ribosomal protein L13
824524	0.6424419	UDP-galactose transporter related
593431	0.6422897	ESTs, Moderately similar to CEGT_HUMAN CERAMIDE GLUCOSYLTRANSFERASE [H.sapiens]
812994	0.6420817	retinoid X receptor, alpha
1631735	0.6404771	Homo sapiens, clone IMAGE:3604336, mRNA, partial cds
278531	0.6404612	cytochrome c oxidase subunit VIc
2302099	0.6386686	sialidase 3 (membrane sialidase)
454896	0.6384516	DnaJ (Hsp40) homolog, subfamily A, member 2
1456701	0.6383709	B-cell CLL/lymphoma 9
2055807	0.638184	protein kinase domains containing protein similar to phosphoprotein C8FW
1518402	0.6378481	KIAA1361 protein
810762	0.6371461	SNARE protein
124447	0.6363079	KIAA1184 protein
49273	0.635859	solute carrier family 27 (fatty acid transporter), member 4
365060	0.6350631	RAB11A, member RAS oncogene family
1591264	0.6337293	transaldolase 1
41698	0.6327738	progesterone binding protein
810316	0.6323393	very long-chain acyl-CoA synthetase; lipidosin
826363	0.6322533	lysophospholipase II

- 135 -

2011515	0.6319712	DKFZP586B0923 protein
770675	0.6315109	Homo sapiens cDNA: FLJ21323 fis, clone COL02374
1461477	0.6300096	Homo sapiens mRNA; cDNA DKFZp586I0324 (from clone DKFZp586I0324)
366834	0.629867	envoplakin
1601947	0.6297475	cytochrome c oxidase subunit VIIa polypeptide 2 (liver)
510575	0.6295917	hypothetical protein FLJ22087
503851	0.6291771	nuclear receptor co-repressor/HDAC3 complex subunit
810063	0.6286184	growth factor, erv1 (S. cerevisiae)-like (augmenter of liver regeneration)
754653	0.6277098	cadherin, EGF LAG seven-pass G-type receptor 3, flamingo (Drosophila) homolog
1518890	0.6251884	metallothionein-like 5, testis-specific (tesmin)
784105	0.6244805	ESTs
205049	0.6231646	protein kinase H11; small stress protein-like protein HSP22
325606	0.6225147	hypothetical protein MGC14353
760299	-1.8441097	dickkopf (Xenopus laevis) homolog 3
200814	-1.8295958	membrane metallo-endopeptidase (neutral endopeptidase, enkephalinase, CALLA, CD10)
1882697	-1.715818	peanut (Drosophila)-like 2
344720	-1.6567437	glycophorin C (Gerbich blood group)
1161564	-1.5877154	desmuslin
45099	-1.566311	regucalcin (senescence marker protein-30)
75859	-1.5303427	N-myc downstream-regulated gene 2
811920	-1.5255258	interleukin 11 receptor, alpha
1569187	-1.4872982	heparan sulfate (glucosamine) 3-O-sulfotransferase 4
796542	-1.4697418	ets variant gene 5 (ets-related molecule)
767202	-1.4573536	latent transforming growth factor beta binding protein 2
285377	-1.4496786	pellino (Drosophila) homolog 2
300632	-1.449587	hypothetical protein FLJ21044 similar to Rbig1

- 136 -

160192	-1.4231519	ESTs, Weakly similar to 2004399A chromosomal protein [H.sapiens]
611532	-1.3689616	troponin I, skeletal, fast
813265	-1.3632094	Homo sapiens mRNA; cDNA DKFZp564H1916 (from clone DKFZp564H1916)
1469377	-1.359583	lipoma HMGIC fusion partner-like 2
810002	-1.3553375	Homo sapiens, clone MGC:19762 IMAGE:3636045, mRNA, complete cds
2056139	-1.3506352	LIM domain protein
377275	-1.3392122	ataxia-telangiectasia group D-associated protein
298122	-1.3364021	frizzled (Drosophila) homolog 7
72778	-1.3299927	caspase 7, apoptosis-related cysteine protease
841308	-1.3246996	myosin, light polypeptide kinase
377461	-1.3214357	caveolin 1, caveolae protein, 22kD
1473274	-1.3200174	myosin regulatory light chain 2, smooth muscle isoform
1554167	-1.3141843	hypothetical protein FLJ14529
131839	-1.3048208	folate receptor 1 (adult)
67741	-1.3014364	PP2135 protein
132857	-1.280579	Homo sapiens mRNA; cDNA DKFZp586N1323 (from clone DKFZp586N1323)
345670	-1.2805684	ESTs, Moderately similar to I59348 CCAAT binding transcription factor CBF subunit C - rat [R.norvegicus]
781014	-1.2659158	suppression of tumorigenicity 5
289760	-1.2610055	ESTs, Highly similar to T00391 hypothetical protein KIAA0612 [H.sapiens]
742685	-1.248683	disabled (Drosophila) homolog 2 (mitogen-responsive phosphoprotein)
52419	-1.2447753	Friedreich ataxia region gene X123
838478	-1.2349342	neurocalcin delta
839736	-1.2337192	crystallin, alpha B
1558675	-1.2313679	SRY (sex determining region Y)-box 10
796181	-1.2276581	growth arrest-specific 6
878836	-1.2200837	secretory granule, neuroendocrine protein 1 (7B2 protein)
130201	-1.2181641	intercellular adhesion molecule 2
1917449	-1.217371	serum amyloid A4, constitutive



- 137 -

22917	-1.2164705	Homo sapiens mRNA; cDNA DKFZp761M0111 (from clone DKFZp761M0111)
1871116	-1.209237	Homo sapiens mRNA; cDNA DKFZp434C1714 (from clone DKFZp434C1714); partial cds
823871	-1.2090693	SPARC-like 1 (mast9, hevin)
811837	-1.2030392	
811848	-1.2011809	hypothetical protein
840266	-1.200789	Homo sapiens cDNA: FLJ22667 fis, clone HSI08385
753071	-1.2005994	Homo sapiens cDNA: FLJ22528 fis, clone HRC12825
2106144	-1.1719133	regulated in glioma
2504881	-1.1674204	signal transducer and activator of transcription 5A
811837	-1.157587	eukaryotic translation elongation factor 1 alpha 1
486683	-1.1548164	Homo sapiens mRNA; cDNA DKFZp564J0323 (from clone DKFZp564J0323)
712139	-1.1491129	ADP-ribosylation factor-like 7
196435	-1.1475545	ESTs
877621	-1.1422087	nGAP-like protein
811088	-1.140941	ephrin-B3
322561	-1.1270333	ribosomal protein L31
712401	-1.1218827	phosphoinositide-3-kinase, catalytic, delta polypeptide
788234	-1.1061158	inhibitor of DNA binding 4, dominant negative helix-loop-helix protein
140574	-1.103064	small inducible cytokine subfamily D (Cys-X3- Cys), member 1 (fractalkine, neurotactin)
490023	-1.1020527	hypothetical protein MGC2648
1584540	-1.0933558	Homo sapiens mRNA; cDNA DKFZp586M2022 (from clone DKFZp586M2022)
1556433	-1.0887923	GRO3 oncogene
47043	-1.0841886	tensin
303109	-1.0807576	purinergic receptor (family A group 5)
343760	-1.0803279	SH3 domain binding glutamic acid-rich protein like 2

- 138 -

248631	-1.0784377	aminomethyltransferase (glycine cleavage system protein T)
740620	-1.071175	tropomyosin 2 (beta)
529843	-1.0698218	ESTs, Moderately similar to JC5238 galactosylceramide-like protein, GCP [H.sapiens]
990881	-1.0640807	
80344	-1.0602825	interleukin 7 receptor
625399	-1.0594452	hypothetical protein similar to beta-transducin family
416676	-1.0568729	pellino (Drosophila) homolog 1
416959	-1.0566462	nuclear factor I/B
307029	-1.0497874	ribosomal protein L26
488404	-1.0484323	Homo sapiens clone TUA8 Cri-du-chat region mRNA
291478	-1.0400846	runt-related transcription factor 3
153760	-1.0340645	EphB1
119290	-1.025797	cortic al thymocyte receptor (X. laevis CTX) like
814443	-1.0250673	hypothetical protein MGC3232
757191	-1.0238476	ESTs
208718	-1.0229324	annexin A1
161456	-1.0220494	serum amyloid A1
1587710	-1.0209983	period (Drosophila) homolog 1
160609	-1.0208819	ESTs
593023	-1.0158099	dystrobrevin, beta
1878409	-1.013872	catechol-O-methyltransferase
781017	-1.0125987	early growth response 2 (Krox-20 (Drosophila) homolog)
753162	-1.0119485	KIAA0603 gene product
897963	-1.0075423	phosphatidic acid phosphatase type 2A
505864	-1.0033263	RaIGDS-like gene
323780	-1.0027215	Homo sapiens cDNA FLJ11177 fis, clone PLACE1007402
343695	-1.002495	hypothetical protein FLJ10875
277571	-0.9977509	KIAA1706 protein
869450	-0.9934083	ribosomal protein L11
280907	-0.9932806	Kruppel-type zinc finger protein
290378	-0.9932643	podocalyxin-like
1635062	-0.9905245	DKFZP586A011 protein

- 139 -

270826	-0.9896888	Homo sapiens cDNA FLJ13329 fis, clone OVARC1001795
1257131	-0.9869928	ESTs

Example VII: Genes for discriminating between ADH and DCIS

As shown in Table 6 below, 350 genes were identified as being able to discriminate between ADH and DCIS. The actual data is shown in Figure 3.

**Table 6**

CloneID	Weight	Description
1404774	-0.5685596	parathyroid hormone-like hormone
823871	-0.9429443	SPARC-like 1 (mast9, hevin)
1882697	-1.4318896	peanut (Drosophila)-like 2
140071	-0.5708303	frizzled-related protein
160192	-1.1751869	ESTs, Weakly similar to 2004399A chromosomal protein [H.sapiens]
796542	-1.2625621	ets variant gene 5 (ets-related molecule)
611532	-1.3158379	troponin I, skeletal, fast
1473274	-1.1172693	myosin regulatory light chain 2, smooth muscle isoform
469306	-0.5113377	gastrin-releasing peptide
2306697	-0.4494025	neuromedin B
132857	-1.0084069	Homo sapiens mRNA; cDNA DKFZp586N1323 (from clone DKFZp586N1323)
2504881	-1.000863	signal transducer and activator of transcription 5A
760299	-1.7010462	dickkopf (Xenopus laevis) homolog 3
293819	-0.4051233	oxidoreductase UCPA
130835	0.3209345	Homo sapiens, Similar to clone FLB3816, clone IMAGE:3454380, mRNA
66532	-0.5773248	endothelin 3
2499829	-0.3619416	zinc finger protein 145 (Kruppel-like, expressed in promyelocytic leukemia)
85840	-0.7754435	nicotinamide N-methyltransferase

sd-71385

FOOTNOTES

- 140 -

859359 -0.4669761 quinone oxidoreductase homolog  
200814 -1.4325283 membrane metallo-endopeptidase (neutral endopeptidase,  
enkephalinase, CALLA, CD10)  
  
825287 -0.2783909 tumor necrosis factor (ligand) superfamily, member 11  
  
202577 -0.2283149 histamine N-methyltransferase  
2014373 0.0232775 HNK-1 sulfotransferase  
153760 -0.9227865 EphB1  
377275 -1.3102234 ataxia-telangiectasia group D-associated protein  
  
745490 -0.6329354 hypothetical protein FLJ20607  
172783 -0.6088873 hypothetical protein FLJ10390  
1558108 0.0044098 ATP-binding cassette, sub-family C (CFTR/MRP), member 8  
  
1587710 -0.8252065 period (Drosophila) homolog 1  
141731 -0.4531468  
490484 -0.3388325 ESTs  
504959 -0.1620065 Homo sapiens mRNA; cDNA DKFZp586G0321 (from clone  
DKFZp586G0321)  
  
1609746 -0.4792809 vitelliform macular dystrophy (Best disease, bestrophin)  
  
882248 -0.8700731 transgelin  
1917449 -1.0894686 serum amyloid A4, constitutive  
2119838 -0.3045374 a disintegrin-like and metalloprotease (repolysin type) with  
thrombospondin type 1 motif, 8  
  
841507 0.0247736 surfactant, pulmonary-associated protein A2  
  
813265 -1.2230435 Homo sapiens mRNA; cDNA DKFZp564H1916 (from clone  
DKFZp564H1916)  
  
767202 -1.0654145 latent transforming growth factor beta binding protein 2  
  
1156538 0.2252628 potassium inwardly-rectifying channel, subfamily J, member  
11  
39600 -0.6540275 adenylate kinase 5  
1630990 -0.7312981 ribosomal protein L29  
41208 -0.865227 bone morphogenetic protein 1  
322561 -0.7970805 ribosomal protein L31  
951008 -0.2221875 ESTs  
841308 -0.9953716 myosin, light polypeptide kinase

sd-71385

"FOOTNOTES" 8/20/2004

- 141 -

119290 -0.8224756 cortic al thymocyte receptor (X. laevis CTX) like  
344959 -0.0998837 gene for serine/threonine protein kinase  
810331 -0.1389612 quiescin Q6  
1161775 -0.7380353 villin 1  
625399 -0.9066615 hypothetical protein similar to beta-transducin family  
  
1470657 -0.3557985 deiodinase, iodothyronine, type II  
160609 -0.8405669 ESTs  
665356 -0.1575968 tumor necrosis factor receptor superfamily, member 11b  
(osteoprotegerin)  
  
1584540 -0.9754693 Homo sapiens mRNA; cDNA DKFZp586M2022 (from clone  
DKFZp586M2022)  
  
726779 -0.4459955 calponin 1, basic, smooth muscle  
296123 -0.2814132 Homo sapiens PRO1851 mRNA, complete cds  
  
190059 -0.634814 guanine nucleotide binding protein (G protein), gamma 7  
  
1471829 -0.7275378 uncharacterized hypothalamus protein HSMNP1  
  
2056139 -1.0673271 LIM domain protein  
190753 -0.4423669 ESTs  
1968422 -0.5802058 Homo sapiens mRNA full length insert cDNA clone  
EUROIMAGE 1968422  
  
725390 -0.4620278 glutathione S-transferase pi  
814826 -0.8443265 ESTs  
878836 -1.037352 secretory granule, neuroendocrine protein 1 (7B2 protein)  
  
27769 -0.8298821 ESTs  
188388 -0.6824191 integrin, alpha 10  
1597813 -0.5406089 hypothetical protein PP1044  
1609625 -0.5920312 selectin P ligand  
810981 -0.1805871 hypothetical protein FLJ20699  
726703 -0.070084 Homo sapiens clone 23736 mRNA sequence  
  
781014 -1.0161379 suppression of tumorigenicity 5  
898222 0.1795078 Homo sapiens clone 24418 mRNA sequence  
  
1475738 -0.6753091 ribosomal protein S25  
1492144 -0.2356446 butyrophilin, subfamily 3, member A2  
1492147 -0.636656 ribosomal protein S4, X-linked

- 142 -

1711456	-0.4896574 H factor (complement)-like 1
1871116	-0.9821709 Homo sapiens mRNA; cDNA DKFZp434C1714 (from clone DKFZp434C1714); partial cds
344720	-1.5196431 glycophorin C (Gerbich blood group)
45099	-1.3515907 regucalcin (senescence marker protein-30)
307029	-0.6500872 ribosomal protein L26
25763	-0.2725443 ankylosis, progressive (mouse) homolog
2502722	-0.3315871 loss of heterozygosity, 11, chromosomal region 2, gene A
1665444	-0.651994 tumor endothelial marker 1 precursor
165837	-0.3443143 translocase of inner mitochondrial membrane 22 (yeast) homolog
502518	-0.4342228 laminin, beta 2 (laminin S)
897963	-0.9775616 phosphatidic acid phosphatase type 2A
120138	0.1570107 J domain containing protein 1
78946	-0.648366 Homo sapiens pyruvate dehydrogenase kinase 4 mRNA, 3' untranslated region, partial sequence
990881	-0.8335353
859192	-0.2265317 vesicle-associated membrane protein 1 (synaptobrevin 1)
712023	-0.5302372 AT-binding transcription factor 1
855586	-0.3944697 nuclear receptor subfamily 3, group C, member 1
52419	-1.0368509 Friedreich ataxia region gene X123
432072	-0.4193898 nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 1
1897947	-0.163146 surfactant, pulmonary-associated protein A2
727229	-0.2668972 mitogen-activated protein kinase kinase kinase 4
810358	-0.5403705 acyl-Coenzyme A dehydrogenase, very long chain
154600	-0.0683999 phospholipase C, delta 1
755855	-0.6298333 solute carrier family 25 (mitochondrial carrier; adenine nucleotide translocator), member 6
323780	-0.8879854 Homo sapiens cDNA FLJ11177 fis, clone PLACE1007402
230560	-0.1803944 ESTs
283173	-0.4293696 EBP50-PDZ interactor of 64 kD

- 143 -

265853 -0.3519023 Homo sapiens cDNA FLJ11298 fis, clone PLACE1009794

161456 -0.8215502 serum amyloid A1

184022 -0.7070266 amyloid beta (A4) precursor protein-binding, family B,  
member 1 (Fe65)

1636523 -0.1496249 glutathione S-transferase subunit 13 homolog

795730 -0.2892776 signal transduction protein (SH3 containing)

1631682 -0.5162406 peptidylprolyl isomerase E (cyclophilin E)

811837 -0.848137

854763 -0.0493902 Homo sapiens cDNA FLJ11341 fis, clone PLACE1010786

50562 -0.0943363 chromosome 8 open reading frame 4

46843 -0.1736687 neuronal Shc adaptor homolog

178137 -0.6839022 ribosomal protein L34

1662279 -0.2290724 plakophilin 1 (ectodermal dysplasia/skin fragility syndrome)

1635062 -0.8243024 DKFZP586A011 protein

293916 -0.3671832 FKBP-associated protein

415613 -0.5528761 DHHC1 protein

80344 -0.8243418 interleukin 7 receptor

1602798 -0.4300808 choline kinase-like

1910516 -0.2845842 ESTs

740620 -0.96496 tropomyosin 2 (beta)

277627 -0.5329107 Human SH3 domain-containing protein SH3P18 mRNA,  
complete cds

854696 0.008949 siah binding protein 1; FBP interacting repressor; pyrimidine  
tract binding splicing factor; Ro ribonucleoprotein-binding  
protein 1

45578 -0.1295485 mitogen-activated protein kinase kinase 6

950710 -0.346361 propionyl Coenzyme A carboxylase, alpha polypeptide

768043 -0.4005598 ECSIT

308539 -0.6726187 Homo sapiens cDNA FLJ12777 fis, clone NT2RP2001720

595637 -0.5373697 Homo sapiens mRNA; cDNA DKFZp586N012 (from clone  
DKFZp586N012)

143661 -0.4713778 netrin 4

248631 -0.9106505 aminomethyltransferase (glycine cleavage system protein T)

- 144 -

826622	0.1339287 KIAA0430 gene product
51981	-0.5554908 ribosomal protein L7a
811848	-1.1350073 hypothetical protein
1533710	-0.5988282 ortholog of mouse integral membrane glycoprotein LIG-1
2017756	-0.2549438 homolog of yeast MOG1
1637296	-0.5105214 ribosomal protein S24
50586	-0.6794282 KIAA1545 protein
877835	-0.5382125 ribosomal protein L35
773319	-0.285447 ribosomal protein S6 kinase, 70kD, polypeptide 1
2014888	-0.2871529 sushi-repeat protein
767495	-0.0366213 GLI-Kruppel family member GLI3 (Greig cephalopolysyndactyly syndrome)
774078	-0.6508021 leiomodulin 1 (smooth muscle)
868400	-0.4928479 glutamyl-tRNA synthetase
240620	-0.5065007 vascular Rab-GAP/TBC-containing
1588791	-0.1606643 O-6-methylguanine-DNA methyltransferase
472186	-0.2687871 RAB32, member RAS oncogene family
725143	-0.3410957 hypothetical protein FLJ22418
714472	-0.3677387 KIAA0397 gene product
1854648	0.0089186 hemopexin
855029	0.076972 Ac-like transposable element
197727	-0.3906866 phosphatidylethanolamine N-methyltransferase
1623016	-0.6263561 EST
813841	0.0885625 plasminogen activator, tissue
2783721	-0.6854649 cholinergic receptor, nicotinic, beta polypeptide 1 (muscle)
549933	-0.3547881 interleukin 8
280907	-0.7307519 Kruppel-type zinc finger protein
796181	-0.8888496 growth arrest-specific 6
415233	-0.6506593 ribosomal protein L37a
811920	-1.2690713 interleukin 11 receptor, alpha
415415	-0.646492 EST, Moderately similar to RL1X_HUMAN 60S RIBOSOMAL PROTEIN L18A [H.sapiens]
2018807	-0.2989936 KIAA0468 gene product
154999	-0.3876757 hypothetical protein FLJ21007
68557	0.0637586 fatty acid binding protein 1, liver
647763	-0.1436566 ESTs



- 145 -

1568967 -0.2938468 ESTs  
210717 -0.4522432 syndecan 2 (heparan sulfate proteoglycan 1, cell surface-associated, fibroglycan)  
1350439 -0.6740833 KIAA0015 gene product  
684582 -0.0089016 tryptophanyl tRNA synthetase 2 (mitochondrial)  
122394 -0.049399 guanine nucleotide binding protein (G protein), alpha 15 (Gq class)  
213136 -0.1139004 BTG family, member 2  
625458 -0.2394796 hypothetical protein MGC3234  
811162 -0.3821116 fibromodulin  
1926246 -0.701769 ESTs, Moderately similar to T46371 hypothetical protein DKFZp434P1018.1 [H.sapiens]  
1554167 -1.0818834 hypothetical protein FLJ14529  
811088 -1.0851546 ephrin-B3  
72778 -1.1752838 caspase 7, apoptosis-related cysteine protease  
220293 -0.5078291 Homo sapiens cDNA: FLJ21800 fis, clone HEP00618  
1569187 -1.2971252 heparan sulfate (glucosamine) 3-O-sulfotransferase 4  
866866 -0.4801351 Ras association (RalGDS/AF-6) domain family 1  
81316 -0.6176265 ESTs  
1161564 -1.3891189 desmuslin  
898044 -0.4691863 metallocarboxypeptidase CPX-1  
293001 -0.4472585 hypothetical protein DKFZp434E2318  
1570502 -0.3314609 Homo sapiens cDNA FLJ12936 fis, clone NT2RP2005018  
839796 -0.6807912 candidate tumor suppressor p33 ING1 homolog  
380883 -0.4438655 Homo sapiens cDNA FLJ10158 fis, clone HEMBA1003463  
180561 -0.3250633 glutathione S-transferase M4  
869450 -0.7659922 ribosomal protein L11  
47043 -0.7729643 tensin  
810463 -0.0829995 DKFZP566O084 protein  
149539 0.4138942 KIAA1700  
200354 -0.0613983 thymidine kinase 2, mitochondrial  
771173 0.1757755 mitochondrial ribosomal protein S21  
270826 -0.9389651 Homo sapiens cDNA FLJ13329 fis, clone OVARC1001795

- 146 -

810017 -0.1076821 plasminogen activator, urokinase receptor  
809838 0.0956351  
344168 -0.386185 polymerase (DNA directed), lambda  
785967 -0.6049171 erythrocyte membrane protein band 4.1-like 2  
  
511831 -0.2912672 hypothetical protein MGC12936  
282404 -0.5528532 Homo sapiens mRNA for KIAA1671 protein, partial cds  
  
730036 -0.079736 Mad4 homolog  
298231 -0.3550893 gamma-aminobutyric acid (GABA) B receptor, 1  
  
2106144 -1.067765 regulated in glioma  
743880 -0.3262292 KIAA0263 gene product  
268234 -0.1535073 Dmx-like 1  
280776 -0.2922639 acyl-Coenzyme A dehydrogenase, C-2 to C-3 short chain  
  
212078 -0.1831462 integrin, alpha 1  
731308 0.5371204 citrate synthase  
774471 -0.1753902 laminin, beta 1  
811837 -0.8764231 eukaryotic translation elongation factor 1 alpha 1  
  
714437 0.0258281 sema domain, immunoglobulin domain (Ig), transmembrane  
domain (TM) and short cytoplasmic domain, (semaphorin) 4B  
  
416959 -0.957248 nuclear factor I/B  
681992 -0.7691299 Homo sapiens cDNA FLJ13384 fis, clone PLACE1001062,  
highly similar to Homo sapiens mRNA for lysine-  
ketoglutarate reductase/saccharopine dehydrogenase  
  
130201 -1.0401997 intercellular adhesion molecule 2  
363144 0.1937891 transcription factor AP-2 beta (activating enhancer-binding  
protein 2 beta)  
857874 -0.3703645 transforming growth factor beta-activated kinase-binding  
protein 1  
377461 -1.2798035 caveolin 1, caveolae protein, 22kD  
156363 0.2685238 hypothetical protein FLJ12934  
589115 -0.0115547 matrix metalloproteinase 1 (interstitial collagenase)  
  
241489 -0.6212938 adrenergic, beta-2-, receptor, surface  
586725 0.0683397 protein phosphatase 2, regulatory subunit B (B56), beta  
isoform

- 147 -

283124	-0.7933772 ESTs, Moderately similar to LONG-CHAIN FATTY ACID TRANSPORT PROTEIN [M.musculus]
307328	-0.4022704 hypothetical protein FLJ10948
37449	-0.3539978 GAS2-related on chromosome 22
50892	-0.6887497 Homo sapiens, clone MGC:9913 IMAGE:3870821, mRNA, complete cds
1897944	-0.4855156 ESTs, Weakly similar to T00366 hypothetical protein KIAA0669 [H.sapiens]
377314	-0.561312 casein kinase 2, alpha prime polypeptide
220851	-0.0351418 crystallin, alpha A
322233	-0.3025167
586803	-0.7423407 placental growth factor, vascular endothelial growth factor- related protein
343695	-0.7706809 hypothetical protein FLJ10875
414999	-0.5002542 ets variant gene 4 (E1A enhancer-binding protein, E1AF)
813823	-0.4463267 lumican
178825	-0.1525189 neurogranin (protein kinase C substrate, RC3)
85582	-0.0716296 target of myb1 (chicken) homolog-like 2
753620	-0.5248489 insulin-like growth factor binding protein 6
1899338	-0.066168 mannosidase, alpha, class 1A, member 2
359250	0.0674486 carbonic anhydrase IV
345034	-1.0600859 small inducible cytokine subfamily B (Cys-X-Cys), member 14 (BRAK)
2164744	-0.2307289 neural cell adhesion molecule 1
613056	-0.3957342 reticulocalbin 1, EF-hand calcium binding domain
1410444	-0.0368027 amphiregulin (schwannoma-derived growth factor)
593023	-0.7516846 dystrobrevin, beta
788511	0.1403148 ribosomal protein S6 kinase, 90kD, polypeptide 1
455269	0.0223607
1650927	-0.1712163 hypothetical protein DKFZp547E052
782427	-0.2818886 inhibin, beta B (activin AB beta polypeptide)
71727	-0.2796517 T-cell acute lymphocytic leukemia 1
840942	-0.6383764 major histocompatibility complex, class II, DP beta 1
825470	0.5218731 topoisomerase (DNA) II alpha (170kD)

- 148 -

595213 0.4824381 hypothetical protein  
796694 0.6893307 baculoviral IAP repeat-containing 5 (survivin)  
  
810711 0.9128832 stearyl-CoA desaturase (delta-9-desaturase)  
  
629944 0.5273447 myosin VB  
66406 0.5966701 hypothetical protein DKFZp762E1312  
785840 0.5401209 SEC24 (*S. cerevisiae*) related gene family, member D  
  
210862 0.4683996 acyl-Coenzyme A oxidase 1, palmitoyl  
84295 0.3989864 interleukin 1 receptor antagonist  
429182 0.2952867 dolichyl-phosphate mannosyltransferase polypeptide 1,  
catalytic subunit  
705064 0.4497555 transforming, acidic coiled-coil containing protein 3  
782513 0.4759571 interferon, alpha-inducible protein (clone IFI-6-16)  
839682 0.5208954 ubiquitin-conjugating enzyme E2N (homologous to yeast  
UBC13)  
768377 0.3809674 activity-dependent neuroprotective protein  
2309073 0.5226599 frizzled (*Drosophila*) homolog 5  
789376 0.4339171 thioredoxin reductase 1  
196992 -0.1015205 aldo-keto reductase family 1, member C1 (dihydrodiol  
dehydrogenase 1; 20-alpha (3-alpha)-hydroxysteroid  
dehydrogenase)  
785707 0.4708376 protein regulator of cytokinesis 1  
1505038 1.1904802 hypothetical protein FLJ20171  
869375 0.4827093 isocitrate dehydrogenase 2 (NADP+), mitochondrial  
  
109221 0.3048287 KIAA0286 protein  
345787 0.4647372 highly expressed in cancer, rich in leucine heptad repeats  
  
46248 0.640944 ADP-ribosyltransferase (NAD+; poly (ADP-ribose)  
polymerase)  
79520 0.6769129 RAB2, member RAS oncogene family  
469383 0.5558732 chromosome 8 open reading frame 1  
509588 0.4157059 TATA box binding protein (TBP)-associated factor, RNA  
polymerase II, J, 20kD  
  
292936 0.4217115 hypothetical protein FLJ10468  
686172 0.6724684 KIAA0008 gene product  
788655 0.2976191 HTPAP protein

- 149 -

770992	0.3331315
1587847	0.1732392 minichromosome maintenance deficient (mis5, <i>S. pombe</i> ) 6
79710	0.4579008 KIAA0174 gene product
2056566	0.2341905 integrin, alpha M (complement component receptor 3, alpha; also known as CD11b (p170), macrophage antigen alpha polypeptide)
454339	0.3738929 thiopurine S-methyltransferase
594226	0.403676 Homo sapiens cDNA FLJ14459 fis, clone HEMBB1002409
897813	0.4749348 polyadenylate binding protein-interacting protein 1
824694	0.0268936 protein tyrosine phosphatase type IVA, member 1
725454	0.6669973 CDC28 protein kinase 2
79254	0.0732432 MHC class I region ORF
1472719	0.2550065 SMT3 (suppressor of mif two 3, yeast) homolog 1
2054635	0.7914755 proteasome (prosome, macropain) subunit, alpha type, 7
289978	0.6308861 ubiquitin-like 4
155806	0.2747101 2'-5'-oligoadenylate synthetase 2 (69-71 kD)
147834	0.4412795 zinc finger protein 217
26171	0.3656147 KIAA0856 protein
2322367	0.5832711 reticulon 4
769921	0.6464245 ubiquitin carrier protein E2-C
73009	0.4822587 ESTs, Weakly similar to A43932 mucin 2 precursor, intestinal [H.sapiens]
2014034	0.6620303 methylene tetrahydrofolate dehydrogenase (NAD+ dependent), methenyltetrahydrofolate cyclohydrolase
359887	0.8692353 translocase of inner mitochondrial membrane 17 (yeast) homolog A
488964	0.9569176 H2A histone family, member O
1476065	0.289904 stathmin 1/oncoprotein 18
1422338	0.5874097 ribonucleotide reductase M2 polypeptide
268946	0.5440003 WD40 protein Ciao1
686552	0.7231319 golgi phosphoprotein 1
149544	0.283642 neuroepithelial cell transforming gene 1
770675	0.454729 Homo sapiens cDNA: FLJ21323 fis, clone COL02374

346257	0.1515899 minichromosome maintenance deficient ( <i>S. cerevisiae</i> ) 4
429799	0.6915155 hypothetical protein FLJ21939 similar to 5-azacytidine induced gene 2
143997	0.1883447 proteasome (prosome, macropain) 26S subunit, non-ATPase, 10
122241	0.7394284 proteasome (prosome, macropain) subunit, beta type, 2
823598	0.9153521 proteasome (prosome, macropain) 26S subunit, non-ATPase, 12
814632	0.0960611 splicing factor, arginine/serine-rich (transformer 2 <i>Drosophila</i> homolog) 10
810316	0.3732635 very long-chain acyl-CoA synthetase; lipidosin
290841	0.8346933 H2B histone family, member A
347373	0.7563599 transcription elongation factor B (SIII), polypeptide 1 (15kD, elongin C)
700792	0.4949149 cyclin-dependent kinase inhibitor 3 (CDK2-associated dual specificity phosphatase)
1554549	0.9764206 hydroxyacyl glutathione hydrolase
897770	0.4270685
504308	0.5564295 hypothetical protein FLJ10540
132828	0.2875519 Down syndrome critical region gene 1-like 2
292388	0.112577
1616253	0.567837 breast carcinoma amplified sequence 1
796469	1.0269115 HSPC150 protein similar to ubiquitin-conjugating enzyme
2139152	-0.4373776 Homo sapiens clone 24473 mRNA sequence
130276	0.2154295 Homo sapiens mRNA; cDNA DKFZp586H0324 (from clone DKFZp586H0324)
34149	0.309569 KIAA0227 protein
150003	0.8251408 hypothetical protein FLJ13187
287749	0.3173706 CDC7 (cell division cycle 7, <i>S. cerevisiae</i> , homolog)-like 1
25380	0.164077 ESTs, Moderately similar to JC5238 galactosylceramide-like protein, GCP [ <i>H.sapiens</i> ]
745083	0.4107735 ubiquitin specific protease 18
810899	0.240336 CDC28 protein kinase 1
768059	0.3139886 hypothetical protein FLJ12619

- 151 -

815026	0.4472842	cleavage stimulation factor, 3' pre-RNA, subunit 3, 77kD
741474	0.7732047	glucose phosphate isomerase
788641	0.3638059	adaptor-related protein complex 1, sigma 2 subunit
2043167	0.7167431	BCL2-associated athanogene 3
1492780	0.6544659	Homo sapiens cDNA FLJ14459 fis, clone HEMBB1002409
2306987	0.4402912	secreted and transmembrane 1
754653	0.4405873	cadherin, EGF LAG seven-pass G-type receptor 3, flamingo (Drosophila) homolog
743810	0.3667087	hypothetical protein MGC2577
126858	0.2389814	sterol O-acyltransferase (acyl-Coenzyme A: cholesterol acyltransferase) 1
2017415	0.43546	centromere protein A (17kD)
280507	0.2846518	hypoxanthine phosphoribosyltransferase 1 (Lesch-Nyhan syndrome)
67237	0.1406869	ESTs
43833	-0.1197546	diacylglycerol kinase, gamma (90kD)
42076	0.3194462	TRK-fused gene
531319	0.1093932	serine/threonine kinase 12
784129	0.2902598	tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein, gamma polypeptide
586895	0.2549108	small nuclear ribonucleoprotein polypeptide G

#### Example VIII: Genes for discriminating between DCIS Grade I and Grade III

As shown in Table 7 below, 350 genes were identified as being able to discriminate between grades of DCIS. The actual data corresponding to this table is shown in Figure 4.

**Table 7**

CloneID	Weight	Description
2460159	4.6268975	tyrosine kinase, non-receptor, 1
358151	2.8783989	zinc finger protein 33a (KOX 31)

795382	2.772572	Rap1 guanine-nucleotide-exchange factor directly activated by cAMP
714472	2.6829714	KIAA0397 gene product
725649	2.6433625	nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 4
51218	2.6200167	ESTs
504959	2.5777963	Homo sapiens mRNA; cDNA DKFZp586G0321 (from clone DKFZp586G0321)
647397	2.5320153	ESTs
814815	2.4198892	plakophilin 4
279720	2.4136316	ESTs, Moderately similar to A47582 B-cell growth factor precursor [H.sapiens]
298231	2.4096172	gamma-aminobutyric acid (GABA) B receptor, 1
172783	2.3339699	hypothetical protein FLJ10390
261609	2.311051	ESTs, Weakly similar to S65657 alpha-1C-adrenergic receptor splice form 2 [H.sapiens]
826668	2.2951137	KIAA0274 gene product
1493383	2.2616737	Homo sapiens mRNA; cDNA DKFZp434H2418 (from clone DKFZp434H2418)
2017756	2.2403622	homolog of yeast MOG1
1455566	2.1756946	adenosine A3 receptor
725321	2.1658095	estrogen receptor 1
180561	2.1393944	glutathione S-transferase M4
32050	2.1165022	Homo sapiens mRNA; cDNA DKFZp586P1124 (from clone DKFZp586P1124)
215000	2.1150123	vasoactive intestinal peptide receptor 1
2019750	2.0957741	SEC14 (S. cerevisiae)-like 2
283124	2.0921779	ESTs, Moderately similar to LONG-CHAIN FATTY ACID TRANSPORT PROTEIN [M.musculus]
490615	2.0635424	tubulin, gamma 2
666138	2.0625667	hypothetical protein DKFZp761J1523
418129	2.0541212	nuclear mitotic apparatus protein 1
1733262	2.052666	BLu protein
1588791	2.044069	O-6-methylguanine-DNA methyltransferase
461761	2.035953	angiogenin, ribonuclease, RNase A family, 5



- 153 -

1031592 1.9930012 kinesin protein 9 gene  
126415 1.9781057 Homo sapiens mRNA; cDNA DKFZp566H0124 (from clone  
DKFZp566H0124)  
  
28643 1.9733043 hypothetical protein DKFZp564D1378  
470261 1.9160486  
1630990 1.9155421 ribosomal protein L29  
810981 1.9132205 hypothetical protein FLJ20699  
767495 1.9052671 GLI-Kruppel family member GLI3 (Greig  
cephalopolysyndactyly syndrome)  
767176 1.8996792 tumor necrosis factor (ligand) superfamily, member 13  
  
1572196 1.8855996 secreted modular calcium-binding protein 2  
  
1706635 1.8754346 bone gamma-carboxyglutamate (gla) protein (osteocalcin)  
  
186301 1.8424458 Homo sapiens cDNA FLJ12924 fis, clone NT2RP2004709  
  
726703 1.842399 Homo sapiens clone 23736 mRNA sequence  
  
214205 1.8416657 Homo sapiens, clone MGC:17687 IMAGE:3865868, mRNA,  
complete cds  
784178 1.822159 Homo sapiens mRNA; cDNA DKFZp586M0723 (from clone  
DKFZp586M0723)  
  
346902 1.8196723 ESTs, Weakly similar to ALU2\_HUMAN ALU SUBFAMILY SB  
SEQUENCE CONTAMINATION WARNING ENTRY  
[H.sapiens]  
  
1367678 1.8184255 KIAA0356 gene product  
190059 1.7988757 guanine nucleotide binding protein (G protein), gamma 7  
  
1456937 1.7980645 oviductal glycoprotein 1, 120kD (mucin 9, oviductin)  
  
45578 1.7935215 mitogen-activated protein kinase kinase 6  
248631 1.7701919 aminomethyltransferase (glycine cleavage system protein T)  
  
1562231 1.7668811 SET binding protein 1  
154466 1.7516237 STIP1 homology and U-Box containing protein 1  
  
2524445 1.7392197 neuronal PAS domain protein 1  
277266 1.7307752 Homo sapiens, clone IMAGE:3625550, mRNA, partial cds  
  
741891 1.7297992 RAB2, member RAS oncogene family-like

- 154 -

206217 1.7227658 nuclear receptor subfamily 1, group H, member 3

2028876 1.7128178 splicing factor, arginine/serine-rich 5  
730036 1.70573 Mad4 homolog

1558233 1.7035129 ESTs  
502518 1.7027287 laminin, beta 2 (laminin S)  
356835 1.7014349 hypothetical protein MGC10500  
744994 1.6938348 hypothetical protein FLJ12242  
810358 1.6924263 acyl-Coenzyme A dehydrogenase, very long chain

813854 1.6914133 purine-rich element binding protein A  
768043 1.6854123 ECSIT  
264632 1.6591646 ESTs  
505243 1.6581518 inositol 1,4,5-triphosphate receptor, type 2  
344073 1.6571604 ESTs, Weakly similar to K1CI\_HUMAN KERATIN, TYPE I  
CYTOSKELETAL 9 [H.sapiens]

343760 1.6435915 SH3 domain binding glutamic acid-rich protein like 2

1569418 1.6408767 Homo sapiens cDNA FLJ11385 fis, clone HEMBA1000520

202577 1.6393518 histamine N-methyltransferase  
823634 1.6366047 ESTs  
839796 1.630887 candidate tumor suppressor p33 ING1 homolog

183440 1.6297915 arylsulfatase A  
344959 1.6268043 gene for serine/threonine protein kinase  
1574252 1.6211224 DKFZP586D0623 protein  
669359 1.6174855 Homo sapiens clone 24405 mRNA sequence

74070 1.6096611 endosulfine alpha  
1660649 1.6051582 suppressor of white apricot homolog 2  
796723 1.6041219 Homo sapiens clone CDABP0014 mRNA sequence

743146 1.597094 hypothetical protein FLJ23403  
789147 1.5937127 enolase 2, (gamma, neuronal)  
342181 1.5926355 B-cell CLL/lymphoma 2  
33076 1.5824352 cholinephosphotransferase 1  
782497 1.5804781 Homo sapiens, clone IMAGE:3010666, mRNA, partial cds

470261 1.5751564 SMA5  
809507 1.5723669 hypothetical protein FLJ20568  
712460 1.5721924 natural killer-tumor recognition sequence

sd-71385

10020018-123101

- 155 -

293569 1.5662317 chromosome 1 open reading frame 21  
1518402 1.5627158 KIAA1361 protein  
155072 1.5557559 ESTs  
455269 1.5538593  
262804 1.5532348 hypothetical protein MGC2941  
1492238 1.5484044 HSPC003 protein  
364865 1.5483958 hypothetical protein FLJ21062  
2325804 1.5461823 95 kDa retinoblastoma protein binding protein  
  
1635062 1.5449944 DKFZP586A011 protein  
344168 1.5421405 polymerase (DNA directed), lambda  
1517171 1.5381921 interleukin 2 receptor, alpha  
769600 1.5335878 uracil-DNA glycosylase 2  
325583 1.5245817 EST  
814826 1.5223296 ESTs  
1570502 1.5184543 Homo sapiens cDNA FLJ12936 fis, clone NT2RP2005018  
  
511831 1.5183298 hypothetical protein MGC12936  
124922 1.5052004 KRAB-zinc finger protein SZF1-1  
2072768 1.4886791 nuclear receptor coactivator 3  
2021882 1.4854251 sodium channel, nonvoltage-gated 1, delta  
627248 1.4827018 SBBI31 protein  
725503 1.476461 D-dopachrome tautomerase  
285312 1.4699425  
141731 1.469088  
1456701 1.4668054 B-cell CLL/lymphoma 9  
898222 1.4667947 Homo sapiens clone 24418 mRNA sequence  
  
725284 1.4638006 phosphorylase kinase, gamma 2 (testis)  
154999 1.4564204 hypothetical protein FLJ21007  
1592530 1.4558873 mammalian inositol hexakisphosphate kinase 2  
  
590310 1.4534138 Homo sapiens mRNA; cDNA DKFZp434E2321 (from clone  
DKFZp434E2321); partial cds  
  
416042 1.446691 ESTs, Weakly similar to ALU1\_HUMAN ALU SUBFAMILY J  
SEQUENCE CONTAMINATION WARNING ENTRY  
[H.sapiens]  
  
2017144 1.4450059 CGI-41 protein  
309449 1.4412699 ribosomal protein S4, Y-linked  
450301 1.4403678 mutL (E. coli) homolog 3

- 156 -

1492468 1.4375076 DEME-6 protein  
121454 1.4361788 arachidonate 12-lipoxygenase  
743880 1.4294838 KIAA0263 gene product  
1568967 1.4109448 ESTs  
1568989 1.4094472 ESTs  
25274 1.4078252 g20 protein  
2413337 1.4058288 sortilin-related receptor, L(DLR class) A repeats-containing  
  
197913 1.3994468 splicing factor proline/glutamine rich (polypyrimidine tract-  
binding protein-associated)  
  
292770 1.3952057 Homo sapiens, clone IMAGE:3627860, mRNA, partial cds  
  
143332 1.3875805 neuropeptide Y receptor Y1  
809779 1.3848708 KIAA0239 protein  
138242 1.3820433 ESTs, Moderately similar to MAS2\_HUMAN MANNAN-  
BINDING LECTIN SERINE PROTEASE 2 PRECURSOR  
[H.sapiens]  
  
270127 1.3786197  
49240 1.3772023 KIAA0460 protein  
826622 1.3734143 KIAA0430 gene product  
1858837 1.3719341 ESTs  
1583198 1.3712112 ESTs, Weakly similar to S65824 reverse transcriptase  
homolog [H.sapiens]  
345858 1.3629187 cisplatin resistance associated  
208387 1.3597205 KIAA1407 protein  
502782 1.3589088 RAN binding protein 3  
26294 1.3578289 RNB6  
669379 1.3559931 Homo sapiens BAC clone RP11-505D17 from 7p22-p21  
  
810728 1.355262 hypothetical gene ZD52F10  
1601845 1.3533864 Ca<sup>2+</sup>-promoted Ras inactivator  
840882 1.350538 nucleotide binding protein  
82173 1.3485351 MYLE protein  
490965 1.3476943 ESTs  
811162 1.3450626 fibromodulin  
61061 1.3437368 hypothetical protein FLJ20585  
49630 1.3412751 calcium channel, voltage-dependent, L type, alpha 1D subunit  
  
825659 -2.8345933 N-myc downstream regulated  
769921 -2.712812 ubiquitin carrier protein E2-C

- 157 -

788232 -2.6550962 sestrin 2  
1476053 -2.529039 RAD51 (S. cerevisiae) homolog (E coli RecA homolog)  
809557 -2.4910488 minichromosome maintenance deficient (S. cerevisiae) 3  
150897 -2.4302462 UDP-GlcNAc:betaGal beta-1,3-N-  
acetylglucosaminyltransferase 3  
814526 -2.37273 seb4D  
796694 -2.3104815 baculoviral IAP repeat-containing 5 (survivin)  
814792 -2.3001334 ubiquitin specific protease 10  
1536236 -2.2971152 hypothetical protein FLJ13154  
2017415 -2.2748559 centromere protein A (17kD)  
210862 -2.2408114 acyl-Coenzyme A oxidase 1, palmitoyl  
795543 -2.2403389 thioredoxin peroxidase (antioxidant enzyme)  
773301 -2.2239527 cadherin 3, type 1, P-cadherin (placental)  
700792 -2.2203557 cyclin-dependent kinase inhibitor 3 (CDK2-associated dual  
specificity phosphatase)  
1883327 -2.2093248 ESTs  
1523225 -2.1936598 oncostatin M receptor  
208718 -2.1882654 annexin A1  
1702742 -2.1567809 solute carrier family 7 (cationic amino acid transporter, y+  
system), member 5  
753378 -2.151145 hypothetical protein FLJ22649 similar to signal peptidase  
SPC22/23  
429222 -2.1351022 CGI-107 protein  
292388 -2.1343367  
781047 -2.1262682 budding uninhibited by benzimidazoles 1 (yeast homolog)  
450854 -2.0819596 craniofacial development protein 1  
823598 -2.0693353 proteasome (prosome, macropain) 26S subunit, non-ATPase,  
12  
1422338 -2.0689327 ribonucleotide reductase M2 polypeptide  
705064 -2.0601901 transforming, acidic coiled-coil containing protein 3  
770675 -2.0272207 Homo sapiens cDNA: FLJ21323 fis, clone COL02374  
345787 -1.9992897 highly expressed in cancer, rich in leucine heptad repeats  
471196 -1.9985751 integral membrane protein 3

- 158 -

753215 -1.9954476 guanine nucleotide binding protein (G protein), alpha inhibiting  
activity polypeptide 1

869375 -1.9803724 isocitrate dehydrogenase 2 (NADP+), mitochondrial

842818 -1.9796507 lysyl-tRNA synthetase

229579 -1.9782493 Golgi apparatus protein 1

1916461 -1.9736332 hypothetical protein

129294 -1.9730334 ESTs

789376 -1.9719575 thioredoxin reductase 1

1897302 -1.9681938 serine/threonine kinase 17b (apoptosis-inducing)

951117 -1.9577807 eukaryotic translation elongation factor 1 gamma

1642496 -1.9484342 hypothetical protein MGC11266

591465 -1.9277062 Homo sapiens, clone MGC:2908 IMAGE:3029644, mRNA,  
complete cds

149355 -1.8999907 translocating chain-associating membrane protein

259950 -1.8958779 hypothetical protein FLJ14991

897770 -1.8887034

878798 -1.8774101 beta-2-microglobulin

746229 -1.873434 mitogen-activated protein kinase kinase kinase kinase 4

624867 -1.8594915 hypothetical protein FLJ20186

504308 -1.8590049 hypothetical protein FLJ10540

727251 -1.8553757 CD9 antigen (p24)

897774 -1.8532125 adenine phosphoribosyltransferase

1901310 -1.8434048 KIAA1209 protein

292936 -1.838542 hypothetical protein FLJ10468

1518591 -1.8382765

321354 -1.8123042 mitochondrial ribosomal protein L15

1903066 -1.8071117 keratin, hair, basic, 1

235180 -1.8004039 ESTs, Moderately similar to ALU1\_HUMAN ALU SUBFAMILY  
J SEQUENCE CONTAMINATION WARNING ENTRY  
[H.sapiens]

740604 -1.7995578 interferon stimulated gene (20kD)

233464 -1.7978669 epithelial V-like antigen 1

951241 -1.7961406 clone HQ0310 PRO0310p1

1587847 -1.7886092 minichromosome maintenance deficient (mis5, S. pombe) 6

624390 -1.7800152 DC13 protein

sd-71385

- 159 -

128711 -1.7791673 anillin (Drosophila Scraps homolog), actin binding protein  
308633 -1.7667341 hypothetical protein FLJ10339  
884425 -1.761486 chaperonin containing TCP1, subunit 5 (epsilon)  
745394 -1.7583344 Homo sapiens cDNA: FLJ23249 fis, clone COL04196  
852829 -1.7579704 karyopherin alpha 3 (importin alpha 4)  
122241 -1.7490794 proteasome (prosome, macropain) subunit, beta type, 2  
307255 -1.7294781 basement membrane-induced gene  
32493 -1.7273675 integrin, alpha 6  
454896 -1.7216328 DnaJ (Hsp40) homolog, subfamily A, member 2  
1876217 -1.7186822 DnaJ (Hsp40) homolog, subfamily A, member 2  
815556 -1.7161709 hypothetical protein FLJ10430  
589869 -1.7120206 transcriptional co-activator with PDZ-binding motif (TAZ)  
2012523 -1.7104314 fatty acid binding protein 5 (psoriasis-associated)  
1946448 -1.7092537 caveolin 2  
841370 -1.7047297 glutamic-oxaloacetic transaminase 2, mitochondrial (aspartate  
aminotransferase 2)  
1600239 -1.6989076 HSPC037 protein  
462926 -1.6981004 NIMA (never in mitosis gene a)-related kinase 2  
144880 -1.6980733 hypothetical protein from EUROIMAGE 1759349  
744047 -1.6910803 polo (Drosophila)-like kinase  
624627 -1.6888444 ribonucleotide reductase M2 polypeptide  
788566 -1.6752673 Purkinje cell protein 4  
858293 -1.6732706 tyrosine 3-monooxygenase/tryptophan 5-monooxygenase  
activation protein, theta polypeptide  
470148 -1.659943 ESTs  
66902 -1.6546363 ESTs  
77533 -1.6505418 inositol polyphosphate-5-phosphatase, 40kD  
825282 -1.6417354 DKFZP586L0724 protein  
785840 -1.6406565 SEC24 (S. cerevisiae) related gene family, member D

- 160 -

824962 -1.6399941 karyopherin alpha 2 (RAG cohort 1, importin alpha 1)  
741139 -1.6396499 eyes absent (Drosophila) homolog 2  
809530 -1.6264151 minichromosome maintenance deficient (S. cerevisiae) 2  
(mitotin)  
2054635 -1.6254343 proteasome (prosome, macropain) subunit, alpha type, 7  
813533 -1.6215197 syndecan binding protein (syntenin)  
267816 -1.6173276 KIAA0551 protein  
306318 -1.6083862 origin recognition complex, subunit 6 (yeast homolog)-like  
796469 -1.6065509 HSPC150 protein similar to ubiquitin-conjugating enzyme  
813256 -1.6027042 ATP-binding cassette, sub-family B (MDR/TAP), member 1  
843121 -1.60128 chloride intracellular channel 1  
713685 -1.6006729 protease, serine, 2 (trypsin 2)  
811590 -1.6001102 hypothetical protein FLJ11100  
823756 -1.5995448 mitogen inducible 2  
549073 -1.597146 capping protein (actin filament) muscle Z-line, alpha 2  
743810 -1.5960999 hypothetical protein MGC2577  
377368 -1.5955102 cell death regulator aven  
1506046 -1.5918448 hypothetical protein FLJ10815  
814270 -1.5890389 polymyositis/scleroderma autoantigen 1 (75kD)  
1604703 -1.5853167 major histocompatibility complex, class I, F  
234736 -1.5765159 GATA-binding protein 6  
429182 -1.5699689 dolichyl-phosphate mannosyltransferase polypeptide 1,  
catalytic subunit  
826355 -1.5677013 vesicle-associated membrane protein 5 (myobrevin)  
34149 -1.5659923 KIAA0227 protein  
746190 -1.5635857 hypothetical protein DKFZp761B1514  
131091 -1.558448 Not56 (D. melanogaster)-like protein  
665384 -1.5550415 KIAA1609 protein  
625683 -1.5501423 neighbor of COX4  
1582738 -1.544427 uncharacterized bone marrow protein BM040  
949988 -1.5437326 Homo sapiens mRNA; cDNA DKFZp586E1124 (from clone  
DKFZp586E1124); complete cds



- 161 -

133213 -1.5433392 fucosyltransferase 4 (alpha (1,3) fucosyltransferase, myeloid-specific)  
1660666 -1.5423907 carbonic anhydrase VB, mitochondrial  
713158 -1.5372729 ESTs  
1614140 -1.5341302 Ris  
201890 -1.5333626 baculoviral IAP repeat-containing 3  
38925 -1.5304604 ESTs, Moderately similar to A47582 B-cell growth factor precursor [H.sapiens]  
511850 -1.5294752 proteasome (prosome, macropain) 26S subunit, ATPase, 1  
489489 -1.5169076 lamin B receptor  
825470 -1.5154306 topoisomerase (DNA) II alpha (170kD)  
42831 -1.5136459 N-terminal kinase-like  
809784 -1.5066053 kallikrein 6 (neurosin, zyme)  
730410 -1.506168 lymphocyte-specific protein tyrosine kinase  
810983 -1.5060386 DKFZP434H132 protein  
731223 -1.5058799 proteasome (prosome, macropain) subunit, beta type, 2  
259017 -1.5053827 ESTs  
340745 -1.5041978 ESTs  
746163 -1.5000358 ESTs, Weakly similar to ALU1\_HUMAN ALU SUBFAMILY J  
SEQUENCE CONTAMINATION WARNING ENTRY  
[H.sapiens]  
1466621 -1.4998321 ATPase, Ca++ transporting, type 2C, member 1  
809588 -1.49754 gamma-glutamyl hydrolase (conjugase,  
folylpolygammaglutamyl hydrolase)  
813707 -1.4946872 regulator of G-protein signalling 16  
188335 -1.4895328 egf-like module containing, mucin-like, hormone receptor-like  
sequence 2  
1493160 -1.4851884 small inducible cytokine subfamily B (Cys-X-Cys), member 10  
531886 -1.4842053 Homo sapiens clone FLB3344 PRO0845 mRNA, complete cds  
753428 -1.4835152 Homo sapiens, clone IMAGE:3542597, mRNA, partial cds  
897731 -1.4833877 latrophilin  
376551 -1.4811859 ETAA16 protein  
256907 -1.4755682 glutathione S-transferase A3  
712139 -1.4735772 ADP-ribosylation factor-like 7

- 162 -

785368 -1.4680554 PDZ-binding kinase; T-cell originated protein kinase  
825606 -1.4667088 kinesin-like 1  
531319 -1.4659871 serine/threonine kinase 12  
66406 -1.4641179 hypothetical protein DKFZp762E1312  
470124 -1.4635711 RAD1 (S. pombe) homolog

Example IX: Genes for discriminating between IDC Grade I and Grade III

As shown in Table 8 below, 300 genes were identified as being able to discriminate between two grades of IDC.

**Table 8**

<b>CloneID</b>	<b>Weight</b>	<b>Description</b>
1706635	3.279305	bone gamma-carboxyglutamate (gla) protein (osteocalcin)
666138	3.1366533	hypothetical protein DKFZp761J1523
795382	3.0005724	Rap1 guanine-nucleotide-exchange factor directly activated by cAMP
364865	2.8512705	hypothetical protein FLJ21062
1500542	2.8021264	regulator of G-protein signalling 11
270127	2.7455118	
588262	2.7127609	Homo sapiens, Similar to RIKEN cDNA 2600001A11 gene, clone MGC:9907 IMAGE:3870073, mRNA, complete cds
277266	2.5004627	Homo sapiens, clone IMAGE:3625550, mRNA, partial cds
2090129	2.4882736	chromobox homolog 2 (Drosophila Pc class)
325583	2.4558352	EST
358151	2.4309122	zinc finger protein 33a (KOX 31)
786675	2.3877233	epididymis-specific, whey-acidic protein type, four-disulfide core; putative ovarian carcinoma marker
854763	2.378956	Homo sapiens cDNA FLJ11341 fis, clone PLACE1010786
1592976	2.3417116	microphthalmia-associated transcription factor

sd-71385

40020019 122401

- 163 -

1562231 2.286418 SET binding protein 1  
767176 2.2472282 tumor necrosis factor (ligand) superfamily, member 13  
  
784178 2.2026739 Homo sapiens mRNA; cDNA DKFZp586M0723 (from clone  
DKFZp586M0723)  
  
811162 2.1926501 fibromodulin  
502518 2.1886432 laminin, beta 2 (laminin S)  
754429 2.1363368 ESTs, Weakly similar to T00084 hypothetical protein  
KIAA0512 [H.sapiens]  
214205 2.1127961 Homo sapiens, clone MGC:17687 IMAGE:3865868, mRNA,  
complete cds  
61061 2.1046094 hypothetical protein FLJ20585  
344959 2.0756965 gene for serine/threonine protein kinase  
814815 2.0694714 plakophilin 4  
262804 2.0673444 hypothetical protein MGC2941  
1455566 2.0489061 adenosine A3 receptor  
1469149 1.9778151 Homo sapiens clone 24606 mRNA sequence  
  
358217 1.9703763 glypican 4  
206217 1.9595484 nuclear receptor subfamily 1, group H, member 3  
  
261609 1.958083 ESTs, Weakly similar to S65657 alpha-1C-adrenergic receptor  
splice form 2 [H.sapiens]  
  
197525 1.9550731 flavin containing monooxygenase 5  
857640 1.9532398 collagen, type VI, alpha 2  
812143 1.9469226 fibronectin leucine rich transmembrane protein 3  
  
898222 1.9409805 Homo sapiens clone 24418 mRNA sequence  
  
418129 1.9385868 nuclear mitotic apparatus protein 1  
2021882 1.9213983 sodium channel, nonvoltage-gated 1, delta  
293819 1.919948 oxidoreductase UCPA  
203003 1.8989675 non-metastatic cells 4, protein expressed in  
  
705274 1.8986548 diacylglycerol kinase, delta (130kD)  
124922 1.8975916 KRAB-zinc finger protein SZF1-1  
2505310 1.8859615 calcium/calmodulin-dependent protein kinase I  
  
1492238 1.8795818 HSPC003 protein  
810358 1.8732889 acyl-Coenzyme A dehydrogenase, very long chain

- 164 -

2391494 1.8698169 ephrin-A4  
1592530 1.8669954 mammalian inositol hexakisphosphate kinase 2  
  
810671 1.8659671 hypothetical protein FLJ22269  
2108048 1.8648249 DNB5  
730036 1.8564918 Mad4 homolog  
810741 1.831265 GABA(A) receptor-associated protein  
45578 1.8274273 mitogen-activated protein kinase kinase 6  
647397 1.8263711 ESTs  
767495 1.8149096 GLI-Kruppel family member GLI3 (Greig  
cephalopolysyndactyly syndrome)  
32050 1.8117146 Homo sapiens mRNA; cDNA DKFZp586P1124 (from clone  
DKFZp586P1124)  
  
811848 1.8106436 hypothetical protein  
186301 1.8021505 Homo sapiens cDNA FLJ12924 fis, clone NT2RP2004709  
  
366526 1.7836259 ESTs, Weakly similar to ALU5\_HUMAN ALU SUBFAMILY SC  
SEQUENCE CONTAMINATION WARNING ENTRY  
[H.sapiens]  
  
813154 1.7815662 nuclear factor I/A  
85195 1.778695 growth arrest and DNA-damage-inducible, gamma  
  
2460159 1.7730834 tyrosine kinase, non-receptor, 1  
504959 1.7723846 Homo sapiens mRNA; cDNA DKFZp586G0321 (from clone  
DKFZp586G0321)  
  
742094 1.7684813 hypothetical protein FLJ20950  
256619 1.7658619 hydroxysteroid (17-beta) dehydrogenase 7  
726699 1.7594888 ESTs, Weakly similar to AAB47496 NG5 [H.sapiens]  
  
2017144 1.7514699 CGI-41 protein  
74070 1.7336683 endosulfine alpha  
1762111 1.7236006 natriuretic peptide receptor C/guanylate cyclase C  
(atrionatriuretic peptide receptor C)  
  
795750 1.7116475 Homo sapiens clone 25056 mRNA sequence  
  
2019750 1.7095011 SEC14 (S. cerevisiae)-like 2  
2325804 1.7039903 95 kDa retinoblastoma protein binding protein

- 165 -

279720 1.693594 ESTs, Moderately similar to A47582 B-cell growth factor precursor [H.sapiens]  
132857 1.6875688 Homo sapiens mRNA; cDNA DKFZp586N1323 (from clone DKFZp586N1323)  
1898619 1.684954 hypothetical protein MGC15737  
681992 1.6802347 Homo sapiens cDNA FLJ13384 fis, clone PLACE1001062, highly similar to Homo sapiens mRNA for lysine-ketoglutarate reductase/saccharopine dehydrogenase  
85450 1.6753438 acyl-Coenzyme A oxidase 2, branched chain  
51218 1.6748931 ESTs  
1925280 1.6563658 homologous to yeast nitrogen permease (candidate tumor suppressor)  
250883 1.6504115 ubiquitin-activating enzyme E1-like  
172783 1.6502064 hypothetical protein FLJ10390  
208387 1.6498608 KIAA1407 protein  
703964 1.6452497 inositol polyphosphate phosphatase-like 1  
298231 1.6428171 gamma-aminobutyric acid (GABA) B receptor, 1  
277848 1.6407233 ADP-ribosylation factor 6  
669359 1.6386236 Homo sapiens clone 24405 mRNA sequence  
321455 1.6327309 Homo sapiens, Similar to RIKEN cDNA 1110002C08 gene, clone MGC:9564 IMAGE:3872267, mRNA, complete cds  
796152 1.6295159 Homo sapiens cDNA FLJ11685 fis, clone HEMBA1004934  
502782 1.614818 RAN binding protein 3  
360778 1.6091782  
950574 1.606032 H3 histone, family 3B (H3.3B)  
788334 1.5816572 mitochondrial ribosomal protein L23  
782497 1.5767626 Homo sapiens, clone IMAGE:3010666, mRNA, partial cds  
1733262 1.5656438 BLu protein  
342181 1.5642436 B-cell CLL/lymphoma 2  
1632248 1.5590794 Homo sapiens cDNA FLJ14181 fis, clone NT2RP2004300  
1526826 1.5576997 homeo box B2  
145132 1.5520317 mannose-P-dolichol utilization defect 1  
183062 1.5400324 ubiquitin specific protease 21

- 166 -

823663 1.5337257 fragile X mental retardation, autosomal homolog 2  
190059 1.5258594 guanine nucleotide binding protein (G protein), gamma 7  
111721 1.5045285 insulin induced protein 2  
840882 1.5039692 nucleotide binding protein  
1902764 1.5037016 6-phosphofructo-2-kinase/fructose-2,6-biphosphatase 3  
414999 1.5021506 ets variant gene 4 (E1A enhancer-binding protein, E1AF)  
1636523 1.4948923 glutathione S-transferase subunit 13 homolog  
594683 1.4862053 ESTs  
1916575 1.4825298 BCL2-interacting killer (apoptosis-inducing)  
1456937 1.4730662 oviductal glycoprotein 1, 120kD (mucin 9, oviductin)  
25274 1.464238 g20 protein  
795288 1.4625618 ubiquitin specific protease 4 (proto-oncogene)  
1569418 1.4599089 Homo sapiens cDNA FLJ11385 fis, clone HEMBA1000520  
753700 1.4522717 Ras-related GTP-binding protein  
1691237 1.4516493 ESTs  
741891 1.450795 RAB2, member RAS oncogene family-like  
1493383 1.4468163 Homo sapiens mRNA; cDNA DKFZp434H2418 (from clone  
DKFZp434H2418)  
1675273 1.445145 RAR-related orphan receptor C  
293916 1.4265341 FKBP-associated protein  
39600 1.4224626 adenylate kinase 5  
1573087 1.4200341 KIAA0592 protein  
839796 1.4114662 candidate tumor suppressor p33 ING1 homolog  
296123 1.4057088 Homo sapiens PRO1851 mRNA, complete cds  
727263 1.4046408 heterogeneous nuclear ribonucleoprotein U (scaffold  
attachment factor A)  
265045 1.4041143 Homo sapiens mRNA; cDNA DKFZp586B0918 (from clone  
DKFZp586B0918)  
869450 1.4034983 ribosomal protein L11  
714472 1.4006155 KIAA0397 gene product

- 167 -

825296 1.3963737 low density lipoprotein receptor defect C complementing  
753301 1.3962897 carcinoembryonic antigen-related cell adhesion molecule 1  
(biliary glycoprotein)  
700527 1.394937 glutaredoxin (thioltransferase)  
811565 1.3909974 KIAA1694 protein  
1630990 1.3843882 ribosomal protein L29  
70749 1.3810753 Homo sapiens cDNA: FLJ21874 fis, clone HEP02488  
  
814826 1.3785178 ESTs  
175103 1.376581 cadherin, EGF LAG seven-pass G-type receptor 2, flamingo  
(Drosophila) homolog  
215000 1.3677098 vasoactive intestinal peptide receptor 1  
810331 1.3656069 quiescin Q6  
1681421 1.3645145 EGF-like-domain, multiple 3  
502198 1.3619124 protein phosphatase 1, regulatory (inhibitor) subunit 5  
  
825365 1.361275 hypothetical protein FLJ21919  
1557047 1.3582677 thrombospondin 3  
2067500 1.3579293 Z-band alternatively spliced PDZ-motif  
490615 1.3566637 tubulin, gamma 2  
743880 1.3528093 KIAA0263 gene product  
812099 1.3444147 RNA binding motif protein 5  
868652 1.3432331 complement component 4B  
120138 1.3384804 J domain containing protein 1  
156363 1.3358314 hypothetical protein FLJ12934  
813584 1.3327194 p53 regulated PA26 nuclear protein  
726703 1.3247883 Homo sapiens clone 23736 mRNA sequence  
  
809507 1.3225829 hypothetical protein FLJ20568  
2018423 1.3212314 death-associated protein kinase 2  
292806 1.3201283 chromosome segregation 1 (yeast homolog)-like  
  
741977 1.3200635 B-factor, properdin  
610326 -4.5776738  
322494 -3.6341126 core-binding factor, beta subunit  
824962 -3.6328144 karyopherin alpha 2 (RAG cohort 1, importin alpha 1)  
  
825659 -3.6247426 N-myc downstream regulated  
796694 -3.5038212 baculoviral IAP repeat-containing 5 (survivin)

- 168 -

149355 -3.4959791 translocating chain-associating membrane protein  
814270 -3.3246381 polymyositis/scleroderma autoantigen 1 (75kD)  
884425 -3.3113482 chaperonin containing TCP1, subunit 5 (epsilon)  
1874367 -3.2390912 small inducible cytokine subfamily A (Cys-Cys), member 20  
742707 -3.2093502 ESTs, Weakly similar to MUC2\_HUMAN MUCIN 2  
PRECURSOR [H.sapiens]  
321354 -3.1386611 mitochondrial ribosomal protein L15  
280375 -3.110486 PRO2000 protein  
2018154 -3.0906181 electron-transfer-flavoprotein, beta polypeptide  
814792 -3.0561472 ubiquitin specific protease 10  
826070 -3.0517117 KIAA0948 protein  
624867 -2.9688171 hypothetical protein FLJ20186  
202035 -2.9361687 uncharacterized hypothalamus protein HT010  
1476053 -2.9129535 RAD51 (S. cerevisiae) homolog (E coli RecA homolog)  
194318 -2.8913426 hypothetical protein MGC5585  
769921 -2.8415969 ubiquitin carrier protein E2-C  
1732922 -2.8216808 Homo sapiens mRNA; cDNA DKFZp762H106 (from clone  
DKFZp762H106)  
1536236 -2.7995652 hypothetical protein FLJ13154  
200402 -2.7793137 hypothetical protein dJ616B8.3  
815556 -2.7693 hypothetical protein FLJ10430  
1055607 -2.7538365 SUMO-1 activating enzyme subunit 1  
1466621 -2.7448687 ATPase, Ca++ transporting, type 2C, member 1  
770675 -2.6929924 Homo sapiens cDNA: FLJ21323 fis, clone COL02374  
79520 -2.6909833 RAB2, member RAS oncogene family  
950355 -2.6845343 ESTs, Weakly similar to S13495 pregnancy zone protein  
[H.sapiens]  
79710 -2.6813265 KIAA0174 gene product  
2011138 -2.6569485 KIAA1036 protein  
898333 -2.6509707 ESTs  
878846 -2.6361347 brain protein I3  
842818 -2.63282 lysyl-tRNA synthetase  
1553065 -2.627227 hypothetical protein FLJ14993

sd-71385

1002301-1002301



- 169 -

136722 -2.6269431 ATPase, Na<sup>+</sup>/K<sup>+</sup> transporting, beta 3 polypeptide

308633 -2.6261051 hypothetical protein FLJ10339

854581 -2.6159344 transcription factor 4

951241 -2.5987959 clone HQ0310 PRO0310p1

292936 -2.5940369 hypothetical protein FLJ10468

781047 -2.569753 budding uninhibited by benzimidazoles 1 (yeast homolog)

823907 -2.5587562 hypothetical protein FLJ10511

377368 -2.5495557 cell death regulator aven

292388 -2.5448438

122241 -2.539248 proteasome (prosome, macropain) subunit, beta type, 2

731223 -2.5364769 proteasome (prosome, macropain) subunit, beta type, 2

1660666 -2.534682 carbonic anhydrase VB, mitochondrial

825470 -2.530053 topoisomerase (DNA) II alpha (170kD)

1472719 -2.526978 SMT3 (suppressor of mif two 3, yeast) homolog 1

869375 -2.5249164 isocitrate dehydrogenase 2 (NADP<sup>+</sup>), mitochondrial

302292 -2.5097213 exostoses (multiple) 2

1175007 -2.4998247 tumor necrosis factor receptor superfamily, member 10d,  
decoy with truncated death domain

1422338 -2.488461 ribonucleotide reductase M2 polypeptide

2014034 -2.4843808 methylene tetrahydrofolate dehydrogenase (NAD<sup>+</sup>  
dependent), methenyltetrahydrofolate cyclohydrolase

50884 -2.4377483 dual-specificity tyrosine-(Y)-phosphorylation regulated kinase 2

1474424 -2.4345879 Homo sapiens cDNA FLJ12758 fis, clone NT2RP2001328

753285 -2.4324065 glycogenin

700792 -2.4311609 cyclin-dependent kinase inhibitor 3 (CDK2-associated dual  
specificity phosphatase)

259950 -2.4310983 hypothetical protein FLJ14991

610326 -2.4214624

284004 -2.4185183 ESTs

756595 -2.3963227 S100 calcium-binding protein A10 (annexin II ligand, calpactin  
I, light polypeptide (p11))

- 170 -

128711 -2.3891503 anillin (Drosophila Scraps homolog), actin binding protein

1582738 -2.3882015 uncharacterized bone marrow protein BM040

1915867 -2.3829702 Homo sapiens cDNA: FLJ21086 fis, clone CAS03272

754582 -2.3765311 ecotropic viral integration site 2A

462961 -2.3685047 dihydrofolate reductase

705064 -2.3682586 transforming, acidic coiled-coil containing protein 3

1553696 -2.3605995 vanin 1

129294 -2.3507956 ESTs

809588 -2.3342142 gamma-glutamyl hydrolase (conjugase, folylpolyglutaminyl hydrolase)

1909526 -2.333814 ESTs, Weakly similar to 2004399A chromosomal protein [H.sapiens]

815771 -2.3166739 chromosome 7 open reading frame 2

703707 -2.2865473 aspartate beta-hydroxylase

2017415 -2.2832169 centromere protein A (17kD)

1650942 -2.2797717 Human proteinase activated receptor-2 mRNA, 3'UTR

645079 -2.2761946 3'(2'), 5'-bisphosphate nucleotidase 1

133213 -2.2753496 fucosyltransferase 4 (alpha (1,3) fucosyltransferase, myeloid-specific)

788566 -2.2752004 Purkinje cell protein 4

2009574 -2.2646007 T brachyury (mouse) homolog

1897302 -2.2563379 serine/threonine kinase 17b (apoptosis-inducing)

78869 -2.2539078 cell membrane glycoprotein, 110000M(r) (surface antigen)

590759 -2.2513499 sterol-C4-methyl oxidase-like

785368 -2.2489394 PDZ-binding kinase; T-cell originated protein kinase

1714689 -2.2489064 ESTs

795197 -2.2481741 ubiquitin-conjugating enzyme E2E 2 (homologous to yeast UBC4/5)

781472 -2.2390098 Tax1 (human T-cell leukemia virus type I) binding protein 1

1469966 -2.2375228 delta (Drosophila)-like 3

1587847 -2.2354975 minichromosome maintenance deficient (mis5, S. pombe) 6

209066 -2.2325695

1035796 -2.2308926 ESTs

sd-71385

"PATENT" ST032001

233464	-2.2280935	epithelial V-like antigen 1
753320	-2.2104141	hypothetical protein FLJ20533
486626	-2.2095389	ESTs
1505038	-2.2060272	hypothetical protein FLJ20171
503851	-2.1992378	nuclear receptor co-repressor/HDAC3 complex subunit
503671	-2.1939286	Homo sapiens cDNA FLJ14368 fis, clone HEMBA1001122
610326	-2.1885269	
712139	-2.1839319	ADP-ribosylation factor-like 7
471196	-2.1797647	integral membrane protein 3
753215	-2.1781545	guanine nucleotide binding protein (G protein), alpha inhibiting activity polypeptide 1
183556	-2.1734918	gap junction protein, alpha 4, 37kD (connexin 37)
809557	-2.1667294	minichromosome maintenance deficient (S. cerevisiae) 3
610326	-2.1633594	hypothetical protein MGC12992
1600239	-2.1602202	HSPC037 protein
1049291	-2.1593791	olfactory receptor, family 7, subfamily E, member 47 pseudogene
839682	-2.1475459	ubiquitin-conjugating enzyme E2N (homologous to yeast UBC13)
950897	-2.1419057	Homo sapiens mRNA; cDNA DKFZp586G1922 (from clone DKFZp586G1922)
712577	-2.1411578	holocytochrome c synthase (cytochrome c heme-lyase)
824524	-2.1400285	UDP-galactose transporter related
241348	-2.1301572	prenylcysteine lyase
823930	-2.1259223	actin related protein 2/3 complex, subunit 1A (41 kD)
150314	-2.1206632	lysophospholipase I
795498	-2.1190094	putative transmembrane protein
82710	-2.1184513	glutathione S-transferase A2
1869201	-2.1135906	hypothetical protein MGC2745
1694526	-2.1126121	hypothetical protein FLJ11029
823598	-2.1072985	proteasome (prosome, macropain) 26S subunit, non-ATPase, 12
214162	-2.1065234	metallothionein 1H
811590	-2.1050783	hypothetical protein FLJ11100
796904	-2.1050163	pleiomorphic adenoma gene-like 1

- 172 -

1517595 -2.0965219 KIAA0175 gene product  
282428 -2.0954489 ESTs, Weakly similar to A46010 X-linked retinopathy protein  
[H.sapiens]  
1883219 -2.0949091 ESTs  
746229 -2.0906355 mitogen-activated protein kinase kinase kinase kinase 4  
  
610326 -2.0896786 hypothetical protein MGC12992  
1632225 -2.0828168 ESTs  
743810 -2.0796998 hypothetical protein MGC2577  
1416055 -2.0725881 extra spindle poles, *S. cerevisiae*, homolog of  
  
1626297 -2.0656173 hypothetical protein FLJ20509  
852829 -2.0608067 karyopherin alpha 3 (importin alpha 4)  
840511 -2.0554449 vimentin  
278504 -2.0519206 neurochondrin  
1534700 -2.0506628 KIAA0830 protein  
812276 -2.0458249 synuclein, alpha (non A4 component of amyloid precursor)  
  
306841 -2.0449642 T cell receptor beta locus  
3172883 -2.0437672 ESTs, Weakly similar to 1709359A dopamine D4 receptor  
[H.sapiens]  
624627 -2.0409259 ribonucleotide reductase M2 polypeptide  
486179 -2.0380168 Homo sapiens cDNA FLJ10205 fis, clone HEMBA1004954  
  
376316 -2.0218598 wee1+ (*S. pombe*) homolog  
37671 -2.0190035 hypothetical protein FLJ21610  
629944 -2.0166696 myosin VB  
713158 -2.0133834 ESTs  
504308 -2.0067977 hypothetical protein FLJ10540  
1605426 -1.9957389 hypothetical protein FLJ13352  
810711 -1.9920785 stearyl-CoA desaturase (delta-9-desaturase)  
  
263790 -1.9904129 isocitrate dehydrogenase 3 (NAD+) alpha  
549349 -1.9893582 hypothetical protein PRO1068  
109221 -1.9879196 KIAA0286 protein  
51773 -1.9875155 hypothetical protein MGC3077

Example X: Crossvalidation

- 173 -

The following table shows the results of crossvalidation analysis (as described in Example I above) of various pair-wise comparisons. The two members of the pairwise comparison are indicated as "Class 1" and "Class 2" with the number of samples analyzed indicated under "N". The number of genes used to predict is indicated, along with the accuracy (in percent) for each combination. "FP" and "FN" refer to "false positive" and "false negative", respectively, as incorrectly identified for each of the two classes.

Class 1 Name	N	Class 2 Name	N	Genes	Accuracy	Class 1 FP	Class 1 FN	Class 2 FP	Class 2 FN
Normal	28	Abnormal	57	850	99	0	1	1	0
Normal	28	ADH	7	600	100	0	0	0	0
Normal	28	DCIS	28	1300	100	0	0	0	0
ADH	7	DCIS	28	350	97	0	1	1	0
ADH	7	DCIS	28	10	97	0	1	1	0
N+ADH	36	DCIS+IDC	49	400	98	1	1	1	1
DCIS-I	7	DCIS-III	9	300	100	0	0	0	0
IDC-I	4	IDC-III	8	300	100	0	0	0	0

#### References:

DeRisi, J., et al., *Use of a cDNA microarray to analyse gene expression patterns in human cancer*, Nature Genetics, (1996) 14:457-460.

Hedenfalk, I., et al., *Gene-Expression Profiles In Heredity Breast Cancer*, The New England Journal of Medicine, (February 22, 2001) 344:8:539-548.

Golub, T. R., et al., *Molecular Classification of Cancer: Class Discovery and Class Prediction by Gene Expression Monitoring*, Science, (October 15, 1999) 286:531-537.

- 174 -

Perou, Charles M., et al., *Molecular portraits of human breast tumours*, Nature, (August 17, 2000) 406:747-752.

Garber, Mitchell E., et al., *Diversity of gene expression in adenocarcinoma of the lung*, Proc. Natl. Acad. Sci. USA, (November 20, 2001) 98:24:13784-13789.

Perou, Charles M., et al., *Distinctive gene expression patterns in human mammary epithelial cells and breast cancers*, Proc. Natl. Acad. Sci. USA, (August 1999) 96:9212-9217.

Sgrio, Dennis C., et al., *In Vivo Gene Expression Profile Analysis of Human Breast Cancer Progression*, Cancer Research, (November 15, 1999) 59:5656-5661.

Sorlie, Therese, et al., *Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications*, Proc. Natl. Acad. Sci., (September 11, 2001) 98:19:10869-10874.

Alizadeh, Ash A., et al., *Distinct types of diffuse large B-cell lymphoma identified by gene expression profiling*, Nature, (February 3, 2000) 403:503-511.

Bittner, M., et al., *Molecular classification of cutaneous malignant melanoma by gene expression profiling*, Nature (August 3, 2000) 406:536-540.

West, Mike, et al., *Predicting the clinical status of human breast cancer by using gene expression profiles*, Proc. Natl. Acad. Sci., (September 25, 2001) 98:20:11462-11467.

- 175 -

All references cited herein, including patents, patent applications, and publications, are hereby incorporated by reference in their entireties, whether previously specifically incorporated or not.

Having now fully described this invention, it will be appreciated by those skilled in the art that the same can be performed within a wide range of equivalent parameters, concentrations, and conditions without departing from the spirit and scope of the invention and without undue experimentation.

While this invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications. This application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure as come within known or customary practice within the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth.

10026018 133101  
FOOTNOTES